



2025

Brunswick County EMS
Protocols, Procedures, Policies and References

Be Kind, Be Determined, Be United

North Carolina Medical Board Approved Medications for Credentialed EMS Personnel

EMS personnel at any level who administer medications must do so with medical oversight. Personnel must complete appropriate medical education. All EMS System and SCTP protocols, policies and procedures must be reviewed and approved by the Medical Director of the Office of EMS

All items highlighted in “red” are required by NCCEP in all systems with EMS personnel credentialed at the specified level. Specialty Care (SCTP) required items are not listed here, as they can be found on the Specialized Ambulance Protocol Summary (SAPS) form.

Medications	EMR	EMT	AEMT	MEDIC
ACE inhibitors				X
Acetaminophen	X	X	X ¹⁵	X
Adenosine				X
Aminophylline				X
Amiodarone				X
Anti-arrhythmic				X ¹²
Antibiotics			X	X
Anti-emetic preparations			X	X
Antipsychotic (Typical and Atypical)		X ¹⁹	X ²⁰	X ²⁰
Antivirals			X	X
Aspirin	X	X	X	X
Atropine	X ⁴	X ⁴	X ⁴	X
Barbiturates				X
Benzodiazepine preparations				X ¹⁴
Beta agonist preparations		X ²	X	X
Beta blockers				X ¹³
Bretylium				X
C1 Esterase-Inhibitors				X
Calcium channel blockers				X ¹³
Calcium chloride/gluconate				X
Calcium Paste		X	X	X
Charcoal		X	X	X
Clonidine				X
Clopidogrel				X
CroFab (Crotalidae Polyvalent Immune Fab)				X ⁸
Crystalloid solutions			X	X
Cyanide poisoning antidote kit				X
Digoxin				X
Diphenhydramine	X ³	X ³	X	X
Diuretics				X
Dobutamine				X
Dopamine				X
Droperidol				X
Epinephrine	X ¹	X ¹	X	X
Etomidate				X
Flumazenil				X
Glucagon		X	X	X
Glucose, oral	X	X	X	X
Glucose solutions			X	X
Haloperidol			X	X
Heparin (unfractionated and low molecular weight)				X
Histamine 2 blockers			X	X

Medications	EMR	EMT	AEMT	MEDIC
Hydroxocobalamin				X
Immunizations		X ²¹	X ⁶	X ⁶
Insulin				X
Ipratropium			X	X
Isoproterenol				X
Ketamine				X ⁷
Levetiracetam				X
Lidocaine			X ¹⁸	X
Magnesium sulfate			X	X
Mannitol				X
Methylene blue				X
Milrinone				X
Monoclonal Antibodies			X	X
N-acetylcysteine				X
Narcotic analgesics				X
Narcotic antagonists	X ^{9,10}	X ^{9,10}	X	X
Nasal spray decongestant		X	X	X
Nesiritide				X
Nitroglycerin		X ²	X	X
Nitroprusside sodium				X
Nitrous oxide		X	X	X
Non-prescription medications		X	X	X
Non-steroidal anti-inflammatory		X	X ¹⁵	X
Norepinephrine				X
Octreotide				X
Oxygen	X ⁵	X ⁵	X ⁵	X ⁵
Oxytocin				X
Paralytic agents				X ¹⁷
Phenothiazine preparations				X
Phenylephrine				X
Phenytoin preparations				X
Plasma protein fraction				X
Platelet g-II/IIIa inhibitors				X
Potassium chloride				X
Pralidoxime	X ⁴	X ⁴	X ⁴	X
Procainamide				X
Procaine				X
Proparacaine				X
Propofol				X ⁸
Proton pump inhibitors				X
Sodium bicarbonate				X
Steroid preparations			X	X
Thiamine			X	X
Thrombolytic agents				X
Topical hemostatic agents	X	X	X	X
Total Parenteral Nutrition				X
Tranexamic Acid (TXA)			X	X ¹¹
Tuberculosis skin test			X ⁶	X ⁶
Valproic Acid				X
Vasopressin			X	X
Vasopressor				X ¹⁶
Whole blood and components				X
Ziprasidone				X

- ¹ EMR and EMT use of epinephrine is limited to the treatment of anaphylaxis and may be administered only by auto injector, unless approved by EMS System Medical Director and OEMS.
- ² EMT use of beta-agonists and nitroglycerine is limited to patients who currently are prescribed the medication unless approved by the EMS System Medical Director and OEMS as part of the expanded scope. EMTs may administer these medications from EMS supplies.
- ³ EMR/EMT administration of diphenhydramine is limited to the oral route.
- ⁴ As a component of preparedness for domestic terrorism, EMS personnel, public safety officers, and other first responders recognized by the EMS system, may carry, self-administer, or administer to a patient atropine and/or pralidoxime, based on written protocols and medical direction. All personnel except for Paramedics must administer these medications by an auto injector.
- ⁵ Administration of oxygen does not require medical direction.
- ⁶ Administration of immunizations and TB skin tests are not limited to public health initiatives.
- ⁷ Ketamine use is restricted to programs that have been approved by the OEMS State Medical Director. It can be used as an induction or post intubation sedation agent in approved DAI programs. Use outside of DAI programs must meet all the requirements outlined in Medical Policy 2 'Ketamine Program Requirements'.
- ⁸ Propofol use is restricted to programs that have been approved by the OEMS State Medical Director. EMS Systems and SCTP's must submit a policy and education plan to the OEMS prior to approval. EMS personnel cannot initiate Propofol, it can only be used for interfacility transport where infusion has already been started at transferring facility. **EMS units cannot stock Propofol or CroFab. This medication must be provided by the transferring hospital.**
- ⁹ FR, EMR, and EMT administration of Naloxone is limited to the intra-nasal (IN), intra-muscular (IM), and auto-injector routes.
- ¹⁰ First Responders (FR) who administer Naloxone must do so under the medical oversight of the County EMS Medical Director, following protocols and procedures approved by the OEMS State Medical Director. FR administration must be monitored by the EMS Systems peer review program.
- ¹¹ For an EMS System to use Tranexamic Acid (TXA), they must submit for approval by the OEMS State Medical Director a signed letter from any Trauma Centers that would be the recipient of the patient that the destination Trauma Center agrees with its use and will give the 2nd required dose of Tranexamic Acid (TXA).
- ¹² All Paramedic systems must carry some form of anti-arrhythmic agent. This must either be amiodarone, lidocaine, **or** procainamide.
- ¹³ Paramedic systems must carry either a calcium channel blocker **or** beta-blocker.
- ¹⁴ All Paramedic systems must carry some form of injectable benzodiazepine.
- ¹⁵ AEMT systems must carry either acetaminophen **or** a non-steroidal anti-inflammatory.
- ¹⁶ All Paramedic systems must carry an approved vasopressor. This must either be dobutamine, dopamine, epinephrine, norepinephrine, phenylephrine, **or** vasopressin.
- ¹⁷ Paralytic agent use is restricted to Drug Assisted Intubation (DAI) programs approved by the OEMS State Medical Director. They require the submission of; signed NCCEP DAI policy by local medical director, unaltered NCCEP DAI protocols, training documentation, and process for peer review of cases. All DAI must have an EMS Airway Evaluation form completed and signed by local medical director in accordance with the NCCEP DAI policy. Systems utilizing must submit monthly airway forms and cases to the OEMS for review.
- ¹⁸ AEMT administration of Lidocaine is allowed for analgesic use only.
- ¹⁹ EMTs may only administer antipsychotic medications orally and if the patient has a current prescription.
- ²⁰ Long-acting antipsychotics may only be used in pilot programs that are approved by the OEMS State Medical Director in conjunction with the State Mental Health Medical Director.
- ²¹ EMTs may administer immunizations in conjunction with public health initiatives.

North Carolina Medical Board Approved Skills for Credentialed EMS Personnel

All items highlighted in “red” are required by NCCEP in all systems with EMS personnel credentialed at the specified level. Specialty Care (SCTP) required items are not listed here, as they can be found on the Specialized Ambulance Protocol Summary (SAPS) form.

Skills	EMR	EMT	AEMT	MEDIC
12-Lead ECG Acquisition & Transmission		X	X	X
12-Lead ECG Interpretation				X
15-Lead ECG Acquisition				X
Airway Adjuncts (NPA/OPA)	X	X	X	X
Arterial Access - Blood Draw				X
Arterial Line maintenance				X
Blind Insertion Airway Device (BIAD)	X ¹	X ¹	X	X
Capnography (Waveform)	X ⁶	X ⁶	X ⁶	X ⁶
Carbon Monoxide Measurement (non-invasive)	X	X	X	X
Cardiac Monitoring		X ⁴	X ⁴	X
Cardiac Pacing				X
Cardiopulmonary Resuscitation	X	X	X	X
Cardioversion				X
Carotid Massage				X
Central Venous Pressure Line Maintenance				X
Chest Compression-External Device	X	X	X	X
Chest Decompression-Needle			X ¹¹	X
Chest Tube Maintenance				X
Childbirth	X	X	X	X
Cricothyrotomy-Needle				X
Cricothyrotomy-Surgical				X ⁵
Decontamination	X	X	X	X
Defibrillation-Automated	X	X	X	X
Defibrillation-Manual			X ¹²	X
Direct Laryngoscopy			X	X
Drug Assisted Intubation (DAI)				X ^{5,6}
Endotracheal Tube Introducer			X	X
Epidural Catheter Maintenance				X
Foreign Body Airway Obstruction	X	X	X	X
Gastric Intubation		X ³	X ³	X
Glucose Measurement	X	X	X	X
Hemostatic Agent	X	X	X	X
Injections – Subcutaneous and Intramuscular		X ²	X	X
Intra-Ventricular Catheter Maintenance				X
Intubation - Nasotracheal			X	X
Intubation - Orotracheal			X ⁶	X ^{6,7}
Intubation Confirmation - Capnometry (color)			X	X
Medication Administration	X ²	X ²	X ²	X ²
Nebulizer Inhalation Therapy		X	X	X
Non-Invasive Positive Pressure Ventilation	X ⁹	X	X	X
Orthostatic Blood Pressure	X	X	X	X
Oxygen Administration	X	X	X	X
Patient Assessment	X	X	X	X
Pulse Oximetry	X	X	X	X

Skills	EMR	EMT	AEMT	MEDIC
Reperfusion Checklist	X	X	X	X
Respirator Operation		X	X	X
Restraints		X	X	X
Specimen Collection		X	X	X
Spinal Motion Restriction	X	X	X	X
Splinting	X	X	X	X
Stroke Screen	X	X	X	X
Suction-Basic	X	X	X	X
Suction-Advanced			X ¹⁰	X ¹⁰
Swan-Ganz Catheter maintenance				X
Taser Probe Removal	X	X	X	X
Temperature Measurement	X	X	X	X
Tourniquet Application	X	X	X	X
Tracheostomy Tube Change			X	X
Urinary Catheterization				X
Venous Access-Blood Draw			X	X
Venous Access-Existing catheters				X
Venous Access-Femoral Line				X
Venous Access-Intraosseous			X	X
Venous Access-Peripheral			X	X
Ventilator Operation		X ⁸	X ⁸	X
Wound Care	X	X	X	X

¹ EMRs and EMTs using blind insertion airway devices must be functioning in EMS systems with medical direction and written treatment protocols.

² EMS personnel educated in approved programs, credentialed by the OEMS, and functioning under physician medical oversight may perform acts and administer intravenous fluids and medications as allowed by the North Carolina Medical Board pursuant to G.S. 143-514. The administration of oxygen does not require medical direction.

³ Gastric tube insertion may be performed only when utilized in conjunction with a blind insertion airway device.

⁴ EMT and AEMT may use the cardiac monitor for vital sign monitoring and EKG transmission.

⁵ Systems performing drug assisted intubation (DAI) must have the ability to perform surgical cricothyrotomy. Commercial cricothyrotomy or tracheostomy kits that create an airway comparable to a surgical cricothyrotomy are acceptable.

⁶ End-tidal (EtCO₂) monitoring is mandatory following placement of an endotracheal tube. EtCO₂ monitoring is mandatory following placement of a BIAD once available on scene.

⁷ Pediatric intubation is an optional skill/procedure.

⁸ Ventilator patients may be transported by EMT/AEMT when all the following conditions are met:

- Patient is receiving home (or skilled nursing) ventilator therapy.
- The ventilator is portable and can continue to ventilate the patient during transport.
- The patient is accompanied by a non-EMS adult (from either the home or facility) who is knowledgeable, capable, and willing to maintain the ventilator during the EMS transport.
- While in transit, the patient is monitored using pulse oximetry.

⁹ Bag Valve Mask ONLY

¹⁰ For a patient currently being assisted by an airway adjunct such as a naso-tracheal tube, endotracheal tube, BIAD, tracheostomy tube or a cricothyrotomy tube.

¹¹ Use of needle chest decompression at the AEMT level is for traumatic arrest only.

¹² AEMT use of manual defibrillation is for pulseless arrest only.

-EMD personnel are responsible for:

- 1) Pre-arrival instructions to callers
- 2) Determining and dispatching appropriate EMS resources
- 3) All EMD skills must be performed in EMS systems with medical oversight and written EMS protocols.

Introduction

The following medical treatment protocols are developed for North Carolina EMS agencies. The process has evolved since 2007 and continues with input from Medical Directors, EMS Administration, North Carolina Chapter of Emergency Physicians Protocol Committee, North Carolina Office of EMS, EMS field personnel and the public at large through on-line surveys, public meetings across North Carolina and direct communication with stakeholders. Updates expand on previous versions and continue to incorporate evidence-based guidelines, expert opinion and historically proven practices meant to ensure that citizens and visitors of North Carolina will continue to be provided the highest quality pre-hospital patient care available. The North Carolina Chapter of Emergency Physicians develops and provides final approval.

The purpose of the protocol section is to provide treatment protocols outlining permissible and appropriate assessment, delivery of care, reassessment and procedures which may be rendered by pre-hospital providers. The protocols also outline which medical situations require direct voice communication with medical control. In general treatment protocols are specific orders which may and should be initiated prior to contact with Medical Control.

Please note the medical protocols are divided into three (3) to four (4) sections. The upper section includes three (3) boxes (History, Signs and Symptoms and Differential) which serves as a guide to assist in obtaining pertinent patient information and exam findings as well as considering multiple potential causes of the patients complaint. It is not expected that every historical element or sign / symptom be recorded for every patient. It is expected that those elements pertinent to your patient encounter will be included in the patient evaluation.

The algorithm section describes the essentials of patient care. Virtually every patient should receive the care outlined in this section, usually in the order described. However each medical emergency must be dealt with individually and appropriate care determined accordingly. Professional judgment is mandatory in determining treatment modalities within the parameters of these protocols. Circumstances will arise where treatment may move ahead in the algorithm, move outside to another protocol and then re-enter later. While protocols are written based on body systems and primary complaints the patient should be treated as a whole and therefore the protocols should be considered as a whole in providing care.

Professional judgment hierarchy:

The pre-hospital provider may determine that no specific treatment is needed;

Or

The pre-hospital provider may follow the appropriate treatment protocols and then consult Medical Control;

Or

The pre-hospital provider may consult Medical Control before initiating any specific treatment.

Some protocols will encompass two (2) pages. Protocols which exist in a single page format may have page 2 added by the local medical director. The PEARLS section will either be located at the bottom of page 1 (single page protocol) or page 2 (double page protocol). The PEARLS section provides points regarding the main protocol based on evidence to date, common medical knowledge and expert medical opinion.

Information boxes highlighted in purple. These areas are editable at the local level. They will mainly involve specific medications and dosages utilized by the local EMS agency. Page 2 will have a large section highlighted in purple where the local Medical Director may edit as they see fit to provide expanded points and treatment not otherwise specified in the algorithm. If the box is not to be utilized – add “***This Space Left Blank Intentionally.***”

Finally these medical treatment protocols are established to ensure safe, efficient and effective interventions to relieve pain and suffering and improve patient outcomes without inflicting harm. They also serve to ensure a structure of accountability for Medical Directors, EMS agencies, pre-hospital providers and facilities to provide continual performance improvement. A recent report of the Institute of Medicine calls for the development of standardized, evidence-based pre-hospital care protocols for the triage, treatment and transport of patients. These protocols establish expectations of pre-hospital care in North Carolina.

Key to Protocol Utilization

History

- Important history items
- Circumstances of event
- SAMPLE
- Time of onset
- Duration

Signs and Symptoms

- Important Signs and Symptoms specific to each protocol

Differential

- A list of other disease or injury which should be considered

Black Box

Highlights Important Information



Universal Patient Care Protocol
Assumed all protocols utilize and will not appear on individual protocols

Red Box

Highlights Critical Information

May direct to another protocol



Signals protocol within a protocol

Information box

Indicates Entry / Exit from / to another protocol(s)



Decision Point
 Darker outline to highlight



Highlights medication after Contact Medical Control
 May be added by Local Medical Director

Purple Shading of Information Box

Indicates items changeable at local agency level, including medications / dosages on NCMB formulary
 Local Medical Director may add / change at his / her discretion
 Local medical director may add page 2 to any protocol where none exists for additional comments

Algorithm Legend

	Emergency Medical Responder
B	Emergency Medical Technician
A	Advanced Emergency Medical Technician
P	Paramedic
	Notify Destination or Contact Medical Control

Pearls

- Important information specific to each protocol will appear here.
- Will usually appear on page.
- Important exam items listed here specific to protocol.

2025 Table of Contents

Treatment Protocols

Protocol introduction and Key to Utilization from NCCEP EMS Committee

*Denotes protocol unique to Brunswick County EMS

Universal Protocols (UP)	
Universal Patient Care	UP1
Triage	UP2
Abdominal Pain / Vomiting and Diarrhea	UP3
Altered Mental Status	UP4
Back Pain	UP5
IV / IO Access	UP6
Dental Emergencies	UP7
Emergencies Involving Indwelling Central Lines	UP8
Epistaxis	UP9
Fever / Infection Control	UP10
Pain Control*	UP11*
Police Custody	UP12
Seizure*	UP13*
Suspected Stroke	UP14
Suspected Sepsis	UP15
Syncope	UP16
Behavioral Health Crisis	UP17
Behavioral Agitation / Sedation Guide	UP18
Behavioral Hyperactive Delirium Syndrome / Violent	UP19
Sickle Cell Crisis	UP 20

Airway Respiratory Section (AR)	
Adult Airway*	AR1*
Adult Failed Airway	AR2
Airway, Drug Assisted*	AR3*
Adult COPD /Asthma*	AR4*
Pediatric Airway*	AR5*
Pediatric Failed Airway	AR6
Pediatric Asthma Respiratory Distress*	AR7*
Post-Intubation / BIAD Management*	AR8*
Ventilator Emergencies	AR9
Tracheostomy Tube Emergencies	AR10

Adult Cardiac Section (AC)	
Asystole / Pulseless Electrical Activity	AC1
Bradycardia; Pulse Present	AC2
Cardiac Arrest; Adult	AC3
Chest Pain; Cardiac and STEMI	AC4
CHF / Pulmonary Edema	AC5
Adult Tachycardia Narrow Complex (≤ 0.11 sec)	AC6
Adult Monomorphic Tachycardia Wide Complex (≥ 0.12 sec)	AC7
Adult Polymorphic Tachycardia Wide Complex, Torsades	AC8
Ventricular Fibrillation; Pulseless Ventricular Tachycardia	AC9
Post Resuscitation	AC10*
Team Focused CPR	AC11
On Scene Resuscitation / Termination of CPR	AC12
LVAD, RVAD and Bi-VAD	AC14
Total Artificial Heart	AC15
Wearable Cardioverter Defibrillator Vest	AC16

Adult Medical Section (AM)	
Allergic Reaction / Anaphylaxis*	AM1*
Diabetic; Adult	AM2
Dialysis / Renal Failure	AM3
Hypertension	AM4
Hypotension / Shock	AM5

Adult Obstetrical Section (AO)	
Childbirth / Labor	AO1
Newly Born	AO2
Obstetrical Emergencies	AO3

Trauma and Burn Section (TB)	
Blast Injury / Incident	TB1
Chemical and Electrical Burn	TB2
Crush Syndrome Trauma	TB3
Extremity Trauma	TB4
Head Trauma	TB5
Multiple Trauma	TB6
Radiation Incident	TB7
Spinal Motion Restriction	TB8
Thermal Burn	TB9
Traumatic Arrest	TB10

Pediatric Cardiac Section (PC)	
Pediatric Asystole / PEA	PC1
Pediatric Bradycardia	PC2
Pediatric CHF / Pulmonary Edema	PC3
Pediatric Cardiac Arrest	PC4
Pediatric Tachycardia Narrow Complex (≤ 0.09 sec)	PC5
Pediatric Tachycardia Wide Complex (>0.09 sec)	PC6
Pediatric Ventricular Fibrillation / Pulseless Ventricular Tach	PC7
Pediatric Post Arrest	PC8

Pediatric Medical Section (PM)	
Pediatric Allergic Reaction*	PM1*
Pediatric Diabetic	PM2
Pediatric Hypotension / Shock	PM3

Toxin-Environmental Section (TE)	
Bites and Envenomation's	TE1
Carbon Monoxide / Cyanide	TE2
Drowning	TE3
Hyperthermia	TE4
Hypothermia / Frostbite	TE5
Marine Envenomation / Injury	TE6
Overdose / Toxic Ingestion	TE7
WMD – Nerve Agent	TE8

Special Circumstances Section (SC)	
Suspected Viral Hemorrhagic Fever - Ebola	SC1
High Consequence Pathogen	SC2
Hospice or Palliative Care Patient	SC3
Mass Vaccination / Immunization Med. Distribution	SC4
SARS CoV2 Monoclonal Antibody Administration	SC5

Special Operations (SO)	
Scene Rehabilitation: General	SO1
Scene Rehabilitation: Responder	SO2

Community Paramedic (CP)	
Medication Assisted Treatment (MAT)*	CP1*

Triage and Destination Plans	
General*	*
Trauma and Burn*	*
STEMI*	*
Stroke/ LVO*	*
Pediatric*	*

Policy Section

Policy Section	
Criteria for Death or Withholding Resuscitation	DS1
Deceased Subjects	DS2
Discontinuation of Prehospital Resuscitation	DS3
Disposition (Patient Instructions)	DS4
DNR or MOST	DS5
Patient Without a Protocol	DS6
Physician on Scene	DS7
Opioid Overdose Misuse	DS8
Organ Procurement Agency Notification	DS9
EMS Offload/ Facility Transition of Care	DS10
Child and/or Adult Abuse Recognition and Reporting*	DS11*
Blood Draw on LEO Request*	DS12*
Refusal of Care, Treatment and/or Transport*	DS13*
Transfer of Care to Southport EMS or St. James EMS*	DS14*
EMS to ED Triage Criteria Novant Facilities*	DS15*
Documentation and Data Quality	DC1
Documentation of Vital Signs	DC2
Insurance and Billing Signatures*	DC3*
Transfer of Care to BCEMS Provider at ED*	DC4*
Drug Assisted Airway*	MP1*
Ketamine Program Requirements	MP2
Children with Special Healthcare Needs (NC Kidbase)	Ped1
Infant Abandonment	Ped2
EMS Back in Service Time	SM1
EMS Wheels Rolling Turn-Out (Chute) Time	SM2
EMS Dispatch Center Time	EMD1
State Poison Center	TE1
Air Transport*	TP1*
Safe Transport of Pediatric Patients	TP2
Transport Policy	TP3
Transport of Weapons and Explosives	TP 4

Procedure Section

Airway Procedure	
Biad-I-Gel	AP4
Cricothyrotomy - Surgical	AP5
Intubation: Oral Tracheal	AP6
Intubation: Nasotracheal	AP7
Video Laryngoscope – McGrath*	AP8
Drug Assisted Airway*	AP9*
RSI Checklist*	AP9.1*
Tracheostomy Tube Change	AP10
Endotracheal Intubation Bougie	AP11
Foreign Body Obstruction	AP13
Post Intubation Checklist	AP 14

Assessment Screening	
Assessment: Adult	ASP1
Pain Assessment and Documentation	ASP2
Assessment: Pediatric	ASP3
Blood Glucose Analysis	ASP4
Capnography	ASP5
Pulse Oximetry	ASP6
Reperfusion Checklist	ASP7
Stroke Screen: MEND Exam and RACE*	ASP8*
Temperature Measurement*	ASP9*
Orthostatic Blood Pressure Measurement	ASP10
Verbal De-escalation	ASP11
Fall Risk Screen (Med Frat) *	ASP12*

Cardiac Procedures	
12 Lead ECG and Code STEMI Activation*	CSP1*
Cardioversion	CSP2
External Pacing	CSP3
CPR	CSP4
Defibrillation - Automated	CSP5
Defibrillation - Manual	CSP6
Defibrillation – Dual or Double	CSP7
15 Lead ECG*	CSP8*
Mechanical CPR – LUCAS*	CSP9*

Parenteral Access Procedures	
Arterial Line Maintenance	PAS2
Venous Blood Draw	PAS3
Central Line Maintenance	PAS4
Epidural Catheter Maintenance	PAS5
Ventricular Catheter Maintenance	PAS6
Existing Catheters	PAS7
External Jugular Access	PAS8
Venous - Extremity	PAS9
Intraosseous*	PAS11*
Swan-Ganz Catheter Maintenance	PAS12

Respiratory Procedures	
Suctioning - Advanced	RSP1
Suctioning - Basic	RSP2
Nebulizer Inhalation Therapy	RSP3
NIPPV	RSP4


Universal Procedures	
Childbirth	USP1
Decontamination	USP2
Gastric Tube Insertion	USP3
Gastric Tube Insertion w/ i-gel	USP3.1*
Subcutaneous and Intramuscular	USP4
Restraints: Physical	USP5

Wound and Trauma Procedures	
Chest Decompression	WTP1
Chest Decompression – SPEAR*	WTP1.1*
Spinal Motion Restriction	WTP2
Splinting	WTP3
General	WTP4
Hemostatic Agent	WTP5
Conducted Electrical Weapon Removal	WTP6
Tourniquet*	WTP7*
Trauma Hand-Off Tool*	WTP8*
Pelvic Binder	WTP9*
TXA Topical	WTP10*

Appendix	
Refusal Form Instructions	A
On-Scene Physician Form	B
Apgar Score	C
Los Angeles Prehospital Stroke Screen	D
Pain Scale Forms	E
Restraint Checklist	F
Approved Medical Abbreviations	G
Difficult Airway Evaluation	I
Burn Resources	J
Drug List & Drip Charts	



Universal Patient Care





Universal Patient Care

** At no time should the pediatric weight based dose exceed the adult dose of any medication.**

Scene Safety evaluation:

Identify potential hazards to rescuers, patient and public.

Identify number of patients and utilize SMART triage bag along with START protocol (UP2) if indicated.

General:

All patient care must be appropriate to your level of certification and documented in the PCR.

The PCR should be considered a story of the circumstances, events and care of the patient and should allow the reader to understand the complaint, the assessment, the treatment, why procedures were performed and why indicated procedures were not performed as well as ongoing assessments and responses to treatments and interventions.

Adult Patients:

An adult is considered hypotensive when systolic blood pressure is less than 90 mmHg.

Diabetic patients and women may have atypical presentations of cardiac related problems such as MI.

General weakness can be the symptom of a very serious underlying process.

Geriatric Patients:

Hip fractures and dislocations have high mortality.

AMS is not always dementia. Always check blood sugar and assess signs of stroke, trauma, etc. with any alteration in a patient's baseline mental status.


Minor or moderate injury in typical adults may be very serious in the elderly.

Pearls

- **Recommended Exam: Minimal exam if not noted on the specific protocol is vital signs, mental status with GCS, and location of injury or complaint.**
- Any patient contact, which does not result in an EMS transport, must have a completed Patient Care Report.
- Vital signs should be obtained before, 10 minutes after, and at patient hand off with all pain medications.
- Two complete vital sign acquisitions should occur at a minimum with any patient encounter.
- **Patient Refusal (Declining Treatment and/ or Transport):**
 - Patient refusal is a high risk situation. Encourage patient to accept transport to medical facility.
 - Encourage patient to allow an assessment, including vital signs. Documentation of the event is very important including a mental status assessment describing the patient's capacity to refuse care.
- **Guide to Assessing capacity:**
 - **C: Patient should be able to communicate a clear choice:** This should remain stable over time. Inability to communicate a choice or an inability to express the choice consistently demonstrates incapacity.
 - **R: Relevant information is understood:** Patient should be able to voice a factual understanding of the illness/ injury, the options, and the risks and benefits of recommended treatment or transport.
 - **A: Appreciation of the situation:** Ability to communicate an understanding of the facts of the situation. The patient should be able to recognize the significance of the outcome potentially from their decision.
 - **M: Manipulation of information in a rational manner:** Demonstrate a rational process to come to a decision. Should be able to describe the logic they are using to come to the decision, though you may not agree with decision.
- **Pediatric Patient General Considerations:**
 - A pediatric patient is defined by fitting with a Pediatric Medication/ Skill Resuscitation System, Age ≤ 15, weight ≤ 49 kg.
 - Special needs children may require continued use of Pediatric based protocols regardless of age and weight.
 - Initial assessment should utilize the **Pediatric Assessment Triangle** which encompasses Appearance, Work of Breathing and Circulation to skin.
 - The order of assessment may require alteration dependent on the developmental state of the pediatric patient.
 - Generally the child or infant should not be separated from the caregiver unless absolutely necessary during assessment and treatment.
- Timing of transport should be based on patient's clinical condition and the agency transport policy.
- Consider consultation with Medical Control for patient(s) refusing treatment/ transport.
- Blood Pressure is defined as a Systolic/ Diastolic reading. A palpated Systolic reading may be necessary at times.
- SAMPLE: Signs/ Symptoms; Allergies; Medications; PMH; Last oral intake; Events leading to illness/ injury



Triage





Triage

Pearls

- **When approaching a multiple casualty incident where resources are limited:**
Triage decisions must be made rapidly with less time to gather information
Emphasis shifts from ensuring the best possible outcome for an individual patient to ensuring the best possible outcome for the greatest number of patients.
- **Scene Size Up:**
 1. **Conduct a scene size up. Assure well being of responders. Determine or ensure scene safety before entering. If there are several patients with the same complaints consider HazMat, WMC or CO poisoning.**
 2. **Take Triage system kit.**
 3. **Determine number of patients. Communicate the number of patients and nature of the incident and establish incident command.**
 4. **Direct incoming resources. Identify ingress and egress path. Establish a staging area. Assign a medical officer, triage officer, transportation officer, and staging officer as personnel become available.**
- **Triage is a continual process and is a continuous process in each section as resources allow.**
- **Step 1: Global sorting:**
Call out to those involved in the incident to walk to a designated area and assess group last.
For those who cannot walk, have them wave/ indicate a purposeful movement and assess them second.
Those involved who are not moving, or have an obvious life threat, assess first.
- **Step 2: Individual assessments:**
Control major hemorrhage.
Open airway and if child, give 2 rescue breaths.
Perform Needle Chest Decompression Procedure if indicated.
Administer injector antidotes if indicated.
- **Assess the first patient you encounter using the three objective criteria which can be remembered by RPM.**
R: Respiratory (*Respiratory rates are difficult to measure quickly, use work of breathing and respiratory distress*)
P: Perfusion (*Capillary refill can be altered by many factors including skin temperature – use age appropriate heart rates*)
M: Mental Status (*Motor component of GCS score is important indicator – ability to follow commands*)
- If your patient falls into the RED TAG category, stop, place RED TAG and move on to next patient. Attempt only to correct airway problems, treat uncontrolled bleeding, or administer an antidote before moving to next patient.
- **Treatment:**
Once casualties are triaged, a focus on treatment can begin. You may need to move patients to treatment areas.
RED TAGs are moved/ treated first, followed by YELLOW TAGs. BLACK TAGs should remain in place.
You may also indicate deceased patients by pulling their shirt/ clothing over their head.
As more help arrives, then the triage/ treatment process may proceed simultaneously.
- **Lightning strike (Reverse Triage):**
Lightning strike victims are amenable to airway, breathing, cardiac compressions as well as early defibrillation.
Use concept of reverse triage with multiple casualties. Resuscitate lightning strikes as the priority.
Lightning strike victims found alive do not often deteriorate quickly.
- **SMART triage tag system is utilized in NC.**



Abdominal Pain Vomiting and Diarrhea

History

- Age
- Time of last meal
- Last bowel movement/emesis
- Improvement or worsening with food or activity
- Duration of problem
- Other sick contacts
- Past medical history
- Past surgical history
- Medications
- Menstrual history (pregnancy)
- Travel history
- Bloody emesis / diarrhea

Signs and Symptoms

- Pain
- Character of pain (constant, intermittent, sharp, dull, etc.)
- Distention
- Constipation
- Diarrhea
- Anorexia
- Radiation

Associated symptoms:

Fever, headache, blurred vision, weakness, malaise, myalgias, cough, headache, dysuria, mental status changes, rash

Differential

- CNS (increased pressure, headache, stroke, CNS lesions, trauma or hemorrhage, vestibular)
- Myocardial infarction
- Drugs (NSAID's, antibiotics, narcotics, chemotherapy)
- GI or Renal disorders
- Diabetic ketoacidosis
- OB-Gyn disease (ovarian cyst, PID, Pregnancy)
- Infections (pneumonia, influenza)
- Electrolyte abnormalities
- Food or toxin induced
- Medication or Substance abuse
- Psychological

	Consider Blood Glucose Analysis Procedure
B	12 Lead ECG Procedure
	IV or IO Access Protocol UP 6
P	Cardiac Monitor
	Age Appropriate Diabetic Protocol AM 2/ PM 2 if indicated
	Pain Control Protocol UP 11 if indicated
	Age Appropriate Cardiac Protocol(s) if indicated

Age Specific Blood Pressure indicating possible shock

Age 0 – 28 days: SBP < 60
 Ages ≥ 1 month: SBP < 70
 Age 1 – 9: SBP < 70 + (2x Age)

Ages 10 – 64: SBP < 90
 Ages ≥ 65: SBP < 110

All ages Shock Index: HR > SBP

Serious Signs/ Symptoms
 Hypotension, poor perfusion, shock

NO

A	Normal Saline IV TKO Or Saline Lock
A	Ondansetron 4 mg IV / IO / ODT / PO / IM Peds: 0.2 mg/kg Peds Maximum 4 mg May repeat in 15 minutes
A	<u>If no response in adults</u> Promethazine 12.5 mg PO / IM / IVPB May repeat x 1 as needed

	IV or IO Access Protocol UP 6 Consider 2 Large Bore sites
A	Normal Saline 500 mL Bolus Repeat as needed Titrate SPB ≥ 90 mmHg Maximum 2 L Peds: 20 mL/kg IV / IO Repeat as needed Titrate to Age Appropriate SBP ≥ 70 + (2 x Age) Maximum 60 mL/kg
A	Ondansetron 4 mg IV / IO / ODT / PO / IM Peds: 0.2 mg/kg Peds Maximum 4 mg May repeat in 15 minutes
	Age Appropriate Hypotension/ Shock Protocol AM 5/ PM 3 if indicated

Monitor and Reassess

Notify Destination or Contact Medical Control

Promethazine:
 FDA advises that deep intramuscular (IM) administration or intravenous piggyback (IVPB) are acceptable under strict adherence to the new guidelines (see pearls)

Universal Protocol Section



Abdominal Pain Vomiting and Diarrhea

Promethazine (Phenergan)

- IM is preferred
- Use with caution as it has a strong sedative effect.
- Consider combined effects with narcotic co-administration.
- If giving IV, dilute in 100cc saline and administer slowly over 20 minutes.
- May be administered first line when PT is allergic to Zofran

Dystonic reactions:

- can occur with co-administration of psych meds.
- involuntary contractions of muscles in the extremities, face, neck, abdomen, pelvis, or larynx in either sustained or intermittent patterns that lead to abnormal movements or postures
- treat with Benadryl 25-50mg

Pearls

- **Recommended Exam: Mental Status, Skin, HEENT, Neck, Heart, Lungs, Abdomen, Back, Extremities, Neuro**
- **Abdominal/ back pain in women of childbearing age should be treated as pregnancy related until proven otherwise.**
- **The diagnosis of abdominal aneurysm should be considered with abdominal pain, with or without back and/ or lower extremity pain or diminished pulses, especially in patients over 50 and/ or patients with shock/ poor perfusion. Notify receiving facility early with suspected abdominal aneurysm.**
- **Consider cardiac etiology in patients > 35, diabetics and/ or women, especially with upper abdominal complaints.**
- **Heart Rate: Tachycardia is one of the first clinical signs of dehydration and volume depletion and typically increases as dehydration becomes more severe.**
- **Nausea without vomiting should be treated like vomiting. Patient will benefit from symptom control with antiemetic even if not actively vomiting.**
- **Promethazine (Phenergan):**
May cause sedative effects in pediatric patients and in ages ≥ 65 , and the debilitated, etc.)
When giving promethazine, PO and IM is preferred over IV administration. If giving IV, dilute with 100 mL of normal saline and administer slowly over at least 20 minutes as it can cause severe chemical irritation and tissue damage. Promethazine should be administered through large patent veins, avoid veins in the hand and wrist.
- Isolated vomiting in children is common but can be a sign of more serious pathology. Pyloric stenosis, bowel obstruction, and CNS processes (bleeding, tumors, or increased CSF pressures) all often present with vomiting.
- Vomiting and diarrhea are common symptoms, but can be the symptoms of uncommon and serious pathology such as stroke, CO poisoning, acute MI, new onset diabetes, diabetic ketoacidosis (DKA), and organophosphate poisoning. Maintain a high index of suspicion for serious pathology.



Altered Mental Status

History

- Known diabetic, medic alert tag
- Drugs, drug paraphernalia
- Report of illicit drug use or toxic ingestion
- Past medical history
- Medications
- History of trauma
- Change in condition
- Changes in feeding or sleep habits


Signs and Symptoms

- Decreased mental status or lethargy
- Change in baseline mental status
- Bizarre behavior
- Hypoglycemia (cool, diaphoretic skin)
- Hyperglycemia (warm, dry skin; fruity breath; Kussmaul respirations; signs of dehydration)
- Irritability

Differential

- Head trauma
- CNS (stroke, tumor, seizure, infection)
- Cardiac (MI, CHF)
- Hypothermia
- Infection (CNS and other)
- Thyroid (hyper / hypo)
- Shock (septic, metabolic, traumatic)
- Diabetes (hyper / hypoglycemia)
- Toxicological or Ingestion
- Acidosis / Alkalosis
- Environmental exposure
- Pulmonary (Hypoxia)
- Electrolyte abnormality
- Psychiatric disorder

Age Appropriate Airway Protocol(s) AR 1, 2, 3, 5, 6 if indicated	
	Blood Glucose Analysis Procedure
B	12 Lead ECG Procedure
IV or IO Access Protocol UP 6	
Age Appropriate Diabetic Protocol(s) AM 2/ PM 2 if indicated	





Altered Mental Status

Refusal Checklist

- o Is the patient >17 y/o/a or an emancipated minor?
- o Can the patient retain and comprehend relevant information?
- o Can the patient believe information?
- o Can the patient use information to make a choice?
- o Is the patient NOT DANGEROUS to self or others (i.e. no suicidal or homicidal ideation)?

If all are "YES" then the patient has the capacity to decline further care/transport. If any are "NO" then the patient does not have capacity to make his/her own medical decisions. Document these concepts clearly in your assessment - simply stating "alert and oriented" is not sufficient. Do not hesitate to contact medical control if there are questions about capacity.

Pearls

- **Recommended Exam: Mental Status, HEENT, Skin, Heart, Lungs, Abdomen, Back, Extremities, Neuro.**
- **AMS may present as a sign of an environmental toxin or Haz-Mat exposure, protect personal safety.**
- **General:**
 - The patient with AMS poses one of the most significant challenges.**
 - A careful assessment of the patient, the scene, and the circumstances should be undertaken.**
 - Assume the patient has a life threatening cause of their AMS until proven otherwise.**
 - Pay careful attention to the head exam for signs of bruising or other injury.**
 - Information found at the scene must be communicated to the receiving facility.**
 - Patients not able to communicate with you coherently require a complete secondary survey (head-to-toe) exam to assess for trauma, infection, or signs of maltreatment/ abuse, or neglect.**
 - Acute Stroke should be considered in all patients with acute AMS when < 24 hours from onset.**
- **Substance misuse:**
 - Patients ingesting substances can pose a great challenge.
 - DO NOT assume recreational drug use and/ or alcohol are the sole reasons for AMS.
 - Misuse of alcohol/ recreational drugs may lead to hypoglycemia or occult trauma.
 - More serious underlying medical and trauma conditions may be the cause.
- **Behavioral health:**
 - The behavioral health patient may present a great challenge in forming a differential.
 - DO NOT assume AMS is the result solely of an underlying psychiatric etiology.
 - Often an underlying medical or trauma condition precipitates a deterioration of a patients underlying disease.
- **Spinal Motion Restriction/ Trauma:**
 - Only utilize spinal immobilization if the situation warrants.
 - The patient with AMS may worsen with increased agitation when immobilized.
- **It is safer to assume hypoglycemia than hyperglycemia if doubt exists. Recheck blood glucose after Dextrose or Glucagon**
- Consider Restraints if necessary for patient's and/ or personnel's protection per USP 5 Restraints: Physical procedure.



Back Pain

History


- Age
- Past medical history
- Past surgical history
- Medications
- Onset of pain / injury
- Previous back injury
- Traumatic mechanism
- Location of pain
- Fever
- Improvement or worsening with activity

Signs and Symptoms

- Pain (paraspinous, spinous process)
- Swelling
- Pain with range of motion
- Extremity weakness
- Extremity numbness
- Shooting pain into an extremity
- Bowel / bladder dysfunction

Differential

- Muscle spasm / strain
- Herniated disc with nerve compression
- Sciatica
- Spine fracture
- Kidney stone
- Pyelonephritis
- Aneurysm
- Pneumonia
- Spinal Epidural Abscess
- Metastatic Cancer
- AAA





Back Pain

Pearls

- **Recommended Exam: Mental Status, Heart, Lungs, Abdomen, Neuro, Lower extremity perfusion, Back**
- **Back pain is one of the most common complaints in medicine and affects more than 90% of adults at some point in their life. Back pain is also common in the pediatric population. Most often it is a benign process but in some circumstances can be life or limb threatening.**
- **Consider pregnancy or ectopic pregnancy with abdominal or back pain in women of childbearing age.**
- **Consider abdominal aortic aneurysm with abdominal pain especially in patients over 50 and/ or patients with shock/ poor perfusion. Patients may have abdominal pain and/ or lower extremity pain with diminished pulses. Notify receiving facility early with suspected abdominal aneurysm.**
- **Consider cardiac etiology in patients > 35, diabetics and/ or women especially with upper abdominal complaints.**
- **Red Flags which may signal a more serious process associated with back pain:**
 - Age > 50 or < 18
 - Neurological deficit (leg weakness, urinary retention, or bowel incontinence)
 - IV Drug use
 - Fever
 - History of cancer, either current or remote
 - Night time pain in pediatric patients
- **Cauda equina syndrome is where the terminal nerves of spinal cord are being compressed (Symptoms include):**
 - Saddle anesthesia (numbness between the genitalia and rectum)
 - Recent onset of bladder and bowel dysfunction. (Urine retention and bowel incontinence)
 - Severe or progressive neurological deficit in the lower extremity.
 - Motor weakness of thigh muscles or foot drop
- **Back pain associated with infection:**
 - Fever/ chills.
 - IV Drug user (consider spinal infection)
 - Recent bacterial infection like pneumonia.
 - Immune suppression such as HIV or patients on chronic steroids like prednisone.
 - Meningitis.
- **Spinal motion restriction in patients with underlying spinal deformity should be maintained in their functional position.**
- **Kidney stones typically present with an acute onset of flank pain which radiates around to the groin area.**



IV or IO Access

History


- Chronic medical conditions requiring recurrent need for IV access for medication, hydration, or blood sampling.
- Medical condition requiring administration of IV medications at home.
- End-stage renal disease requiring hemodialysis.
- Chronic medical condition requiring IV nutrition.

Signs and Symptoms

- Fever
- Bleeding
- Hypotension
- Redness, swelling, and/or pain at IV catheter site
- Shortness of breath
- Chest pain
- IV catheter patency

Differential

- Infection or sepsis
- Infection of catheter
- Clotted IV catheter
- Air embolism
- Pneumothorax
- Overdose of home medication
- Shock



Universal Protocol Section

Pearls

- Frequent encounter of patients with IV access devices and confusion as to which device can be accessed and used by EMS providers are common.
- If unclear about device use, always ask "Is this device used for dialysis?"
- When accessing central catheter, always ensure sterility of catheter connection point by cleaning port with alcohol, or similar disinfectant, 2 – 3 times prior to access.
- Central line catheters placed for administration of chemotherapy, medications, electrolytes, antibiotics, and blood are available to EMS providers for access and administration of fluids, medications, antibiotics, and blood products.
- Central line catheters placed for hemodialysis are NOT available for access by EMS providers unless the patient is in cardiac arrest.
- Long term IV access is frequently needed for a variety of indications:
 - Medication administration such as antibiotics, pain relief, or chemotherapy.
 - Administration of IV nutrition or feeding.
 - Need for multiple IV line access or recurrent blood sampling.
 - Poor vasculature requiring repeated attempts at IV access.
 - End-stage renal disease requiring hemodialysis.
- Common complications of central access devices:

Infection	Loss of patency due to clogging or clotting
Damage to vasculature	Pneumothorax
Air embolism	

IV or IO Access

Types of IV catheters:


Port-a-Cath® :

Surgically implanted device allowing easy access to venous system. The port and the catheter are all placed beneath the skin. Requires a special kit and a specific needle to access.

Paramedic does NOT routinely access this device.

Paramedic may utilize if already accessed with needle/ extension.

Paramedic may access if trained on procedure with access to proper equipment.



Dialysis Catheter:

Surgically implanted device used to access the vasculature for hemodialysis.

May be tunneled under the skin with access on outside of skin surface or may be non-tunneled with greater portion of catheter on outside of skin surface.

Catheter has a RED port indicating use for dialysis:


Most catheters have a RED port and a BLUE port.

Some catheters have a RED port and a WHITE port.

Dialysis catheters may be used for both short and long-term dialysis and should not be accessed or used for delivery of fluids, medications, antibiotics, or blood products as it increases risk of infection, which then requires removal and subsequent loss of dialysis access.

Paramedic and AEMT do NOT routinely access this device.

Paramedic and AEMT MAY access during cardiac arrest only (Only if IV or IO access cannot be established.)



PICC (Peripherally Inserted Central Catheters):


Long catheter inserted into a vein in arm or leg (less common) with the tip of the catheter positioned into the central circulation.

Used for long-term IV fluids, medication administration, blood administration or blood draws.

May have 1 or 2 ports (possibly more, but less common.)

Port ends usually white, blue, or purple. (May be red, less common and is not used for dialysis.)

Paramedic and AEMT may access and utilize following clean technique.



Central Lines:


Catheter placed in large vein in the neck, under the clavicle, or in the groin.

Used for long-term IV fluids, medication administration, blood administration or blood draws.

May have 1 - 4 ports (possibly more, but less common.)

Port ends usually white, blue, or purple. (May be red, less common and is not used for dialysis.)

Paramedic and AEMT may access and utilize following sterile technique.





Dental Problems

History


- Age
- Past medical history
- Medications
- Onset of pain / injury
- Trauma with "knocked out" tooth
- Location of tooth
- Whole vs. partial tooth injury

Signs and Symptoms

- Bleeding
- Pain
- Fever
- Swelling
- Tooth missing or fractured

Differential

- Decay
- Infection
- Fracture
- Avulsion
- Abscess
- Facial cellulitis
- Impacted tooth (wisdom)
- TMJ syndrome
- Myocardial infarction



Universal Protocol Section

Pearls

- **Recommended Exam: Mental Status, HEENT, Neck, Chest, Lungs, Neuro**
- Significant soft tissue swelling to the face or oral cavity can represent a cellulitis or abscess.
- Scene and transport times should be minimized in complete tooth avulsions. Reimplantation is possible within 4 hours if the tooth is properly cared for, but unlikely when > 1 hour from time of injury.
- Cardiac chest pain may radiate to the jaw and teeth mimicking dental pain.
- **Avulsed tooth:**
Handle tooth by the crown, do not touch the root.
Rinse tooth if soiled but do not scrub, as this can damage the ligaments vital for possible reimplantation.
Rinse with mild, commercial tooth solution, normal saline or lactated ringers, or the patient's own saliva if dry.
Transport tooth in milk, commercial solution, patient's own saliva, or IV solution in a container to protect.
- **TXA Use in Dental Bleeding:**
May be used topically and is approved by agency Medical Director.
Procedure (WTP 10) has been created with specific guidance on how to administer TXA for dental bleeding.
TXA offers modest benefit as TXA instilled gauze combined with direct pressure.



Emergencies Involving Indwelling Central Lines

History


- Central Venous Catheter Type
Tunneled Catheter (Broviac / Hickman)
- PICC (peripherally inserted central catheter)
- Implanted catheter (Mediport / Hickman)
- Occlusion of line
- Complete or partial dislodge
- Complete or partial disruption

Signs and Symptoms

- External catheter dislodgement
- Complete catheter dislodgement
- Damaged catheter
- Bleeding at catheter site
- Internal bleeding
- Blood clot
- Air embolus
- Erythema, warmth or drainage about catheter site indicating infection

Differential

- Fever
- Hemorrhage
- Reactions from home nutrient or medication
- Respiratory distress
- Shock



Universal Protocol Section

Pearls

- **Always talk to family / caregivers as they have specific knowledge and skills.**
- **Use strict sterile technique when accessing / manipulating an indwelling catheter.**
- **Cardiac arrest: May access central catheter and utilize if functioning properly.**
- Do not attempt to force catheter open if occlusion evident.
- Some infusions may be detrimental to stop. Ask family or caregiver if it is appropriate to stop or change infusion.
- Hyperalimentation infusions (IV nutrition): If stopped for any reason monitor for hypoglycemia.



Epistaxis

History


- Age
- Past medical history
- Medications (HTN, anticoagulants, aspirin, NSAIDs)
- Previous episodes of epistaxis
- Trauma
- Duration of bleeding
- Quantity of bleeding

Signs and Symptoms

- Bleeding from nasal passage
- Pain
- Nausea
- Vomiting

Differential

- Trauma
- Infection (viral URI or Sinusitis)
- Allergic rhinitis
- Lesions (polyps, ulcers)
- Hypertension



Age Specific Blood Pressure indicating possible shock

Age 0 – 28 days: SBP < 60
 Ages ≥ 1 month: SBP < 70
 Age 1 – 9: SBP < 70 + (2x Age)

Ages 10 – 64: SBP < 90
 Ages ≥ 65: SBP < 110

All ages Shock Index:
 HR > SBP

Pearls

- **Recommended Exam: Mental Status, HEENT, Heart, Lungs, Neuro**
- **TXA Use in Epistaxis:**
 May be used topically and is approved by agency Medical Director.
 Procedure (WTP10) has been created with specific guidance on how to administer TXA for epistaxis.
 No clear evidence that TXA provides benefit over conventional vasoconstrictors and sustained direct pressure.
- It is very difficult to quantify the amount of blood loss with epistaxis.
- Bleeding may also be occurring posteriorly. Evaluate for posterior blood loss by examining the posterior pharynx.
- Anticoagulants include warfarin (Coumadin), Apixaban (Eliquis), heparin, enoxaparin (Lovenox), dabigatran (Pradaxa), rivaroxaban (Xarelto), and many over the counter headache relief powders.
- Anti-platelet agents like aspirin, clopidogrel (Plavix), aspirin/ dipyridamole (Aggrenox), and ticlopidine (Ticlid) can contribute to bleeding.



Fever/ Infection Control

History

- Age
- Duration of fever
- Severity of fever
- Past medical history
- Medications
- Immunocompromised (transplant, HIV, diabetes, cancer)
- Environmental exposure
- Last acetaminophen or ibuprofen

Signs and Symptoms

- Warm
- Flushed
- Sweaty
- Chills/Rigors


Associated Symptoms (Helpful to localize source)

- Myalgias, cough, chest pain, headache, dysuria, abdominal pain, mental status changes, rash

Differential

- Infections / Sepsis
- Cancer / Tumors / Lymphomas
- Medication or drug reaction
- Connective tissue disease
 - Arthritis
 - Vasculitis
- Hyperthyroidism
- Heat Stroke
- Meningitis

	Contact, Droplet, and Airborne Precautions See Pearls
B	Temperature Measurement Procedure <i>if available</i>
	IV or IO Access Protocol UP 6 <i>If indicated</i>



Universal Protocol Section

Pearls

- **Recommended Exam: Mental Status, Skin, HEENT, Neck, Heart, Lungs, Abdomen, Back, Extremities, Neuro**
- Febrile seizures are more likely in children with a history of febrile seizures and with a rapid elevation in temperature.
- Patients with a history of liver failure should not receive acetaminophen.
- **Droplet precautions** include standard PPE plus a standard surgical mask for providers who accompany patients in the back of the ambulance and a surgical mask or NRB O2 mask for the patient. This level of precaution should be utilized when influenza, meningitis, mumps, streptococcal pharyngitis, and other illnesses spread via large particle droplets are suspected. A patient with a potentially infectious rash should be treated with droplet precautions.
- **Airborne precautions** include standard PPE plus utilization of a gown, change of gloves after every patient contact, and strict hand washing precautions. This level of precaution is utilized when multi-drug resistant organisms (e.g. MRSA), scabies, or zoster (shingles), or other illnesses spread by contact are suspected.
- **All-hazards precautions** include standard PPE plus airborne precautions plus contact precautions. This level of precaution is utilized during the initial phases of an outbreak when the etiology of the infection is unknown or when the causative agent is found to be highly contagious (e.g. **SARS, SARS-CoV-2, COVID-19, MERS, Monkeypox**).
- Rehydration with fluids increases the patient's ability to sweat and improves heat loss.
- Allergies to NSAIDs (non-steroidal anti-inflammatory medications) are a contraindication to Ibuprofen. Do not give to patients who have renal disease or renal transplant.
- NSAIDs should not be used in the setting of environmental heat emergencies.
- **Do not** give aspirin to a child, age ≤ 15 years.
- Agency Medical Director does not require contact of medical control prior to EMT administering any medication for fever. EMR may administer Acetaminophen.



Pain Control

History

- Age
- Location
- Duration
- Severity (1 - 10)
- If child use Wong-Baker faces scale
- Past medical history
- Medications
- Drug allergies

Signs and Symptoms

- Severity (pain scale)
- Quality (sharp, dull, etc.)
- Radiation
- Relation to movement, respiration
- Increased with palpation of area

Differential

- Per the specific protocol
- Musculoskeletal
- Visceral (abdominal)
- Cardiac
- Pleural / Respiratory
- Neurogenic
- Renal (colic)

Specific Complaint Protocol

Assess Pain Severity

Combination of Pain Scale, MOI, circumstances, Injury or Illness severity

Mild

Moderate to Severe

Ibuprofen 10 mg/kg PO
(400 – 600 mg typical adult)
Maximum 800 mg
Or
Acetaminophen 15 mg/kg PO
(325 – 1000 mg PO/IV typical adult)
Maximum 1000 mg PO
Or
Aspirin 324 to 650 mg PO
(≥ 16 only)

If no improvement
Consider IV Protocol UP 6
if indicated

Acetaminophen 15 mg / kg IV

Monitor and Reassess
Every 10 minutes following sedative

Notify Destination or
Contact Medical Control

IV / IO Protocol UP 6

Acetaminophen 1000mg /100mL IV
over 15 minutes
Peds: 15 mg/kg IV over 15 minutes
Ketorolac 15 mg IV / IO; 30 mg IM
Peds: 0.5 mg/kg IV / IO / IM
Maximum 30 mg

Cardiac Monitor

Nitrous Oxide 70:30 to 50:50 Mix NO/O₂

Fentanyl 50 – 100 mcg IV / IO / IM / IN
Repeat every 5 minutes
Maximum 3 mcg/kg
Peds: 1 mcg/kg IV / IO / IM / IN
May repeat 0.5 mcg/kg every 5 minutes
Maximum 2 mcg/kg
Or
Morphine 4 mg IV / IO / IM
Repeat **2 mg** every 5 minutes as needed
Peds: 0.1 mg/kg IV / IO / IM
May repeat every 5 minutes
Maximum 10 mg

Adult ONLY:
Ketamine 0.3 mg/kg IV /IO
Infuse over 10 minutes
May repeat every 20 minutes
maximum single dose 30 mg
Maximum 3 doses
OR
Ketamine 1 mg/kg IN
Maximum 1 total dose

Universal Protocol Section



Pain Control

Document OPQRST:

- **Onset**- What was the patient doing when it started? (active, inactive, stressed, etc.)
- **Provocation / Palliation** - Does the patient believe that activity prompted the pain, and whether the onset was sudden, gradual or part of an ongoing chronic problem.
- **Quality** - Have the patient describe the pain. Is it dull, throbbing, aching, burning, sharp, crushing, shooting, tearing, constant, intermittent, etc.?
- **Region and Radiation** - Where is the pain and does it move?
- **Severity** - Use the 0-10 pain scale. This can be comparative ("...compared to the worst pain you have ever experienced") or imaginative ("...compared to having your leg broken").
- **Timing** - Identify when the pain started. How long has the condition been going on?

Document - amount of pain medication wasted in ePCR even if equal to zero, along with witness and location.

Pearls

- **Recommended Exam: Mental Status, Area of Pain, Neuro**
- **Pain severity (0-10) is a vital sign to be recorded before and after PO, IV, IO or IM medication delivery and at patient hand off. Monitor BP closely as sedative and pain control agents may cause hypotension.**
- **Ketamine:**
Ketamine may be used in patients who are outside a Pediatric Medication/Skill Resuscitation System product. Ketamine may be used in patients who fit within a Pediatric Medication/Skill Resuscitation System product only with DIRECT ONLINE MEDICAL ORDER, by the system MEDICAL DIRECTOR or ASSISTANT MEDICAL DIRECTOR.
- **Ketamine: appropriate indications for pain control:**
Patients who have developed opioid-tolerance or opioid use disorder (OUD). Sickle cell crisis patients with opioid-tolerance. Patients who have obstructive sleep apnea.
May use in combination with opioids to limit total amount of opioid administration.
- **Ketamine: caution when using for pain control:**
Slow infusion or IV push over 10 minutes is associated with less side effects. Do not administer by rapid IV push.
Avoid in patients who have cardiac disease or uncontrolled hypertension.
Avoid in patients with increased intraocular pressure such as glaucoma.
Avoid use in combination with benzodiazepines due to decreased respiratory drive..
- **Both arms of the treatment may be used in concert. For patients in Moderate pain for instance, you may use the combination of an oral medication and parenteral if no contraindications are present.**
- **Pediatrics:**
For children use Wong-Baker faces scale or the FLACC score (see Assessment Pain Procedure ASP2)
Use Numeric (> 9 yrs), Wong-Baker faces (4-16yrs) or FLACC scale (0-7 yrs) as needed to assess pain
- **Vital signs should be obtained before, 10 minutes after, and at patient hand off with all pain medications.**
- All patients who receive IM or IV medications must be observed 15 minutes for drug reaction in the event no transport occurs.
- Do not administer **Acetaminophen** to patients with a history of liver disease.
- Burn patients may require higher than usual opioid doses to titrate adequate pain control.
- Consider agency-specific anti-emetic(s) for nausea and/or vomiting.



Police Custody

History


- Traumatic Injury
- Drug Abuse
- Cardiac History
- History of Asthma
- Psychiatric History

Signs and Symptoms

- External signs of trauma
- Palpitations
- Shortness of breath
- Wheezing
- Altered Mental Status
- Intoxication/Substance Abuse

Differential

- Agitated Delirium Secondary to Psychiatric Illness
- Agitated Delirium Secondary to Substance Abuse
- Traumatic Injury
- Closed Head Injury
- Asthma Exacerbation
- Cardiac Dysrhythmia





Police Custody

Anyone who has been tased **MUST** receive a full assessment including a 12 Lead ECG.

Pearls

- **Patient does not have to be in police custody or under arrest to utilize this protocol.**
- **Local EMS agencies should formulate a policy with local law enforcement agencies concerning patients requiring EMS and Law Enforcement services simultaneously.**
- **Agencies should work together to formulate a disposition in the best interest of the patient.**
- **Patients restrained by law enforcement devices must be transported and accompanied by a law enforcement officer in the patient compartment who is capable of removing the devices.**

- **All patients who receive either physical and chemical restraint must be continuously observed by ALS personnel on scene or immediately upon their arrival.**
- The responsibility for patient care rests with the highest authorized medical provider on scene per North Carolina law.
- If an asthmatic patient is exposed to irritant/ pepper spray and released to law enforcement, all parties should be advised to immediately contact EMS if wheezing/ difficulty breathing occurs.
- All patients with decision-making capacity in police custody retain the right to participate in decision-making regarding their care and may request care or refuse care of EMS.
- If extremity/ chemical/ law enforcement restraints are applied, follow USP 5 Restraints: Physical.
- **Consider Haldol or Droperidol for patients with history of psychosis or a benzodiazepine for patients with presumed substance misuse.**
- **Haldol is acceptable treatment in pediatric patients ≥ 12 years old. Safety and efficacy is not established in younger ages. Contact Medical Control for advice as needed.**
- **Hyperactive Delirium with Severe Agitation:**
 - Medical emergency: Combination of delirium, psychomotor agitation, anxiety, hallucinations, speech disturbances, disorientation, violent/ bizarre behavior, insensitivity to pain, hyperthermia and increased strength. Potentially life-threatening and associated with use of physical control measures, including physical restraints and Tasers.
 - Most commonly seen in male subjects with a history of serious mental illness and/or acute or chronic drug abuse, particularly stimulant drugs such as cocaine, crack cocaine, methamphetamine, amphetamines or similar agents. Alcohol withdrawal or head trauma may also contribute to the condition.
 - If patient suspected of EDS suffers cardiac arrest, consider a fluid bolus and sodium bicarbonate early.**
- Do not position or transport any restrained patient in such a way that could impact the patient's respiratory or circulatory status.
- Patients exposed to chemical spray, with or without history of respiratory disease, may develop respiratory complaints up to 20 minutes post exposure.



Seizure

History

- Reported / witnessed seizure activity
- Previous seizure history
- Medical alert tag information
- Seizure medications
- History of trauma
- History of diabetes
- History of pregnancy
- Time of seizure onset
- Document number of seizures
- Alcohol use, abuse or abrupt cessation
- Fever

Signs and Symptoms

- Decreased mental status
- Sleepiness
- Incontinence
- Observed seizure activity
- Evidence of trauma
- Unconscious

Differential

- CNS (Head) trauma
- Tumor
- Metabolic, Hepatic, or Renal failure
- Hypoxia
- Electrolyte abnormality (Na, Ca, Mg)
- Drugs, Medications, Non-compliance
- Infection / Fever
- Alcohol withdrawal
- Eclampsia
- Stroke
- Hyperthermia
- Hypoglycemia

Age Appropriate Airway Protocol(s) AR 1, 2, 3, 5, 6 as indicated
Altered Mental Status Protocol UP 4 if indicated
Childbirth/ Labor Protocol AO 1 Obstetrical Emergency Protocol AO 3 if indicated
Behavioral Protocol UP 17, 18, 19 if indicated
Loosen any constrictive clothing Protect patient

Active Seizure Activity

IV / IO Access

NO

NO

P	≥ 49 kg Midazolam 10 mg IM OR Lorazepam 2mg IM OR Diazepam 10 mg PR <i>(See page 2 for repeat dosing)</i>
	< 49 kg Midazolam 0.2 mg/kg IM Max single dose 5 mg OR Lorazepam 0.1 mg/kg IM, max 1mg OR Diazepam 0.5 mg/kg PR, Max 10 mg <i>(See page 2 for repeat dosing)</i>

P	Midazolam 2 – 2.5 mg IV / IO Peds: Midazolam 0.2 mg/kg IV / IO repeat every 3 - 5 minutes as needed Maximum 10 mg OR Lorazepam 1 mg IV/IO Peds: 0.1 mg/kg IV/IO, max 1mg repeat every 10 minutes as needed up to 4 mg OR Diazepam 5 mg IV/IO Peds: 0.1 mg/kg IV/IO, Maximum 4 mg
	----- -For seizures refractory to Benzodiazepines: Levetiracetam (Keppra) 1 gram in 100cc NS over 15 minutes IV/IO Peds: 15 mg/kg of mixture over 15 minutes Max 1 gram

P	Blood Glucose Analysis Procedure
	IV or IO Access Protocol UP 6 if indicated
P	Cardiac Monitor if indicated
	Monitor and Reassess

Notify Destination or Contact Medical Control

Universal Protocol Section



Seizure

Active seizure in known or suspected pregnancy greater than 20 weeks, administer Magnesium Sulfate 2g IV/IO over 2-3 minutes, may repeat dose x1. See Protocol AO3.

Repeat dosing:

Midazolam:

≥49kg- may repeat in 5 minutes if needed, Maximum 20mg
<49kg- may repeat in 5 minutes if needed, Maximum 10mg

Lorazepam: (IV/IO preferred over IM)

Adult IV/IO/IM- may repeat every 10 minutes as needed up to 4mg
Pediatric IV/IO/IM- may repeat every 10 minutes as needed up to 4 mg, Maximum single dose = 1mg

Pearls

- **Recommended Exam: Mental Status, HEENT, Heart, Lungs, Extremities, Neuro**
- **Items in Red Text are key performance measures used to evaluate protocol compliance and care.**
- **Brief seizure-like activity can be seen following ventricular fibrillation or ventricular tachycardia associated cardiac arrest.**
- **Status epilepticus is defined by seizure activity lasting > 5 minutes or multiple seizures without return to baseline.**
- **Most seizure activity is brief, lasting only 1 – 2 minutes, and is associated with transient hypoventilation.**
- **Be prepared for airway problems and continued seizures.**
- **Seizure activity may be a marker of closed head injury, especially in the very young, examine for trauma.**
- **Adult:**
 - **Midazolam 10 mg IM is effective in termination of seizures.**
 - **Do not delay IM administration with difficult IV or IO access. IM Preferred over IO.**
- **Pediatrics:**
 - **Midazolam 0.2 mg/kg (Maximum 5 mg) IM is effective in termination of seizures.**
 - **Do not delay IM administration with difficult IV or IO access. IM Preferred over IO.**
- **Do not delay administration of anti-epileptic drugs to check for blood glucose.**
- **Grand mal seizures (generalized)** are associated with loss of consciousness, incontinence, and tongue trauma.
- **Focal seizures** affect only a part of the body and are not usually associated with a loss of consciousness, but can propagate to generalized seizures with loss of consciousness.
- **Be prepared to assist ventilations especially if diazepam or midazolam is used.**
- **For any seizure in a pregnant patient, follow the AO 3 Obstetrical Emergencies Protocol.**
- **Midazolam (Versed) is shown to be as effective with IM route as Lorazepam (Ativan) is via the IV or IO route.**
- **Lorazepam (Ativan) is not as effective when administered IM. IV or IO route is preferred.**
- **Diazepam (Valium) is not effective when administered IM. Give IV or Rectally.**
- **Optimal conditions for patients refusing transport following a seizure:**

Known history of seizures/epilepsy	Seizure not associated with drugs or alcohol
Full recovery to baseline mental status	Only 1 seizure episode in the past hour
No injuries requiring treatment or evaluation	Seizure not associated with pregnancy
Adequate supervision	



Suspected Stroke

History


- Previous CVA, TIA's
- Previous cardiac / vascular surgery
- Associated diseases: diabetes, hypertension, CAD
- Atrial fibrillation
- Medications (blood thinners)
- History of trauma
- Sickle Cell Disease
- Immune disorders
- Congenital heart defects
- Maternal infection / hypertension

Signs and Symptoms

- Altered mental status
- Weakness / Paralysis
- Blindness or other sensory loss
- Aphasia / Dysarthria
- Syncope
- Vertigo / Dizziness
- Vomiting
- Headache
- Seizures
- Respiratory pattern change
- Hypertension / hypotension

Differential

- See Altered Mental Status
- TIA (Transient ischemic attack)
- Seizure
- Todd's Paralysis
- Hypoglycemia
- Stroke
 - Thrombotic or Embolic (~85%)
 - Hemorrhagic (~15%)
- Tumor
- Trauma
- Dialysis / Renal Failure



Transport Destination Facility based on:
Stroke
EMS Triage and Destination Plan

- Refer to plan for Last Known Well Times and Stroke Severity Assessment Values
- Combination of factors will drive transport destination decision-making based on your local region

Transport based on:
STROKE
EMS Triage and Destination Plan

Keep Scene Time to ≤ 15 Minutes
Provide Early Notification of Destination Facility Suspected Stroke Alert

	Reperfusion Checklist Procedure <i>if time allows completion</i>
B	Cardiac Monitor
	IV or IO Access Protocol UP 6 Preferably 2 Sites <i>See Pearls</i>
A	Venous Access Blood Draw Procedure <i>if applicable</i>
	Cardiac Monitor
P	Persistent Hypertension: If SBP ≥ 185 or DBP ≥ 110 mmHg After 3 readings, 5 minutes apart Contact Receiving Facility for guidance on hypertension treatment

Universal Protocol Section



Suspected Stroke

Complete the Brunswick County EMS Acute Stroke Activation Criteria and Assessment Form.

Leave a copy at the receiving facility. Attach a copy to your PCR.

Notify receiving facility by stating "Code Stroke Activation". Document this with time stamp in ePCR in Alert Type by selecting "Acute Stroke".

Pediatric Stroke: transport to NHRMC or GSRMC.

Pearls

- **Recommended Exam: Mental Status, HEENT, Heart, Lungs, Abdomen, Extremities, Neuro**
- **Items in Red Text are key performance measures used in the EMS Acute Stroke Care Toolkit.**
- **Acute Stroke care is evolving rapidly. Time of Onset/ Last Seen Normal may be changed at any time depending on the capabilities and resources of your regional hospital(s).**
- **Refer to your Stroke: EMS Triage and Destination Plan which should be updated when community resources change.**
- **Time of Onset or Last Seen Normal:**
 - **One of the most important items the pre-hospital provider can obtain, of which all treatment decisions are based.**
 - **Be very precise in gathering data to establish the time of onset and report as an actual time (i.e. 13:47 NOT "about 45 minutes ago.")**
 - **Without this information patient may not be able to receive thrombolytics at facility.**
 - **Wake up stroke: Time starts when patient last awake or symptom free.**
- **Time of Symptom Discovery:**
 - **Time when symptoms of stroke are first noticed by patient, bystanders, witnesses, or family/ caregivers.**
- **Sources of information pertaining to Last Known Well Time or Symptoms Onset:**
 - **You are often in the best position to determine the actual Time of Onset while you have family, friends or caretakers available.**
 - **Often these sources of information may arrive well after you have delivered the patient to the hospital. Delays in decisions due to lack of information may negatively impact patient care.**
 - **Obtain contact information (phone number and name) of best witnesses and give to hospital providers.**
- **The Reperfusion Checklist should be completed for any suspected stroke patient as time allows.**
- **If possible place 2 IV sites, preferably above the wrists, and if possible both in the left upper extremity.**
- **Blood Draw:**
 - **Many stroke centers utilize EMS venous blood samples. Follow your local policy and procedures.**
 - **The differential listed in the UP 4 Altered Mental Status Protocol should also be considered.**
 - **Be alert for airway problems (swallowing difficulty, vomiting/aspiration).**
 - **Hypoglycemia can present as a localized neurologic deficit, especially in the elderly.**
 - **Document the EMS Stroke Screen, Stroke Severity Score, and Stroke Alert notification time in the ePCR or PCR.**
 - **Agencies may use validated pre-hospital stroke screen of choice.**
- **Pediatrics:**
 - **Strokes do occur in children, they are slightly more common in ages < 2, in boys, and in African-Americans.**
 - **Newborn and infant symptoms consist of seizures, extreme sleepiness, and using only one side of the body.**
 - **Children and teenagers symptoms may consist of severe headaches, vomiting, sleepiness, dizziness, and/or loss of balance or coordination.**



Suspected Sepsis

History

- Duration and severity of fever
- Past medical history
- Medications/ Recent antibiotics
- Immunocompromised (transplant, HIV, diabetes, cancer)
- Indwelling medical device
- Last acetaminophen or ibuprofen
- Recent Hospital/ healthcare facility
- Bedridden or immobile
- Elderly and very young – at risk
- Prosthetic device / indwelling device

Signs and Symptoms

- Warm
- Flushed
- Sweaty
- Chills/ Rigors
- Delayed cap refill
- Mental status changes

Associated Symptoms (Helpful to localize source)

- myalgias, cough, chest pain, headache, dysuria, abdominal pain, rash

Differential

- Infections: UTI, Pneumonia, skin/ wound
- Cancer/ Tumors/ Lymphomas
- Medication or drug reaction
- Connective tissue disease: Arthritis, Vasculitis
- Hyperthyroidism
- Heat Stroke
- Meningitis
- Hypoglycemia/hypothermia
- MI/ CVA

Consider: Contact, Droplet, and Airborne Precautions	
Temperature Measurement Procedure <i>if available</i>	
	Fever/ Infection Control Protocol UP 10 <i>if needed</i>
	Altered Mental Status Protocol UP 4 <i>if needed</i>
B	12 Lead ECG Procedure
	IV or IO Access Protocol UP 6 <i>If indicated</i>
P	Cardiac Monitor

Age Specific Blood Pressure indicating possible shock

Age 0 – 28 days: SBP < 60
 Ages ≥ 1 month: SBP < 70
 Age 1 – 9: SBP < 70 + (2x Age)

Ages 10 – 64: SBP < 90
 Ages ≥ 65: SBP < 110

All ages Shock Index:
 HR > SBP

Exit to
 Age Appropriate
 Condition Appropriate
 Protocol(s)

Sepsis Screen Positive

SEPSIS ALERT
 Notify Receiving Facility Immediately

MAP (Mean Arterial Pressure)

SBP + 2(DBP)
3

Monitor usually calculates this value on screen

Adult SIRS Criteria

+ SUSPECTED INFECTION

Temperature ≥ 100.4° F (38° C)
 Or ≤ 96.8° F (36° C)
AND
 Any 1 of the following:
 HR > 90
 RR > 20
 EtCO < 25 mmHg

Pediatrics SIRS Criteria

Temperature Same as adult

AND

Heart Rate
 1 month – 1 year > 180
 2 – 5 years > 140
 6 – 12 years > 130
 13 – 18 years > 120

A

Venous Access Blood Draw *if supplies available*

Lactated Ringers 30mL/kg
 Titrate SPB ≥ 90 mmHg MAP > 65 mmHg
Maximum 2 L
Peds: 20 mL/kg IV / IO
 Repeat to titrate Age Appropriate SBP ≥ 70 + 2 x Age
Maximum 60 mL/kg

Rocephin IV / IO (age ≥ 18 years)
50 mg /kg up to 2 grams over 10 minutes (to adm IM, see page 2)

P

Epinephrine 2-10 mcg/min
 on infusion to goal of **MAP>65**

Age Appropriate Hypotension/ Shock Protocol AM 5/ PM 3

Notify Destination or Contact Medical Control

Universal Protocol Section



Suspected Sepsis

Sepsis vs. Septic Shock:

Sepsis - organ injury or damage in response to infection

Septic Shock - occurs when the response to the infection leads to dangerously low blood pressure and abnormalities in cellular metabolism.

Rocephin reconstitution and administration:

- 1. Mix w/ 3mL of NS.**
- 2. Roll vial to dissolve the powder. DO NOT SHAKE**

IV / IO: Mix the reconstitution with 100mL NS and give over 10 minutes

IM: administer reconstitution deep gluteal

Pearls

- **Recommended Exam: Mental Status, Skin, HEENT, Neck, Heart, Lungs, Abdomen, Back, Extremities, Neuro**
- **Recommended Exam Pediatrics: In childhood, physical assessment reveals important clues for sepsis. Look for mental status abnormalities such as anxiety, restlessness, agitation, irritability, confusion, or lethargy. Cardiovascular findings to look for include cool distal extremities, capillary refill >3 seconds, or mottled skin.**
- **Sepsis is a life threatening condition where the body's immune response to infection injures its own tissues and organs.**
- **Severe sepsis is a suspected infection with 2 or more SIRS criteria (or qSOFA) along with organ dysfunction, such as AMS, hypotension, or hypoxia.**
- **Septic shock is severe sepsis and poor perfusion unimproved after fluid bolus.**
- **Agencies administering antibiotics should inquire about drug allergies specific to antibiotics or family of antibiotics.**
- **Following each fluid bolus, assess for pulmonary edema. Consider administration of agency specific vasopressor.**
- **Supplemental oxygen should be given and titrated to oxygenation saturation $\geq 92\%$.**
- **EKG should be obtained with suspected sepsis, but should not delay care in order to obtain.**
- **Abnormally low temperatures increase mortality and are found often in geriatric patients.**
- **Quantitative waveform capnography can be a reliable surrogate for lactate monitoring in detecting metabolic distress in sepsis patients. EtCO₂ < 25 mm Hg are associated with serum lactate levels > 4 mmol/L.**
- **Patients with a history of liver failure should not receive acetaminophen.**
- **Droplet precautions** include standard PPE plus a standard surgical mask for providers who accompany patients in the back of the ambulance and a surgical mask or NRB O2 mask for the patient. This level of precaution should be utilized when influenza, meningitis, mumps, streptococcal pharyngitis, and other illnesses spread via large particle droplets are suspected. A patient with a potentially infectious rash should be treated with droplet precautions.
- **Airborne precautions** include standard PPE plus utilization of a gown, change of gloves after every patient contact, and strict hand washing precautions. This level of precaution is utilized when multi-drug resistant organisms (e.g. MRSA), scabies, or zoster (shingles), or other illnesses spread by contact are suspected.
- **All-hazards precautions** include standard PPE plus airborne precautions plus contact precautions. This level of precaution is utilized during the initial phases of an outbreak when the etiology of the infection is unknown or when the causative agent is found to be highly contagious (e.g. **SARS, SARS-CoV-2, COVID-19, MERS, Monkeypox**).
- **Allergies to NSAIDs (non-steroidal anti-inflammatory medications) are a contraindication to Ibuprofen.**
- **Agency Medical Director may require contact of medical control prior to EMT / MR administering any medication.**
- **Sepsis Screen:**
Adult / Pediatric Systemic Inflammatory Response Syndrome (SIRS) criteria



Syncope

History

- Cardiac history, stroke, seizure
- Occult blood loss (GI, ectopic)
- Females: LMP, vaginal bleeding
- Fluid loss: nausea, vomiting, diarrhea
- Past medical history
- Medications

Signs and Symptoms

- Loss of consciousness with recovery
- Lightheadedness, dizziness
- Palpitations, slow or rapid pulse
- Pulse irregularity
- Decreased blood pressure

Differential

- Vasovagal
- Orthostatic hypotension
- Cardiac syncope
- Micturition / Defecation syncope
- Psychiatric
- Stroke
- Hypoglycemia
- Seizure
- Shock (see Shock Protocol)
- Toxicological (Alcohol)
- Medication effect (hypertension)
- PE
- AAA


	Age Appropriate Airway Protocol(s) AR 1, 2, 3, 5, 6 if indicated
	Blood Glucose Analysis Procedure
B	12 Lead ECG Procedure
	IV or IO Access Protocol UP 6
P	Cardiac Monitor
	Altered Mental Status Protocol UP 4 if indicated
	Age Appropriate Cardiac Protocol(s) if indicated
	Age Appropriate Hypotension/ Shock Protocol AM 5/ PM 3 if indicated
	Multiple Trauma Protocol TB 6 Spinal Motion Restriction Procedure/ Protocol TB 8 if indicated

**Age Specific Blood Pressure
indicating possible shock**

Age 0 – 28 days: SBP < 60
Ages ≥ 1 month: SBP < 70
Age 1 – 9: SBP < 70 + (2x Age)

Ages 10 – 64: SBP < 90
Ages ≥ 65: SBP < 110

**All ages Shock Index:
HR > SBP**





Syncope

Pearls

- **Recommended Exam: Mental Status, Skin, HEENT, Heart, Lungs, Abdomen, Back, Extremities, Neuro**
- **Syncope is both loss of consciousness and loss of postural/ muscle tone with collapse. Symptoms preceding the event are important in determining etiology.**
- **Syncope often is due to a benign process but can be an indication of serious underlying disease in both the adult and pediatric patient.**
- **Often patients with syncope are found normal on EMS evaluation. In general patients experiencing syncope require cardiac monitoring and emergency department evaluation.**
- **Differential should remain wide and include:**

Cardiac arrhythmia	Neurological problem	Choking	Pulmonary embolism
Hemorrhage	Stroke	Respiratory	Hypo or Hyperglycemia
GI Hemorrhage	Seizure	Sepsis	
- **High-risk patients:**

Age \geq 60	Syncope with exertion
History of CHF	Syncope with chest pain
Abnormal ECG	Syncope with dyspnea
- **Abdominal/ back pain in women of childbearing age should be treated as pregnancy related until proven otherwise.**
- **The diagnosis of abdominal aneurysm should be considered with abdominal pain, with or without back and/ or lower extremity pain or diminished pulses, especially in patients over 50 and/ or patients with shock/ poor perfusion. Notify receiving facility early with suspected abdominal aneurysm.**
- **Consider cardiac etiology in patients > 35, diabetics, and/ or women especially with upper abdominal complaints.**
- **Heart Rate: Tachycardia is one of the first clinical signs of dehydration, typically increases as dehydration becomes more severe.**
- **Syncope with no preceding symptoms or event may be associated with an arrhythmia.**
- **Assess for signs and symptoms of trauma if associated or questionable fall with syncope.**
- **Consider dysrhythmias, GI bleed, ectopic pregnancy, and seizure as possible causes of syncope.**
- **In general these patients should be transported: Patients who experience syncope associated with headache, neck pain, chest pain, abdominal pain, back pain, dyspnea, or dyspnea on exertion need prompt medical evaluation.**
- **More than 25% of geriatric syncope is cardiac dysrhythmia based.**



Behavioral Health Crisis

History


- Situational crisis
- Psychiatric illness/medications
- Injury to self or threats to others
- Medic alert tag
- Substance abuse / overdose
- Diabetes

Signs and Symptoms

- Anxiety, agitation, confusion
- Affect change, hallucinations
- Delusional thoughts, bizarre behavior
- Combative violent
- Expression of suicidal / homicidal thoughts

Differential

- Altered Mental Status
- Alcohol Intoxication
- Toxin / Substance abuse
- Medication effect / overdose / withdrawal
- Depression
- Bipolar (manic-depressive)
- Schizophrenia
- Anxiety disorders



Universal Protocol Section



Behavioral Health Crisis

Refusal Checklist

- o Is the patient >17 y/o/a or an emancipated minor?
- o Can the patient retain and comprehend relevant information?
- o Can the patient believe information?
- o Can the patient use information to make a choice?
- o Is the patient NOT DANGEROUS to self or others (i.e. no suicidal or homicidal ideation)?

If all are "YES" then the patient has the capacity to decline further care/ transport.
 If any are "NO" then the patient does not have capacity to make his/her own medical decisions. Document these concepts clearly in your assessment - simply stating "alert and oriented" is not sufficient.
 Do not hesitate to contact medical control if there are questions about capacity.


Pearls

- **Recommended Exam: Mental Status, Skin, Heart, Lungs, Neurologic status**
- **Crew/ responders safety is the main priority. Call for assistance, stage, or withdraw from scene if necessary.**
- **Law Enforcement:**
Any patient who is handcuffed or restrained by Law Enforcement and transported by EMS, must be accompanied by law enforcement during transport.
Patient should not be transported with upper extremities hand-cuffed behind back as this prevents proper assessment and could lead to injury.
Consider multidisciplinary coordination with law enforcement to approach verbal de-escalation, restraint, and/ or USP 6 Restraints: Therapeutic Take-down Procedure.
- **Maintain high-index of suspicion for underlying medical or traumatic disorder causing or contributing to behavioral disturbance. Medical causes more likely in ages < 12 or > 40.**
- **General communications techniques**
Ask Open-ended questions (questions that cannot be answered with a yes/no)
"Tell me how we can help you?" "What caused you to call 911 today?"
Active listening (stay engaged, be able to summarize patient's story, use your body language to convey listening)
Eye contact, nodding your head, periodically repeating back part of patient's story
Encouraging (remain positive, convey interest in patient's crisis)
"Tell me more about that..."
Clarifying questions (ask patient to rephrase or repeat if you don't understand)
"I'm not sure I understand, can you...?"
Emotional labeling (naming emotions patient is demonstrating, validating emotions)
"You look upset." "You seem angry."
Conversational pause (okay to allow a period of silence for patient to process information)
- **Behavioral health disturbance incidents are increasing and commonly involve the following:**

Substance misuse	Psychosis
Depression/ Anxiety/ Stress Reactions / Bipolar	Schizophrenia or schizophrenia-like illness
- **Restraints:**
All patients who receive either physical or chemical restraint must be continuously observed by ALS personnel on scene or immediately upon their arrival.
Do not position or transport any restrained patient in such a way that could impact the patient's respiratory or circulatory status (i.e. prone position)
- **Maintain high-index of suspicion for medical, trauma, abuse, or neglect causes:**
 Hypoglycemia, hyperglycemia, overdose, substance abuse, hypoxia, head injury, shock, sepsis, stroke, etc.
 Domestic violence, child or geriatric abuse/ neglect.
- **Extrapyramidal reactions:**
 Condition causing involuntary muscle movements or spasms typically of the face, neck and upper extremities. May present with contorted neck and trunk with difficult motor movements. Typically an adverse reaction to antipsychotic drugs like Haloperidol and may occur with your administration. When recognized, give **Diphenhydramine 50 mg IV / IO / IM / PO** in adults or **1 mg/kg IV / IO / IM / PO** in pediatrics, **Maximum 50 mg.**
- **May add page 3 to protocol for specific for local mental health and / or substance misuse resources or destinations.**



Behavioral CIT Paramedic (Optional)




This is a continuation of the Evaluation and Screening for Mental Health and Substance Abuse Procedure

Alternative Destinations other than Hospital ED
Call Access Line for all referrals:
Contact Number

If appropriate, contact Mobile Crisis:
Contact Number

- Blood/Vomit/Difficulty Breathing/Acute Medical Issue
- Blood Glucose <70 or >250 & symptomatic
- Head Trauma or fall in past 7 days
- In and out of consciousness
- Seizure activity in past 24 hours
- Unable to speak or walk
- On IVC or Emergency Custody
- Violent or Aggressive
- Provider can't see within 2 hours at alternative site
- Acute Withdrawal
- Hospital transport requested by patient
- BP ≥210/130 or ≥180/110 and symptomatic
- BAC > .35
- Oxygen Dependent




NO

- Suicidal (with plan and/or intent)
- Homicidal Ideation
- Psychosis (auditory/visual hallucinations or delusions)
- Request for Detox
- Med Request during 8:00 am and 3:00pm and client can not wait until next day for meds



NO

- No current suicidal ideation
- Anxiety/Panic
- Tearful crying with no suicidal plan or intent
- Med request in which EMS/Mobile Crisis able to triage and client is "ok" to go M-F for meds.




Alternative Destinations / Crisis Providers For Centerpoint

County		
Resource Agency	Resource Agency	Resource Agency
Hours of Operation	Hours of Operation	Hours of Operation




Behavioral Agitation/ Sedation Guide



Universal Protocol Section



Behavioral Hyperactive Delirium With Severe Agitation



- General Guidance**
- Use of protocol is indicated when a medical emergency requires a combination of both physical restraint and chemical restraint in order to prevent imminent injury to a patient and/ or providers.
 - Use of this protocol requires medical judgement and consultation with Medical Control where indicated.
 - Non-medical personnel requests or opinions should not be used as a factor when implementing this protocol.



Behavioral Hyperactive Delirium With Severe Agitation

IM administration of Midazolam is more effective and is the preferred route of administration if IV / IO access is not obtained, but may be administered IN.

Midazolam may be administered at 2-2.5mg IV as opposed to IM when an IV is established.

Pearls

- **Ketamine for sedation purposes:**
Ketamine may be used in pediatric patients who fit within a Pediatric Medication/ Skill Resuscitation System product, ≤ 15 years of age, or ≤ 49 kg) with DIRECT ONLINE MEDICAL ORDER by the system MEDICAL DIRECTOR or ASSISTANT MEDICAL DIRECTOR only.
- **Hyperactive Delirium with Severe Agitation:**
Medical emergency: Combination of delirium, psychomotor agitation, anxiety, hallucinations, speech disturbances, disorientation, violent/ bizarre behavior, insensitivity to pain, hyperthermia and increased strength.
Potentially life-threatening and associated with use of physical control measures, including physical restraints.
Most commonly seen in male subjects with a history of serious mental illness and/or acute or chronic drug abuse, particularly stimulant drugs such as cocaine, crack cocaine, methamphetamine, amphetamines or similar agents.
Alcohol or substance withdrawal as well as head trauma may also contribute to the condition.
- **Restraint use:**
Physical restraints are not contraindicated in agitated or excited delirium, but you must use caution. Once sedated, prevent patient from continued struggle, which can worsen metabolic condition. Prevent patient from assuming a prone position for prolonged period, move to supine position as quickly as possible.
Team approach for sedation and Restraint Therapeutic Take Down Procedure USP-6:
 - 1 provider for each limb.
 - 1 provider to lead restraint, maintain airway and control head.
 - 1 Provider to administer medication.Do not position prone or prone with restraints, as this can impede respiration and ventilation.
- Hyperthermia: Assess for and treat hyperthermia.



Sickle Cell Crisis

History

- Past medical history
- Medications
- Recent illness
- Prior pain crisis location
- Pain regimen at home

Signs and Symptoms

- Pain
- One sided paralysis / weakness
- Difficulty walking / speaking
- Sudden vision changes
- Unexplained numbness
- Severe headache
- Fever
- SOB
- Chest Pain
- Abdominal Pain
- Pallor

Differential

- Sickle Celle Pain Crisis
- Aplastic Crisis
- Acute Chest Syndrome
- Alcohol / drug use
- Toxic ingestion
- Seizure
- Stroke
- Altered baseline mental status
- Sepsis
- Pneumonia

Apply Hot Packs to affected areas: especially joints and areas of increased pain

Provide emotional support
Calm and continual reassurance

B	Blood Glucose Analysis as needed Assess Pain Severity 12 Lead ECG Procedure
	IV or IO Access Protocol UP 6
P	Cardiac Monitor
	Altered Mental Status Protocol UP 4 <i>if indicated</i>
	Manage Airway as appropriate Protocol AR 1 & AR 5 <i>if indicated</i>
	Suspected Stroke Protocol UP 14 <i>if indicated</i>
	Suspected Seizure Protocol UP 13 <i>if indicated</i>
	Hypotension/ Shock Protocol AM 5 & PM 3 <i>if indicated</i>

Fluids indicated if signs of shock or hypotensive

Sickle Cell Crisis
- Acute Chest Syndrome
- Abdominal Crisis
- Joint Crisis

YES

Pain Control
Protocol UP 11

Notify Destination or Contact Medical Control



Sickle Cell Crisis

PEARLS

Patterns of an acute sickle cell crisis are now recognizable. They are based on the part of the body where the crisis occurs.

Is this their typical pain crisis? If not, what is different about it?

Any fever, SOB, pleuritic chest pain?

Acute chest syndrome:

Sudden acute chest pain with coughing up of blood can occur. Low-grade fevers can be present. The person is usually short of breath. If a cough is present, it often

is nonproductive. Acute chest syndrome is common in a young person with sickle cell disease.

Chronic (long-term) sickle cell lung disease develops over time because the acute and subacute lung crisis leads to scarred lungs as well as other problems.

Abdominal crisis:

The pain associated with the abdominal crisis of sickle cell disease is constant and sudden. It becomes unrelenting. The pain may or may not be localized to any one area of the abdomen. Nausea, vomiting, and diarrhea may or may not occur.

Joint crisis:


Acute and painful joint crisis may develop without a significant traumatic history. Its focus is either in a single joint or in multiple joints. Often the connecting bony parts of the joint are painful. Range of motion is often restricted because of the pain. Avascular necrosis of the hips can occur, causing permanent damage.

Stroke:

Many sickle cell patients can present with strokes at a younger age than average. Ensure hospital pre-notification indicating it is a sickle cell patient as the standard of care for a Sickle Cell Stroke can involve exchange therapy as opposed to other treatments such as tPA.



Adult Airway



Capnography Monitoring

- End-tidal (EtCO₂) monitoring is mandatory following placement of an endotracheal tube.
- EtCO₂ monitoring is mandatory following placement of a BIAD once available on scene.

Protocols AR 1, 2, and 3 should be utilized together (even if agency is not using Drug Assisted Airway Protocol) as they contain useful information for airway management.

Failed Airway:
Unable to Ventilate and Oxygenate ≥ 90% during or after one (1) or more unsuccessful intubation attempts
and/ or Anatomy inconsistent with continued attempts
and/ or Three (3) unsuccessful attempts by most experienced Paramedic/ AEMT.
Exit to **Adult Failed Airway Protocol AR 2**



Adult Airway

A c-collar should be placed, especially when moving the patient, to better maintain ETT placement.

Pearls

- See Pearls section of protocols AR 2 and 3.
- For the purposes of this protocol a secure airway is when the patient is receiving appropriate oxygenation and ventilation.
- If an effective airway is being maintained by BVM with continuous pulse oximetry values of $\geq 90\%$, it is acceptable to continue with basic airway measures.
- Ventilation rate should be 10 - 12 per minute to maintain a EtCO₂ of 35 – 45 and avoid hyperventilation.
- **Anticipating the Difficult Airway and Airway Assessment**
 - **Difficult BVM Ventilation (ROMAN):** Radiation treatment/ Restriction; Obese/ Obstruction/ OB – 2d and 3d trimesters/ Obstructive sleep apnea; Mask seal difficulty (hair, secretions, trauma); Age ≥ 55 ; No teeth.
 - **Difficult Laryngoscopy (LEON):** Look externally for anatomical problems; Evaluate 3-3-2 (Mouth opening should equal 3 of patients finger's width, mental area to neck should equal 3 of patient's finger's width, base of chin to thyroid prominence should equal 2 of patients finger's width); Obese, obstruction, OB – 2d and 3d trimesters; Neck mobility limited.
 - **Difficulty BIAD (RODS):** Radiation treatment/ Restriction; Obese/ Obstruction/ OB – 2d and 3d trimesters/ Obstructive sleep apnea; Distorted or disrupted airway; Short thyromental distance/ Small mandible.
 - **Difficulty Cricothyrotomy / Surgical Airway (SMART):** Surgery scars; Mass or hematoma, Access or anatomical problems; Radiation treatment to face, neck, or chest; Tumor.
- Complete an Airway Evaluation Form with any BIAD or Intubation procedure where medications are used to facilitate.
- **Nasotracheal intubation:**
 - Procedure requires spontaneous breathing and may require considerable time, exposing patient to critical desaturation.
 - Contraindicated in combative, anatomically disrupted or distorted airways, increased ICP, severe facial trauma, basal skull fracture, and head injury. Orotracheal route is preferred.
- Intubation attempt defined as laryngoscope blade passing the teeth or endotracheal tube passed into the nostril.
- If First intubation attempt fails, make an adjustment and try again: (Consider change of provider in addition to equipment).
- AEMT and Paramedics should consider using a BIAD if oral-tracheal intubation is unsuccessful.
- During intubation attempts use External Laryngeal Manipulation to improve view of glottis.
- Gastric tube placement should be placed in all intubated patients if available or time allows.
- It is important to secure the endotracheal tube well to better maintain ETT placement. Manual stabilization of endotracheal tube should be used during all patient moves / transfers.
- **DOPE:** Displaced tracheostomy tube / ETT, Obstructed tracheostomy tube / ETT, Pneumothorax and Equipment failure.



Adult, Failed Airway

Definition of Failed Airway:
 Unable to Ventilate and Oxygenate $\geq 90\%$ during or after one (1) or more unsuccessful intubation attempts
 and/ or
 Anatomy inconsistent with continued attempts
 and/ or
 Three (3) unsuccessful attempts by most experienced Paramedic/AEMT.
Each attempt should include change in approach or equipment
 NO MORE THAN THREE (3) ATTEMPTS TOTAL

- Capnography Monitoring**
- End-tidal (EtCO₂) monitoring is mandatory following placement of an endotracheal tube.
 - EtCO₂ monitoring is mandatory following placement of a BIAD once available on scene.

Protocols AR 1, 2, and 3 should be utilized together (even if agency is not using Drug Assisted Airway as they contain useful information for airway management).

Failed Airway

Call for additional resources if available

BVM
 Adjunctive Airway NPA/ OPA
 Maintains
 Oxygen Saturation $\geq 92\%$

Continue BVM
 Supplemental Oxygen
 Exit to
 Appropriate Protocol(s)

NO

B	Attempt Airway Blind Insertion Airway Device Procedure
A	Airway Video Laryngoscopy Device Procedure <i>if available</i> Optional
P	Airway Cricothyrotomy Surgical Procedure
	Supplemental oxygen BVM with Airway Adjuncts Maintain Oxygen Saturation $\geq 92\%$
	Post-intubation BIAD Management Protocol AR 8

Notify Destination or Contact Medical Control



Adult, Failed Airway

A c-collar should be placed, especially when moving the patient, to better maintain ETT placement.

Pearls

- **For the purposes of this protocol a secure airway is when the patient is receiving appropriate oxygenation and ventilation.**
- **If an effective airway is being maintained by BVM with continuous pulse oximetry values of $\geq 90\%$, it is acceptable to continue with basic airway measures.**
- **Ventilation rate should be 10 - 12 per minute to maintain a EtCO₂ of 35-45 and avoid hyperventilation.**
- **Anticipating the Difficult Airway and Airway Assessment**
 - Difficult BVM Ventilation (ROMAN):** Radiation treatment/ Restriction; **Obese/ Obstruction/ OB** – 2d and 3d trimesters/ Obstructive sleep apnea; **Mask seal difficulty** (hair, secretions, trauma); **Age ≥ 55 ; No teeth.**
 - Difficult Laryngoscopy (LEON):** Look externally for anatomical problems; Evaluate 3-3-2 (Mouth opening should equal 3 of patients finger's width, mental area to neck should equal 3 of patient's finger's width, base of chin to thyroid prominence should equal 2 of patients finger's width); **Obese, obstruction, OB** – 2d and 3d trimesters; **Neck mobility limited.**
 - Difficulty BIAD (RODS):** Radiation treatment/ Restriction; **Obese/ Obstruction/ OB** – 2d and 3d trimesters/ Obstructive sleep apnea; **Distorted or disrupted airway; Short thyromental distance/ Small mandible.**
 - Difficulty Cricothyrotomy / Surgical Airway (SMART):** Surgery scars; **Mass or hematoma, Access or anatomical problems; Radiation treatment to face, neck, or chest; Tumor**
- **Complete an Airway Evaluation Form with any BIAD or Intubation procedure where medications are used to facilitate.**
- **Nasotracheal intubation:**
 - Procedure requires spontaneous breathing and may require considerable time, exposing patient to critical desaturation.**
 - Contraindicated in combative, anatomically disrupted or distorted airways, increased ICP, severe facial trauma, basal skull fracture, and head injury. Orotracheal route is preferred.**
- **Intubation attempt defined as laryngoscope blade passing the teeth or endotracheal tube passed into the nostril.**
- **If First intubation attempt fails, make an adjustment and try again: (Consider change of provider in addition to equipment)**
- **AEMT and Paramedics should consider using a BIAD if oral-tracheal intubation is unsuccessful.**
- **During intubation attempts use External Laryngeal Manipulation to improve view of glottis.**
- **Gastric tube placement should be placed in all intubated patients if available or time allows.**
- **It is important to secure the endotracheal tube well to better maintain ETT placement. Manual stabilization of endotracheal tube should be used during all patient moves/ transfers.**
- **DOPE: Displaced tracheostomy tube/ ETT, Obstructed tracheostomy tube/ ETT, Pneumothorax and Equipment failure.**



Airway, Drug Assisted (OPTIONAL)

Indications for Drug Assisted Airway
 Failure to protect the airway and/or
 Unable to oxygenate and/or
 Unable to ventilate and/or
 Impending airway compromise

Capnography Monitoring

- End-tidal (EtCO₂) monitoring is mandatory following placement of an endotracheal tube.
- EtCO₂ monitoring is mandatory following placement of a BIAD once available on scene.

Protocols AR 1, 2, 3, 5, and 6 should be utilized together (even if agency is not using Drug Assisted Airway Protocol) as they contain useful information for airway management.

Preoxygenate 100% O₂
 IV or IO Access Protocol UP 6
2 points of access
 Assemble Airway Equipment
 Suction equipment
 Alternative Airway Device

Hypoxic Or
 Hypotensive Or
 Dangerously Combative?

Ketamine 2 mg/kg IV / IO
No IV or IO Access:
Ketamine 4 mg/kg IM
Maximum 400 mg
Pediatric: See Pearls
2 mg/kg IM
Maximum 400 mg
 Correct Hypoxia and/ or Hypotension
 Age Appropriate Airway Protocol(s) AR 1, 2, 5, 6, as indicated
 Hypotension/ Shock Protocol AM 5/ PM 3 as indicated

Etomidate 0.3 mg/kg IV / IO
Or
Ketamine 2 mg/kg IV / IO
May repeat x 1
Succinylcholine 2 mg/kg IV / IO
Or
Rocuronium 1.5 mg/kg IV / IO
(if Succinylcholine contraindicated)
May repeat x 1
Intubate trachea
Placement Verified
Continuous Waveform Capnography

Consider Restraints Physical Procedure
 Consider Gastric Tube Insertion Procedure

Awakening or Moving after intubation

Exit to Post-intubation/ BIAD Management Protocol AR 8

Exit to Appropriate Protocol(s)

Procedure will remove patient's protective airway reflexes and ability to breath.
 You must be sure of your ability to intubate before beginning this procedure.
 Must have two (2) Paramedics on scene

Red Text
 Key performance indicators used to evaluate protocol compliance.
 A NCOEMS Airway Evaluation Form must be completed for every patient who receives Drug Assisted Airway.

Airway Respiratory Protocol Section



Airway, Drug Assisted (OPTIONAL)

Pearls

- Agencies must maintain a separate Performance Improvement Program specific to Drug Assisted Airway.
- This procedure requires at least 2 Paramedics. See Pearls section of protocols AR 1 and 2.
- For the purposes of this protocol, a secure airway is when the patient is receiving appropriate oxygenation and ventilation.
- If an effective airway is being maintained by BVM with continuous pulse oximetry values of $\geq 90\%$, it is acceptable to continue with basic airway measures.
- **Ventilation rate:**
30 for Neonates, 25 for Toddlers, 20 for School Age, and for Adolescents the normal Adult rate of 10 - 12 per minute. Maintain EtCO₂ between 35 - 45 and avoid hyperventilation.
- **Hypoxia and/ or Hypotension:**
Increased risk of cardiac arrest when a sedative with paralytic medications are administered while hypoxic and/ or hypotensive. Resuscitation and correction of hypoxia and/ or hypotension are paramount prior to use of these combined agents. Ketamine administration allows time for appropriate resuscitation of hypoxia and/or hypotension while managing the airway.
- **PEDIATRIC POPULATION (fit within a Pediatric Medication/ Skill Resuscitation System product, ≤ 15 years of age, or ≤ 49 kg):**
This protocol may be used in the pediatric population ONLY with real time, direct online medical control by the system MEDICAL DIRECTOR or ASSISTANT MEDICAL DIRECTOR.
Agencies using Ketamine in the pediatric population must also be using in their adult population.
- **KETAMINE:**
Ketamine may be used with or without a paralytic agent in conjunction with either an OPA, NPA, BIAD, or endotracheal tube. (BIAD is preferred over endotracheal tube until hypoxia and/ or hypotension are corrected).
Ketamine may be used during the resuscitation of hypoxia or hypotension in conjunction with airway management. Once hypoxia and hypotension are corrected, use of a sedative and paralytic can proceed if indicated.
Ketamine may be used in the dangerously combative patient requiring airway management IM. IV/ IO should be established as soon as possible.
Ketamine may be used for sedation once a BIAD or endotracheal tube are established and confirmed.
Agencies using Ketamine must follow Standards Policy: Medial Policy Section Ketamine Program Requirements. Medical Policy 2.
- Intubation attempt defined as laryngoscope blade passing the teeth or endotracheal tube passed into the nostril.
- If First intubation attempt fails, make an adjustment and try again: (Consider change of provider in addition to equipment)
- **NC EMS Airway Evaluation Form:**
Fully complete and have receiving healthcare provider sign confirming BIAD or endotracheal tube placement.
Complete online in region specific *ReadyOp* and upload completed form.
Complete when Ketamine, Etomidate, Succinylcholine and/ or Rocuronium or used to facilitate use of a BIAD and/ or endotracheal intubation.
- Paramedics/ AEMT should consider using a BIAD if endotracheal intubation is unsuccessful.
- Drug Assisted Airway is not recommended in an urban setting (short transport) when able to maintain oxygen saturation $\geq 90\%$.
- **DOPE:** Displaced tracheostomy tube/ ETT, Obstructed tracheostomy tube/ ETT, Pneumothorax and Equipment failure.



Adult COPD/ Asthma Respiratory Distress

History


- Asthma; COPD -- chronic bronchitis, emphysema, congestive heart failure
- Home treatment (oxygen, nebulizer)
- Medications (theophylline, steroids, inhalers)
- Toxic exposure, smoke inhalation

Signs and Symptoms

- Shortness of breath
- Pursed lip breathing
- Decreased ability to speak
- Increased respiratory rate and effort
- Wheezing, rhonchi
- Use of accessory muscles
- Fever, cough
- Tachycardia

Differential

- Asthma
- Anaphylaxis
- Aspiration
- COPD (Emphysema, Bronchitis)
- Pleural effusion
- Pneumonia
- Pulmonary embolus
- Pneumothorax
- Cardiac (MI or CHF)
- Pericardial tamponade
- Hyperventilation
- Inhaled toxin (Carbon monoxide, etc.)





Adult COPD/ Asthma Respiratory Distress

Known or Suspected COVID

Terbutaline can be administered alone without Albuterol / Atrovent if COVID is suspected or known. This decision is left up to the Provider.

If Albuterol / Atrovent are administered a N95 mask or P100 mask **MUST** be worn by the Provider.

Dosage:

Adult: 0.25 mg SQ, injected in lateral deltoid. May repeat dosage every 15 minutes as needed.


Pediatric: 0.01 mg/kg (up to 0.25mg) SQ. May repeat dosage every 15 minutes as needed.

Pearls

- **Recommended Exam: Mental Status, HEENT, Skin, Neck, Heart, Lungs, Abdomen, Extremities, Neuro**
- **Items in Red Text are key performance measures used to evaluate protocol compliance and care.**
- **This protocol includes all patients with respiratory distress, COPD, Asthma, Reactive Airway Disease, or bronchospasm.**
- **Patients may also have wheezing and respiratory distress with viral upper respiratory tract infections and pneumonia.**
- Pulse oximetry should be monitored continuously and consider End-tidal CO₂ monitoring if available.
- **Combination nebulizers containing albuterol and ipratropium (DuoNeb):**
 - Patients may require more than 3 nebulizer treatments, treatments should continue until improvement.
 - Following 3 combination nebulizers (DuoNeb), it is preferable to continue albuterol solely with subsequent treatments as there is no proven benefit to continual use of ipratropium.
- **Epinephrine:**
 - If allergic reaction or anaphylaxis is suspected, give immediately and repeat until improvement.
 - If allergic reaction is not suspected, administer with failure to improve and/ or impending respiratory failure.
- Consider Magnesium Sulfate with no improvement and/ or impending respiratory failure. Likely more effective with asthmatic exacerbation and less so with COPD exacerbation.
- **Non-Invasive Positive Pressure Ventilation (NIPPV: CPAP or Bi-Level/ BiPap):**
 - May be used with COPD, Asthma, Allergic reactions, and/ or CHF.
 - Consider early in treatment course.
 - Consider removal if SBP remains < 100 mmHg and not responding to other treatments.
- In patients using levalbuterol (Xopenex) you may use Albuterol for the first treatment then use the patient's supply for repeat nebulizers or agency's supply.
- A silent chest in respiratory distress is a pre-respiratory arrest sign.
- **EMR/EMT:**
 - The use of Epinephrine IM is limited to the treatment of anaphylaxis.
 - Administration of diphenhydramine is limited to the oral route only.
- EMT may administer a beta-agonist (Albuterol) to patients currently prescribed the medication and patients who are not currently prescribed the medication.



Pediatric Airway





Pediatric Airway

Ketamine:

can ONLY be given in the pediatric population for hypoxia/ hypotension/ dangerously combative IF:

1. age >15 years old OR weight >49 kg OR
2. Direct order from Medical Director - Dr. Sherrod

Pearls

This protocol is for use in patients who FIT within a Pediatric Medication/ Skill Resuscitation System Product.

- For the purposes of this protocol, a secure airway is when the patient is receiving appropriate oxygenation and ventilation.
- If an effective airway is being maintained by BVM with continuous pulse oximetry values of $\geq 90\%$, it is acceptable to continue with basic airway measures.
- Ventilation rate:
30 for Neonates, 25 for Toddlers, 20 for School Age, and for Adolescents the normal Adult rate of 10 - 12 per minute. Maintain EtCO₂ between 35 - 45 and avoid hyperventilation.
- Ketamine for airway intervention and/ or sedation purposes:
Ketamine may be used in pediatric patients (fit within a Pediatric Medication/Skill Resuscitation System product, ≤ 15 years of age, or ≤ 49 kg) with DIRECT ONLINE MEDICAL ORDER by the system MEDICAL DIRECTOR or ASSISTANT MEDICAL DIRECTOR only.
Agencies using Ketamine in the pediatric population must also be using in their adult population.
- KETAMINE:
Ketamine may be used with or without a paralytic agent in conjunction with either an OPA, NPA, BIAD or endotracheal tube. BIAD is preferred over endotracheal tube until hypoxia and/ or hypotension are corrected.
Ketamine may be used during the resuscitation of hypoxia or hypotension in conjunction with airway management. Once hypoxia and hypotension are corrected, use of a sedative and paralytic can proceed if indicated.
Ketamine may be used in the dangerously combative patient requiring airway management IM. IV/ IO should be established as soon as possible.
Ketamine may be used for sedation once a BIAD or endotracheal tube are established and confirmed.
Agencies using Ketamine must follow Standards Policy: Medial Policy Section Ketamine Program Requirements. Medical Policy 2.
- Intubation:
Attempt defined as laryngoscope blade passing the teeth or endotracheal tube passed into the nostril.
Use of a stylet is recommended in all pediatric intubations.
Endotracheal tube: Depth = 3 x the diameter of the ETT. Estimated Size = 16 + age (years) / 4. Term newborn = 3.5 mm.
If First intubation attempt fails, make an adjustment and try again: (Consider change of provider in addition to equipment)
- NC EMS Airway Evaluation Form:
Fully complete and have receiving healthcare provider sign confirming BIAD or endotracheal tube placement.
Complete online in region specific *ReadyOp* and upload completed form.
Complete when Ketamine, Etomidate, Succinylcholine and/ or Rocuronium or used to facilitate use of a BIAD and/ or endotracheal intubation. Paramedics/ AEMT should consider using a BIAD if endotracheal intubation is unsuccessful.
- Secure the endotracheal tube well and consider c-collar in pediatric patients (even in absence of trauma) to better maintain ETT placement.
Manual stabilization of endotracheal tube should be used during all patient moves / transfers.
- Airway Cricothyrotomy Percutaneous Needle Procedure:
Indicated as a lifesaving / last resort procedure in pediatric patients < 10 years of age.
Very little evidence to support it's use and safety.
A variety of alternative pediatric airway devices now available make the use of this procedure rare.
Agencies who utilize this procedure must develop a written procedure, establish a training program, maintain equipment and submit procedure and training plan to the State Medical Director/ Regional EMS Office.
 ≥ 10 years: Surgical cricothyrotomy or commercial kits based on agency preference recommended.
- DOPE: Displaced tracheostomy tube/ ETT, Obstructed tracheostomy tube/ ETT, Pneumothorax and Equipment failure.



Pediatric Failed Airway

Definition of Failed Airway:

- Unable to Ventilate and Oxygenate $\geq 90\%$ during or after one (1) or more unsuccessful intubation attempts.
and/ or
- Anatomy inconsistent with continued attempts.
and/ or
- Three (3) unsuccessful attempts by most experienced Paramedic/AEMT.
Each attempt should include change in approach or equipment

NO MORE THAN THREE (3) ATTEMPTS TOTAL

Call for additional resources if available

Failed Airway

BVM Adjunctive Airway NP/ OP
Maintains
Oxygen Saturation $\geq 92\%$

Continue BVM
Supplemental Oxygen

Exit to
Appropriate
Protocol(s)

NO

A Airway Video Laryngoscopy
Device Procedure
if available
Optional

B Attempt
Airway Blind Insertion Airway
Device Procedure

P Airway Cricothyrotomy
Needle Procedure
See Pearls Section

BIAD / Cricothyrotomy
Successful
Or
Oxygenation / Ventilation
Adequate

YES

Exit to
Post-intubation/
BIAD Management
Protocol AR 8


NO

Capnography Monitoring

- End-tidal (EtCO₂) monitoring is mandatory following placement of an endotracheal tube.
- EtCO₂ monitoring is mandatory following placement of a BIAD once available on scene.

Protocols AR 3, 5, and 6 should be utilized together (even if agency is not using Drug Assisted Airway Protocol) as they contain useful information for airway management.

Supplemental oxygen
BVM with Airway Adjuncts
Maintain Oxygen Saturation $\geq 92\%$

 **Notify Destination or Contact Medical Control** 



Pediatric Failed Airway

Ketamine:

can ONLY be given in the pediatric population for hypoxia/ hypotension/ dangerously combative IF:

1. age >15 years old OR weight >49 kg OR
2. Direct order from Medical Director - Dr. Sherrod

Pearls

This protocol is for use in patients who FIT within a Pediatric Medication/ Skill Resuscitation System Product.

- For the purposes of this protocol, a secure airway is when the patient is receiving appropriate oxygenation and ventilation.
- If an effective airway is being maintained by BVM with continuous pulse oximetry values of $\geq 90\%$, it is acceptable to continue with basic airway measures.
- Ventilation rate:
30 for Neonates, 25 for Toddlers, 20 for School Age, and for Adolescents the normal Adult rate of 10 - 12 per minute. Maintain EtCO₂ between 35 - 45 and avoid hyperventilation.
- Ketamine for airway intervention and/ or sedation purposes:
Ketamine may be used in pediatric patients (fit within a Pediatric Medication/Skill Resuscitation System product, ≤ 15 years of age, or ≤ 49 kg) with DIRECT ONLINE MEDICAL ORDER by the system MEDICAL DIRECTOR or ASSISTANT MEDICAL DIRECTOR only.
Agencies using Ketamine in the pediatric population must also be using in their adult population.
- KETAMINE:
Ketamine may be used with or without a paralytic agent in conjunction with either an OPA, NPA, BIAD or endotracheal tube. BIAD is preferred over endotracheal tube until hypoxia and/ or hypotension are corrected.
Ketamine may be used during the resuscitation of hypoxia or hypotension in conjunction with airway management. Once hypoxia and hypotension are corrected, use of a sedative and paralytic can proceed if indicated.
Ketamine may be used in the dangerously combative patient requiring airway management IM. IV/ IO should be established as soon as possible.
Ketamine may be used for sedation once a BIAD or endotracheal tube are established and confirmed.
Agencies using Ketamine must follow Standards Policy: Medial Policy Section Ketamine Program Requirements. Medical Policy 2.
- Intubation:
Attempt defined as laryngoscope blade passing the teeth or endotracheal tube passed into the nostril.
Use of a stylet is recommended in all pediatric intubations.
Endotracheal tube: Depth = 3 x the diameter of the ETT. Estimated Size = 16 + age (years) / 4. Term newborn = 3.5 mm.
If First intubation attempt fails, make an adjustment and try again: (Consider change of provider in addition to equipment)
- NC EMS Airway Evaluation Form:
Fully complete and have receiving healthcare provider sign confirming BIAD or endotracheal tube placement.
Complete online in region specific *ReadyOp* and upload completed form.
Complete when Ketamine, Etomidate, Succinylcholine and/ or Rocuronium or used to facilitate use of a BIAD and/ or endotracheal intubation. Paramedics/ AEMT should consider using a BIAD if endotracheal intubation is unsuccessful.
- Secure the endotracheal tube well and consider c-collar in pediatric patients (even in absence of trauma) to better maintain ETT placement.
Manual stabilization of endotracheal tube should be used during all patient moves / transfers.
- Airway Cricothyrotomy Percutaneous Needle Procedure:
Indicated as a lifesaving / last resort procedure in pediatric patients < 10 years of age.
Very little evidence to support it's use and safety.
A variety of alternative pediatric airway devices now available make the use of this procedure rare.
Agencies who utilize this procedure must develop a written procedure, establish a training program, maintain equipment and submit procedure and training plan to the State Medical Director/ Regional EMS Office.
 ≥ 10 years: Surgical cricothyrotomy or commercial kits based on agency preference recommended.
- DOPE: Displaced tracheostomy tube/ ETT, Obstructed tracheostomy tube/ ETT, Pneumothorax and Equipment failure.



Pediatric Asthma Respiratory Distress

History


- Time of onset
- Possibility of foreign body
- Past Medical History
- Medications
- Fever / Illness
- Sick Contacts
- History of trauma
- History / possibility of choking
- Ingestion / OD
- Congenital heart disease

Signs and Symptoms

- Wheezing / Stridor / Crackles / Rales
- Nasal Flaring / Retractions / Grunting
- Increased Heart Rate
- AMS
- Anxiety
- Attentiveness / Distractability
- Cyanosis
- Poor feeding
- JVD / Frothy Sputum
- Hypotension

Differential

- Asthma / Reactive Airway Disease
- Aspiration
- Foreign body
- Upper or lower airway infection
- Congenital heart disease
- OD / Toxic ingestion / CHF
- Anaphylaxis
- Trauma





Pediatric Asthma Respiratory Distress

Known or Suspected COVID

Terbutaline can be administered alone without Albuterol / Atrovent if COVID is suspected or known.

This decision is left up to the Provider.

If Albuterol / Atrovent are administered a N95 mask or P100 mask **MUST** be worn by the Provider.

Dosage:

Pediatric: 0.01 mg/kg (up to 0.25mg) SQ. May repeat dosage every 15 minutes as needed.

DO NOT exceed adult dose of 0.25mg.

Pearls

- **Recommended Exam: Mental Status, HEENT, Skin, Neck, Heart, Lungs, Abdomen, Extremities, Neuro**
- **Items in Red Text are key performance measures used to evaluate protocol compliance and care.**
- **This protocol includes all patients with respiratory distress, Asthma, Reactive Airway Disease, croup, or bronchospasm.**
- **Patients may also have wheezing and respiratory distress with viral upper respiratory tract infections and pneumonia.**
- Pulse oximetry should be monitored continuously and consider End-tidal CO₂ monitoring if available.
- **Combination nebulizers containing albuterol and ipratropium (DuoNeb):**
 - Patients may require more than 3 nebulizer treatments, treatments should continue until improvement.
 - Following 3 combination nebulizers (DuoNeb), it is preferable to continue albuterol solely with subsequent treatments as there is no proven benefit to continual use of ipratropium.
- **Epinephrine:**
 - If allergic reaction or anaphylaxis is suspected, give immediately and repeat until improvement.
 - If allergic reaction is not suspected, administer with no improvement and/ or impending respiratory failure.
- Consider Magnesium Sulfate with impending respiratory failure and/ or no improvement.
- Consider IV access when Pulse oximetry remains $\leq 92\%$ after first beta-agonist nebulizer treatment.
- **Do not force a child into a position, allow them to assume position of comfort, typically the tripod position.**
- Bronchiolitis is a viral infection typically affecting infants which results in wheezing which may not respond to beta-agonists. Consider Epinephrine nebulizer if patient < 18 months and not responding to initial beta-agonist treatment.
- Croup typically affects children < 2 years of age. It is viral, possible fever, gradual onset, no drooling is noted.
- Epiglottitis typically affects children > 2 years of age. It is bacterial, with fever, rapid onset, possible stridor, patient wants to sit up to keep airway open, drooling is common. Airway manipulation may worsen the condition.
- In patients using levalbuterol (Xopenex) you may use Albuterol for the first treatment then use the patient's supply for repeat nebulizers or agency's supply.
- A silent chest in respiratory distress is a pre-respiratory arrest sign.
- **EMR/ EMT:**
 - The use of Epinephrine IM is limited to the treatment of anaphylaxis.
 - Administration of diphenhydramine is limited to the oral route only.
- EMT may administer a beta-agonist (Albuterol) to patients currently prescribed the medication and patients who are not currently prescribed the medication.




Post-intubation/ BIAD Management

Capnography Monitoring

- End-tidal (EtCO₂) monitoring is mandatory following placement of an endotracheal tube.
- EtCO₂ monitoring is mandatory following placement of a BIAD once available on scene.

Protocols AR 1, 2, 3, 5, and 6 should be utilized together (even if agency is not using Drug Assisted Airway Protocol) as they contain useful information for airway management.





Post-intubation/ BIAD Management

Lengthy transports may require additional doses of sedatives to maintain patients condition avoiding evidence of Anxiety / Agitation. Changes in vital signs may also determine additional doses.

Epi infusion for Blood Pressure

Mix 1mg of Epi(1:1) into a 500cc bag of NS
This gives you a concentration of 1 mcg per 1/2 cc
Using a 10gtt set: 10gtts = 1cc = 2mcg
Dose range is 1-10mcg, titrate to SBP > 90

Example:

5gtts per min = 1 mcg per min
10 gtts per min = 2 mcg per min
15 gtts per min = 3 mcg per min
20 gtts per min = 4 mcg per min

Epi used as a "push dose" would be 10mcg per min. This can be achieved by drawing up 5cc from infused bag.

Pearls

- **Recommended Exam: Mental Status, HEENT, Heart, Lungs, Neuro**
- **Patients requiring advanced airways and ventilation commonly experience pain and anxiety.**
- **Unrelieved pain can lead to increased catecholamine release, ischemia, immunosuppression, and prolonged hospitalization.**
- **Ventilated patients cannot communicate pain/ anxiety and providers are poor at recognizing pain/ anxiety.**
- **Vital signs such as tachycardia and/ or hypertension can provide clues to inadequate sedation, however they are not always reliable indicators of a patient's lack of adequate sedation.**
- **Sedation strategy:**
 - Pain is the primary reason patients experience agitation and must be addressed first.
 - Opioids and/ or Ketamine are the first line agents, alone or in combination.
 - Benzodiazepines may be utilized if patient is not responding to adequate opioid and/ or Ketamine doses.
 - Paralysis is considered a last resort, only when patients are not responding to opioid, Ketamine, or benzodiazepines.
 - Patients that have received paralytics may be experiencing pain with no obvious signs or symptoms.
 - Consider sedation early after giving paralytics, especially in patients receiving Rocuronium.
- **Ventilation rate:**
 - Guidelines: 30 for Neonates, 25 for Toddlers, 20 for School Age, and for Adolescents the normal Adult rate of 10 – 12 per minute.
 - Maintain EtCO₂ between 35 - 45 and avoid hyperventilation.
- **Ventilator/ Ventilation strategies will need to be tailored to individual patient presentations. Medical director can indicate different strategies above.**
- **Propofol:**
 - Use restricted to agencies approved by the OEMS State Medical Director.
 - Agencies must submit a use policy and education plan to the OEMS.
 - Infusion must be supplied and initiated by a medical facility and may be used only during interfacility transfer.
 - Paramedic may titrate infusion to maintain appropriate sedation but cannot initiate or bolus the medication.
- In general, ventilation with BVM should cause chest rise. With mechanical ventilation a reasonable tidal volume should be about 6 - 8 mL/kg and peak pressures should be < 30 cmH₂O. Plateau Pressures should be < 30 cmH₂O.
- Head of bed should be maintained at least 10 – 20 degrees of elevation when possible, to decrease aspiration risk.
- With abrupt clinical deterioration, if mechanically ventilated, disconnect from ventilator to assess lung compliance.
- **DOPE:** Displaced tracheostomy tube/ ETT, **O**bststructed tracheostomy tube/ ETT, **P**neumothorax and **E**quipment failure.



Ventilator Emergencies

History


- Birth defect (tracheal atresia, tracheomalacia, craniofacial abnormalities)
- Surgical complications (damage to phrenic nerve)
- Trauma (post-traumatic brain or spinal cord injury)
- Medical condition (bronchopulmonary dysplasia, muscular dystrophy)

Signs and Symptoms

- Transport requiring maintenance of a mechanical ventilator
- Power or equipment failure at residence

Differential

- Disruption of oxygen source
- Dislodged or obstructed tracheostomy tube
- Detached or disrupted ventilator circuit
- Cardiac arrest
- Increased oxygen requirement / demand
- Ventilator failure



Airway Respiratory Protocol Section

Pearls

- **Always talk to family/ caregivers as they have specific knowledge and skills.**
- **If using the patient's ventilator bring caregiver knowledgeable in ventilator operation during transport.**
- **Take patient's ventilator to hospital even if not functioning properly.**
- Always use patient's equipment if available and functioning properly.
- Continuous pulse oximetry and End Tidal CO₂ monitoring must be utilized during assessment and transport.
- Unable to correct ventilator problem: Remove patient from ventilator and manually ventilate using BVM.
- Typical alarms:
 - Low Pressure/ Apnea: Loose or disconnected circuit, leak in circuit or around tracheostomy site.
 - Low Power: Internal battery depleted.
 - High Pressure: Plugged/ obstructed airway or circuit.
- **DOPE: Displaced tracheostomy tube/ ETT, Obstructed tracheostomy tube/ ETT, Pneumothorax and Equipment failure.**



Tracheostomy Tube Emergencies

History


- Birth defect (tracheal atresia, tracheomalacia, craniofacial abnormalities)
- Surgical complications (accidental damage to phrenic nerve)
- Trauma (post-traumatic brain or spinal cord injury)
- Medical condition (bronchial or pulmonary dysplasia, muscular dystrophy)

Signs and Symptoms

- Nasal flaring
- Chest wall retractions (with or without abnormal breath sounds)
- Attempts to cough
- Copious secretions noted coming out of the tube
- Faint breath sounds on both sides of chest despite significant respiratory effort
- AMS
- Cyanosis

Differential

- Allergic reaction
- Asthma
- Aspiration
- Septicemia
- Foreign body
- Infection
- Congenital heart disease
- Medication or toxin
- Trauma



Pearls

- Always talk to family/ caregivers as they have specific knowledge and skills.
- Important to ask if patient has undergone laryngectomy. This does not allow mouth/ nasal ventilation by covering stoma.
- Use patients equipment if available and functioning properly.
- Estimate suction catheter size by doubling the inner tracheostomy tube diameter and rounding down.
- Suction depth: Ask family/ caregiver. No more than 3 to 6 cm typically. Instill 2 – 3 mL of NS before suctioning.
- Do not suction more than 10 seconds each attempt and pre-oxygenate before and between attempts.
- DO NOT force suction catheter. If unable to pass, then tracheostomy tube should be changed.
- Always deflate tracheal tube cuff before removal. Continual pulse oximetry and EtCO2 monitoring if available.
- **DOPE:** Displaced tracheostomy tube/ ETT, **O**bststructed tracheostomy tube/ ETT, **P**neumothorax and **E**quipment failure.



Mechanical Ventilation; Adult (Optional)

History

- Multiple etiologies leading to need for advanced airway control
- Requires ventilation support
- Height and underlying lung conditions

Signs and Symptoms

- Loss of consciousness or AMS with inability to protect airway
- Difficult oxygenation and/or ventilation

Differential

- ROSC
- Trauma
- Stroke
- Seizure
- Shock (see Shock Protocol)
- Toxicological

Age Appropriate
 Airway Protocol(s) AR 1, 2, 3, 5, 6
if indicated

Post-intubation/BIAD Management
 Protocol AR 8
if indicated

History of COPD or Asthma?

Alarming Ventilator and unsure how to troubleshoot

- Immediately disconnect patient and use BVM.
- Once oxygenation and ventilation stabilized, restart ventilator set-up procedure.

Home Ventilator Inter-facility Transfer with Ventilator

- Set initial parameters to home or facility settings
- Titrate to oxygenation, work of breathing, SpO₂, and EtCO₂.
- Use home ventilator if functioning properly.

MODE:
Volume – Assist Control

FiO₂: 100%

PEEP: 5 cmH₂O

TIDAL VOLUME (V_t):
8 mL/kg
Follow PBW and V_t on page 3

BPM: RESPIRATORY RATE:
18 BPM

FLOW RATE:
60 mL/min
(preset)

Check Plateau Pressure
Press Manual Breath
P Pressure button
Goal Pressure < 30 cm/H₂O

Decrease Tidal Volume
1 mL/kg increments
Until ≤ 29 cm/H₂O
(DO NOT DECREASE < 4 mL/kg)

MODE:
Volume – Assist Control

FiO₂: 100%

PEEP: 5 cmH₂O

TIDAL VOLUME:
8 mL/kg
Follow PBW and V_t on page 3

BPM: RESPIRATORY RATE:
12 BPM

FLOW RATE:
60 mL/min
(preset)

I:E Ratio
Increase to 1:4 or 1:5

Check Plateau Pressure
Press Manual Breath
P Pressure button
Goal Pressure < 30 cm/H₂O

Decrease Tidal Volume
1 mL/kg increments
Until ≤ 29 cm/H₂O
(DO NOT DECREASE < 4 mL/kg)

After 10 minutes
Decrease FiO₂ down to 50%
Then adjust PEEP and FiO₂
Goal SpO₂ 92 – 98%

Step 1: PEEP =10 FiO₂ =50%

Step 2: PEEP =10 FiO₂ =60%

Step 3: PEEP =10 FiO₂ =70%

Step 4: PEEP =12 FiO₂ =70%

Step 5: PEEP =14 FiO₂ =70%

Check Peak Inspiratory Pressure (PIP)
Goal V_t is 8 mL/kg

ADJUST PIP Alarm Settings

- Up until full exhalation achieved on 8 mL/kg Tidal Volume

Notify Destination or Contact Medical Control

Universal Protocol Section



Mechanical Ventilation; Adult (Optional)

Pearls

- **Recommended Exam: Mental Status, Skin, HEENT, Heart, Lungs, Abdomen, Back, Extremities, Neuro**
- **Mechanical ventilation may be used in any patient ≥ 1 year old.**
- **MODE:**
 - In all adult patients use Volume – Assist Control.
 - This mode requires adequate sedation as it can be uncomfortable in a patient who is awakening.
- **TIDAL VOLUME:**
 - Tidal volume is very important in preventing lung injury and calculated by height and predicted body weight, or ideal body weight, and NOT actual body weight.
 - Follow Tidal Volume by Height Table on page 3.
 - Follow Tidal Volume by Height Table on page 3 when adjusting Peak Inspiratory Pressure alarms to allow full exhalation.
 - High Tidal Volumes are well known to cause alveolar damage and lung injury.
- **FLOW RATE:**
 - A normal breath (non-mechanical ventilation) has highest flow and volume at the beginning and both decrease as inspiration comes to an end.
 - Setting Flow Rate at 60 L/minute allows patient to take full breath without air hunger toward end of inspiration. This is more comfortable for the patient.
 - If patient looks like they are trying to take in more volume initially, the Flow Rate can be increased by increments of 5 as needed to improve patient comfort.
- **FiO₂ and PEEP Adjustments:**
 - Seems intuitive that when SpO₂ is less than desired the FiO₂ should be increased.
 - When FiO₂ is $\geq 50\%$ and SpO₂ remains low, this indicates a shunt, and PEEP must be used in conjunction with FiO₂ to correct the shunt and increase oxygenation.
 - Follow PEEP adjustment recommendations on page 1.
- **EtCO₂:**
 - EtCO₂ and arterial CO₂ do not always correlate well in patients with lung disease or during serious illness or injury.
 - Use caution in adjusting respiratory rate to reach a goal of 35 – 45 mmHg. Most intubated patients do not need tight control in this range.
 - Patients with suspected head injury do need EtCO₂ with a target of 35 – 45 mmHg.
 - Allowing patients with COPD and asthma exacerbations to have higher EtCO₂ outside the 35 – 45 mmHg range is acceptable. Lower ventilation rates allow more time for exhalation and prevents auto-PEEP and/ or air trapping.
- **DOPE:** Displaced tracheostomy tube / ETT, Obstructed tracheostomy tube / ETT, Pneumothorax and Equipment failure.



Mechanical Ventilation; Adult (Optional)

TIDAL VOLUME INITIAL SETTINGS By HEIGHT

FEMALE Height / Predicted body weight / Vt

MALE Height / Predicted body weight / Vt

HEIGHT	PBW	4 ml	5 ml	6 ml	7 ml	8 ml
4' 0" (48)	17.9	72	90	107	125	143
4' 1" (49)	20.2	81	101	121	141	162
4' 2" (50)	22.5	90	113	135	158	180
4' 3" (51)	24.8	99	124	149	174	198
4' 4" (52)	27.1	108	136	163	190	217
4' 5" (53)	29.4	118	147	176	206	235
4' 6" (54)	31.7	127	159	190	222	254
4' 7" (55)	34	136	170	204	238	272
4' 8" (56)	36.3	145	182	218	254	290
4' 9" (57)	38.6	154	193	232	270	309
4' 10" (58)	40.9	164	205	245	286	327
4' 11" (59)	43.2	173	216	259	302	346
5' 0" (60)	45.5	182	228	273	319	364
5' 1" (61)	47.8	191	239	287	335	382
5' 2" (62)	50.1	200	251	301	351	401
5' 3" (63)	52.4	210	262	314	367	419
5' 4" (64)	54.7	219	274	328	383	438
5' 5" (65)	57	228	285	342	399	456
5' 6" (66)	59.3	237	297	356	415	474
5' 7" (67)	61.6	246	308	370	431	493
5' 8" (68)	63.9	256	320	383	447	511
5' 9" (69)	66.2	265	331	397	463	530
5' 10" (70)	68.5	274	343	411	480	548
5' 11" (71)	70.8	283	354	425	496	566
6' 0" (72)	73.1	292	366	439	512	585
6' 1" (73)	75.4	302	377	452	528	603
6' 2" (74)	77.7	311	389	466	544	622
6' 3" (75)	80	320	400	480	560	640
6' 4" (76)	82.3	329	412	494	576	658
6' 5" (77)	84.6	338	423	508	592	677
6' 6" (78)	86.9	348	435	521	608	695
6' 7" (79)	89.2	357	446	535	624	714
6' 8" (80)	91.5	366	458	549	641	732
6' 9" (81)	93.8	375	469	563	657	750
6' 10" (82)	96.1	384	481	577	673	769
6' 11" (83)	98.4	394	492	590	689	787
7' 0" (84)	100.7	403	504	604	705	806

HEIGHT	PBW	4 ml	5 ml	6 ml	7 ml	8 ml
4' 0" (48)	22.4	90	112	134	157	179
4' 1" (49)	24.7	99	124	148	173	198
4' 2" (50)	27	108	135	162	189	216
4' 3" (51)	29.3	117	147	176	205	234
4' 4" (52)	31.6	126	158	190	221	253
4' 5" (53)	33.9	136	170	203	237	271
4' 6" (54)	36.2	145	181	217	253	290
4' 7" (55)	38.5	154	193	231	270	308
4' 8" (56)	40.8	163	204	245	286	326
4' 9" (57)	43.1	172	216	259	302	345
4' 10" (58)	45.4	182	227	272	318	363
4' 11" (59)	47.7	191	239	286	334	382
5' 0" (60)	50	200	250	300	350	400
5' 1" (61)	52.3	209	262	314	366	418
5' 2" (62)	54.6	218	273	328	382	437
5' 3" (63)	56.9	228	285	341	398	455
5' 4" (64)	59.2	237	296	355	414	474
5' 5" (65)	61.5	246	308	369	431	492
5' 6" (66)	63.8	255	319	383	447	510
5' 7" (67)	66.1	264	331	397	463	529
5' 8" (68)	68.4	274	342	410	479	547
5' 9" (69)	70.7	283	354	424	495	566
5' 10" (70)	73	292	365	438	511	584
5' 11" (71)	75.3	301	377	452	527	602
6' 0" (72)	77.6	310	388	466	543	621
6' 1" (73)	79.9	320	400	479	559	639
6' 2" (74)	82.2	329	411	493	575	658
6' 3" (75)	84.5	338	423	507	592	676
6' 4" (76)	86.8	347	434	521	608	694
6' 5" (77)	89.1	356	446	535	624	713
6' 6" (78)	91.4	366	457	548	640	731
6' 7" (79)	93.7	375	469	562	656	750
6' 8" (80)	96	384	480	576	672	768
6' 9" (81)	98.3	393	492	590	688	786
6' 10" (82)	100.6	402	503	604	704	805
6' 11" (83)	102.9	412	515	617	720	823
7' 0" (84)	105.2	421	526	631	736	842

Universal Protocol Section

TROUBLESHOOTING Hypoxia or Deterioration DOPEs		RESPONSE to Hypoxia or Deterioration DOTT	
D	Dislodged ETT or cuff leak	D	Disconnect ventilator, squeeze chest if auto-PEEP, Decompress if pneumothorax
O	Obstruction of ETT or circuit	O	Oxygen 100% FiO2, BVM and check compliance
P	Pneumothorax, Pneumonia, Pulmonary embolism or edema, Plug (mucous)	T	Tube position and function, check EtCO2
E	Equipment problem	T	Tweak ventilator settings or equipment
S	Stacked breaths, air trapping, or auto-PEEP		

Pressure Alarm Troubleshooting			Problem Location	Consider
High PIP	+	High Plateau > 30	Alveoli	Compliance problem: Pneumothorax, Pneumonia Pulmonary Edema or Embolism, CHF
High PIP	+	Normal Plateau < 30	Airway problem	Airway, ventilator, or circuit problem: DOPE, Right Main stem intubation, Air trapping or auto-PEEP, Mucous plug, Patient out of synchrony with ventilator



Adult Asystole / Pulseless Electrical Activity

History

- SAMPLE
- Estimated downtime
- See Reversible Causes below
- DNR, MOST, or Living Will

Signs and Symptoms

- Pulseless
- Apneic
- No electrical activity on ECG
- No heart tones on auscultation

Differential

- See Reversible Causes below



Cardiac Arrest Protocol AC 3

Criteria for Death / No Resuscitation
Review DNR / MOST Form

YES

Decomposition
Rigor mortis
Dependent lividity
Blunt force trauma
Injury incompatible with life
Extended downtime with asystole

Do not begin resuscitation

Follow
Deceased Subjects
Policy

NO

Begin Continuous CPR Compressions
Push Hard (≥ 2 inches)
Push Fast (100 - 120 / min)
Change Compressors every 2 minutes
(sooner if fatigued)
(Limit changes / pulse checks ≤ 10 seconds)

Ventilate 1 breath every 6 seconds
30:2 Compression:Ventilation if no Advanced Airway
Monitor EtCO2
if available

AED Procedure
if available

P

Cardiac Monitor



IV or IO Access Protocol UP 6

A

Epinephrine (1:10,000) 1 mg IV / IO
Repeat every 5 minutes

Normal Saline Bolus 500 mL IV / IO
May repeat as needed
Maximum 2 L

Search for Reversible Causes

Blood Glucose Analysis Procedure
if applicable

P

Available for Agency Medications



On Scene Resuscitation / Termination of Resuscitation
Protocol(s) AC 12
as indicated

Reversible Causes

Hypovolemia
Hypoxia
Hydrogen ion (acidosis)
Hypothermia
Hypo / Hyperkalemia

Tension pneumothorax
Tamponade; cardiac
Toxins
Thrombosis; pulmonary (PE)
Thrombosis; coronary (MI)

Suspected Opioid Overdose

Administer Naloxone per
Overdose / Toxic Ingestion
Protocol TE 7

AT ANY TIME

Return of
Spontaneous
Circulation



Go to
Post Resuscitation
Protocol AC 10



Notify Destination or
Contact Medical Control





Adult Asystole / Pulseless Electrical Activity

Pearls

- **Team Focused Approach / Pit-Crew Approach recommended; assigning responders to predetermined tasks. Refer to optional Team Focused CPR Protocol AC 11 or development of local agency protocol.**
- **Efforts should be directed at high quality and continuous compressions with limited interruptions and early defibrillation when indicated.**
- **DO NOT HYPERVENTILATE: If no advanced airway (BIAD, ETT), compression to ventilation ratio is 30:2. If advanced airway in place, ventilate 10 breaths per minute with continuous, uninterrupted compressions.**
- **Do not interrupt compressions to place endotracheal tube. Consider BIAD first to limit interruptions.**
- **Passive oxygenation optional in agencies practicing Team Focused Approach / Pit-Crew Approach.**
- **Reassess and document BIAD and / or endotracheal tube placement and EtCO2 frequently, after every move, and at transfer of care.**
- **IV / IO access and drug delivery are secondary to high-quality chest compressions and early defibrillation.**
- **IV access is preferred route. Follow IV or IO Access Protocol UP 6.**
- **Defibrillation:** Follow manufacture's recommendations concerning defibrillation / cardioversion energy when specified.
- **End Tidal CO2 (EtCO2)**
 - If EtCO2 is < 10 mmHg, improve chest compressions. Goal is ≥ 20 mmHg.
 - If EtCO2 spikes, typically > 40 mmHg, consider Return of Spontaneous Circulation (ROSC)
- **Special Considerations**
 - **Maternal Arrest** - Treat mother per appropriate protocol with immediate notification to Medical Control and rapid transport preferably to obstetrical center if available and proximate. Place mother supine and perform Manual Left Uterine Displacement moving uterus to the patient's left side. IV/IO access preferably above diaphragm. Defibrillation is safe at all energy levels.
 - **Renal Dialysis / Renal Failure** - Refer to Dialysis / Renal Failure Protocol AM 3 caveats when faced with dialysis / renal failure patient experiencing cardiac arrest.
 - **Opioid Overdose** - If suspected, administer Naloxone per Overdose / Toxic Ingestion Protocol TE 7 while ensuring airway, oxygenation, ventilations, and high-quality chest compressions.
 - **Drowning / Suffocation / Asphyxiation / Hanging / Lightning Strike** – Hypoxic associated cardiac arrest and prompt attention to airway and ventilation is priority followed by high-quality and continuous chest compressions and early defibrillation.
- **Transcutaneous Pacing:**
 - Pacing is NOT effective in cardiac arrest and pacing in cardiac arrest does NOT increase chance of survival
 - Success is based on proper planning and execution. Procedures require space and patient access. Make room to work.
 - Discussion with Medical Control can be a valuable tool in developing a differential diagnosis and identifying possible treatment options.



Bradycardia; Pulse Present

History


- Past medical history
- Medications
 - Beta-Blockers
 - Calcium channel blockers
 - Clonidine
 - Digoxin
- Pacemaker

Signs and Symptoms

- HR < 60/min with hypotension, acute altered mental status, chest pain, acute CHF, seizures, syncope, or shock secondary to bradycardia
- Chest pain
- Respiratory distress
- Hypotension or Shock
- Altered mental status
- Syncope

Differential

- Acute myocardial infarction
- Hypoxia / Hypothermia
- Pacemaker failure
- Sinus bradycardia
- Head injury (elevated ICP) or Stroke
- Spinal cord lesion
- Sick sinus syndrome
- AV blocks (1°, 2°, or 3°)
- Overdose



YES

	Airway Protocol(s) AR 1, 2, 3 <i>if indicated</i>
	Respiratory Distress Protocol AR 4 <i>if indicated</i>
	Chest Pain: Cardiac and STEMI Protocol AC 4 <i>if indicated</i>
B	Search for Reversible Causes
	12 Lead ECG Procedure
	IV / IO Protocol UP 6
P	Cardiac Monitor
A	Normal Saline Fluid Bolus 500 mL – 2 L NS IV / IO (Unless Acute CHF) Maximum 2 L
	Atropine 1 mg IV / IO May repeat every 3 – 5 minutes Maximum 3 mg
P	Epinephrine 1 - 10 mcg/min IV / IO Titrate to SBP ≥ 90 mmHg
	OR Norepinephrine 0.01-1.0 mcg/kg/min IV/IO via pump Titrate to SBP ≥ 90mmHg
	If No Improvement Transcutaneous Pacing Procedure (<i>Consider earlier in 2nd or 3rd AVB</i>)
	Notify Destination or Contact Medical Control

Suspected Beta-Blocker or Calcium Channel Blocker

Follow Overdose/ Toxic Ingestion Protocol TE 7

Reversible Causes

Hypovolemia
Hypoxia
Hydrogen ion (acidosis)
Hypothermia
Hypo / Hyperkalemia

Tension pneumothorax
Tamponade; cardiac
Toxins
Thrombosis; pulmonary (PE)
Thrombosis; coronary (MI)

Consider Sedation

Midazolam 2 – 2.5 mg
IV / IO / IM / IN

Maximum 10 mg



Bradycardia; Pulse Present

Epi. Infusion

1mg (1:1) of Epi in 500cc bag of NS Concentration = 1mcg per 1/2 cc Use 10gtt set: 10gtts = 1cc = 2mcg Dose = 1-10mcg, titrate to SBP>90

5 gtts per min = 1 mcg per min
10 gtts per min = 2 mcg per min
15 gtts per min = 3 mcg per min
20 gtts per min = 4 mcg per min
25 gtts per min = 5 mcg per min
etc.

Norepinephrine Infusion

*** IV Pump is REQUIRED ***

Mix 4mg in 250cc bag of D5W

This gives you a concentration of 16 mcg per cc.
begin at 0.01 mcg/kg/min and titrate SBP

*** IV Pump is REQUIRED ***

Pearls

- **Recommended Exam: Mental Status, HEENT, Skin, Heart, Lungs, Abdomen, Back, Extremities, Neuro**
- **Identifying signs and symptoms of poor perfusion caused by bradycardia are paramount.**
- **Rhythm should be interpreted in the context of symptoms and pharmacological treatment given only when symptomatic, otherwise monitor and reassess.**
- **Consider hyperkalemia with wide complex, bizarre appearance of QRS complex, and bradycardia. Give Calcium Chloride or Gluconate in addition to Sodium Bicarbonate if hyperkalemia suspected.**
- **12-Lead ECG:**
12 Lead ECG not necessary to diagnose and treat
Obtain when patient is stable and/or following rhythm conversion.
- **Unstable condition**
Condition which acutely impairs vital organ function and cardiac arrest may be imminent.
If at any point patient becomes unstable move to unstable arm in algorithm.
- Hypoxemia is a common cause of bradycardia. Ensure oxygenation and support respiratory effort.
- **Atropine:**
Atropine is considered a first line agent in symptomatic bradycardia.
Ineffective and potentially harmful in cardiac transplantation. May cause paradoxical bradycardia.
- **Symptomatic bradycardia causing shock or peri-arrest condition:**
If no IV or IO access immediately available start Transcutaneous Pacing, establish IV / IO access, and then administer atropine and/or epinephrine.
Epinephrine or Dopamine may be considered if no response to Atropine.
- **Symptomatic condition**
Arrhythmia is causing symptoms such as palpitations, lightheadedness, or dyspnea, but cardiac arrest is not imminent.
Symptomatic bradycardia usually occurs at rates < 50 beats per minute.
Search for underlying causes such as hypoxia or impending respiratory failure.
- **Serious Signs / Symptoms:**
Hypotension. Acutely altered mental status. Signs of shock / poor perfusion. Chest pain with evidence of ischemia (STEMI, T wave inversions or depressions.) Acute CHF.
- **Transcutaneous Pacing Procedure (TCP)**
Indicated with unstable bradycardia unresponsive to medical therapy.
If time allows transport to specialty center because transcutaneous pacing is a temporizing measure.
Transvenous / permanent pacemaker will probably be needed.
Immediate TCP with high-degree AV block (2d or 3d degree) with no IV / IO access.
- Consider treatable causes for bradycardia (Beta Blocker OD, Calcium Channel Blocker OD, etc.)



Cardiac Arrest; Adult

AT ANY TIME

Return of Spontaneous Circulation



Go to Post Resuscitation Protocol AC 10

Criteria for Death / No Resuscitation
Review DNR / MOST Form

YES

Decomposition
Rigor mortis
Dependent lividity
Blunt force trauma
Injury incompatible with life
Extended downtime with asystole

Do not begin resuscitation

Follow Deceased Subjects Policy

NO

Begin Continuous CPR Compressions
Push Hard (≥ 2 inches)
Push Fast (100 - 120 / min)
Change Compressors every 2 minutes (sooner if fatigued)
(Limit changes / pulse checks ≤ 10 seconds)
Change to LUCAS if available

Ventilate 1 breath every 6 seconds
30:2 Compression:Ventilation if no Advanced Airway
Monitor EtCO2 if available

AED Procedure *if available*

ALS Available

P Cardiac Monitor

Shockable Rhythm

NO

YES

AED Procedure

Shockable Rhythm

NO

YES

Asystole / PEA
Protocol AC 1
as indicated

Airway
Protocol(s) AR 1, 2, 3

VF / VT
Protocol AC 9
Tachycardia
Protocol(s) AC 6, 7
as indicated

Airway
Protocol(s) AR 1, 2, 3

Continue CPR
2 Minutes

Repeat and reassess

Airway
Protocol(s) AR 1, 2, 3

Arrest secondary to Opioid OD?

NO

YES

A

Naloxone 0.4 – 2 mg IN / IM
Peds: 0.1 mg/kg IN

Maximum 4 mg

Naloxone 0.4 – 2 mg
Peds: 0.1 mg/kg
IV / IO / IM / IN / ETT

Maximum 4 mg

Termination on Scene
Protocol AC 12
as indicated

 **Notify Destination or Contact Medical Control** 



Cardiac Arrest; Adult

LUCAS

1. Power on the LUCAS while rescuer 2 provides manual CPR.
2. Place the LUCAS back plate with assistance from rescuer 2, then rescuer 2 resumes CPR.
3. Attach the upper part, listen for "click".
4. Push down suction cup, lower edge over end of sternum. Adjust if needed.
5. Push PAUSE to lock START POSITION.
6. Push ACTIVE to start compressions.
7. Attach stabilization strap.
8. Place patients arms in Writ Straps provided at top of device.
9. Follow protocol.

Pearls

- **Team Focused Approach / Pit-Crew Approach recommended; assign responders to predetermined tasks. Refer to optional protocol or development of local agency protocol.**
- **Efforts should be directed at high quality and continuous compressions with limited interruptions and early defibrillation when indicated.**
- **DO NOT HYPERVENTILATE: If no advanced airway (BIAD, ETT) compression to ventilation ratio is 30:2. If advanced airway in place, ventilate 10 breaths per minute with continuous, uninterrupted compressions.**
- **Do not interrupt compressions to place endotracheal tube. Consider BIAD first to limit interruptions.**
- **Passive oxygenation optional in agencies practicing Team Focused Approach / Pit-Crew Approach.**
- **Reassess and document BIAD and / or endotracheal tube placement and EtCO2 frequently, after every move, and at transfer of care.**
- **IV / IO access and drug delivery is secondary to high-quality chest compressions and early defibrillation.**
- **IV access is preferred route. Follow IV or IO Access Protocol UP 6.**
- **Defibrillation:**
 - Follow manufacture's recommendations concerning defibrillation / cardioversion energy when specified.
 - Charge defibrillator during chest compressions, near the end of 2-minute cycle, to decrease peri-shock pause.
 - Following defibrillation, provider should immediately restart chest compressions with no pulse check until end of next cycle.
- **End Tidal CO2 (EtCO2)**
 - If EtCO2 is < 10 mmHg, improve chest compressions. Goal is ≥ 20 mmHg.
 - If EtCO2 spikes, typically > 40 mmHg, consider Return of Spontaneous Circulation (ROSC)
- **Special Considerations**
 - Maternal Arrest** - Treat mother per appropriate protocol with immediate notification to Medical Control and rapid transport preferably to obstetrical center if available and proximate. Place mother supine and perform Manual Left Uterine Displacement moving uterus to the patient's left side. IV/IO access preferably above diaphragm. Defibrillation is safe at all energy levels.
 - Renal Dialysis / Renal Failure** - Refer to Dialysis / Renal Failure Protocol AM 3 caveats when faced with dialysis / renal failure patient experiencing cardiac arrest.
 - Opioid Overdose** - If suspected, administer Naloxone per Overdose / Toxic Ingestion Protocol TE 7 while ensuring airway, oxygenation, ventilations, and high-quality chest compressions.
 - Drowning / Suffocation / Asphyxiation / Hanging / Lightning Strike** – Hypoxic associated cardiac arrest and prompt attention to airway and ventilation is priority followed by high-quality and continuous chest compressions and early defibrillation.
- **Transcutaneous Pacing:**
 - Pacing is NOT effective in cardiac arrest and pacing in cardiac arrest does NOT increase chance of survival
- Success is based on proper planning and execution. Procedures require space and patient access. Make room to work.
- Discussion with Medical Control can be a valuable tool in developing a differential diagnosis and identifying possible treatment options.



Chest Pain: Cardiac and STEMI

History

- Age
- Medications (Viagra / sildenafil, Levitra / vardenafil, Cialis / tadalafil)
- Past medical history (MI, Angina, Diabetes, post menopausal)
- Allergies
- Recent physical exertion
- Onset / Palliation / Provocation
- Quality (crampy, constant, sharp, dull, etc.)
- Region / Radiation / Referred
- Severity (1-10)
- Time (onset/duration / repetition)

Signs and Symptoms

- CP (pain, pressure, aching, vice-like tightness)
- Location (substernal, epigastric, arm, jaw, neck, shoulder)
- Radiation of pain
- Pale, diaphoresis
- Shortness of breath
- Nausea, vomiting, dizziness
- **Time of Onset**
- Women:
 - More likely to have dyspnea, N/V, weakness, back or jaw pain

Differential

- Trauma vs. Medical
- Angina vs. Myocardial infarction
- Pericarditis
- Pulmonary embolism
- Asthma / COPD
- Pneumothorax
- Aortic dissection or aneurysm
- GE reflux or Hiatal hernia
- Esophageal spasm
- Chest wall injury or pain
- Pleural pain
- Overdose: Cocaine or Methamphetamine

	12 Lead ECG Procedure
B	Aspirin 81 mg x 4 PO (chewed) Or 325 mg PO
	Nitroglycerin 0.3 / 0.4 mg Sublingual Repeat every 5 minutes x 3 <i>if prescribed to patient and (BP ≥ 100)</i>
P	Cardiac Monitor

Acute MI / STEMI
See box to right

Transport based on:
STEMI Triage and Destination Plan

Notification of Facility Immediate (≤ 5 minutes)
NHRMC- Activate via Pulsara App
GSRMC / Seacoast- Immediate Transmission of ECG
Keep Scene Time to ≤ 10 Minutes

Acute MI / STEMI

STEMI Definition:

- ≥ 1 mm ST Segment elevation in ≥ 2 contiguous leads
- ≥ 2 mm ST/J point elevation in V2-V3 for men
- ≥ 1.5 mm ST/J point elevation in V2-V3 for women
- ECG software diagnoses Acute MI (symptomatic)

	IV / IO Protocol UP 6
A	Nitroglycerin 0.3 / 0.4 mg SL Repeat every 5 minutes as needed (until pain free)
	Nitroglycerin Paste (pain free) SBP > 100 1inch SBP > 150 1.5 inch SBP > 200 2 inch
P	Morphine 2 – 4 mg IV / IO Repeat every 5 minutes as needed Maximum 10 mg Or Fentanyl 50 – 75 mcg IV / IO Repeat 25 mcg every 20 minutes as needed Maximum 200 mcg
	Hypotension / Shock Protocol AM 5 <i>if indicated</i>
	CHF / Pulmonary Edema Protocol AC 5 <i>if indicated</i>

B	<i>If transporting to Non PCI Center</i> Reperfusion Checklist
A	Nitroglycerin 0.3 / 0.4 mg SL Repeat every 5 minutes as needed (until pain free)
	Nitroglycerin Paste (pain free)
P	Morphine 2 – 4 mg IV / IO Repeat every 5 minutes as needed Maximum 10 mg Or Fentanyl 50 – 75 mcg IV / IO Repeat 25 mcg every 20 minutes as needed Maximum 200 mcg
	Heparin 60 U/kg IV/IO Maximum 10,000 U

Notify Destination or Contact Medical Control



Chest Pain: Cardiac and STEMI

Target oxygen saturation is 94 - 98%

A 12 Lead should be obtained within 5 minutes of arriving on scene.

Code STEMI should be activated within 5 minutes of interpretation of the 12 Lead as a STEMI.

A 15 Lead ECG should be obtained on all patients with an inferior wall MI, normal 12 Lead ECG, and those with a strong suspicion of having a posterior (V8, V9) or right ventricular involvement (V4R).

NTG Spray should be administered every 5 minutes as needed until pain free (monitor for hypotension).
NTG Paste should be placed when patient is pain free (monitor for hypotension).

Document OPQRST:

- **Onset**- What was the patient doing when it started? (active, inactive, stressed, etc.)
- **Provocation / Palliation** - Does the patient believe that activity prompted the pain, and whether the onset was sudden, gradual or part of an ongoing chronic problem.
- **Quality** - Have the patient describe the pain. Is it dull, throbbing, aching, burning, sharp, crushing, shooting, tearing, constant, intermittent, etc.?
- **Region and Radiation** - Where is the pain and does it move?
- **Severity** - Use the 0-10 pain scale. This can be comparative ("...compared to the worst pain you have ever experienced") or imaginative ("...compared to having your leg broken").
- **Timing** - Identify when the pain started. How long has the condition been going on?

Pearls

- **Recommended Exam: Mental Status, Skin, Neck, Lung, Heart, Abdomen, Back, Extremities, Neuro**
- **Items in Red Text are the key performance indicators for the EMS Acute Cardiac (STEMI) Care Toolkit**
- **Nitroglycerin:**
 - Avoid Nitroglycerin in any patient who has used Viagra (sildenafil) or Levitra (vardenafil) in the past 24 hours or Cialis (tadalafil) in the past 36 hours due to potential severe hypotension.
 - Nitroglycerin may cause hypotension during any type myocardial infarction. It is NOT more likely to cause hypotension in an inferior MI and should NOT be avoided unless already hypotensive.
- **STEMI (ST-Elevation Myocardial Infarction)**
 - Positive Reperfusion Checklist should be transported to the appropriate facility based on STEMI EMS Triage and Destination Plan.
 - Consider placing 2 IV sites in the left arm: Many PCI centers use the right radial artery for intervention.
 - Consider placing defibrillator pads on patient as a precaution.
 - Consider Normal Saline or Lactated Ringers bolus of 250 – 500 mL as pre-cath hydration.
 - Scene time goal is < 10 minutes.
 - Document and time-stamp facility STEMI notification and make notification as soon as possible.
 - Document the time of the 12-Lead ECG in the PCR as a Procedure along with the interpretation (Paramedic).
- **Cardiac related symptoms in men and women:**
 - Pressure, squeezing, fullness, or pain in the chest.
 - Pain or discomfort in one or both arms, the back, neck, jaw, or stomach.
 - Shortness of breath with or without chest pain.
 - Sweating, nausea, weakness, and/or lightheadedness.
 - **Women, diabetic patients, and the elderly often experience only weakness, shortness of breath, nausea/vomiting, and back or jaw pain.**
- If patient has taken nitroglycerin without relief, consider potency of the medication.
- Monitor for hypotension after administration of nitroglycerin and opioids.
- **EMT may administer Nitroglycerin to patients already prescribed medication. May give from EMS supply.**



CHF / Pulmonary Edema

History

- Congestive heart failure
- Past medical history
- Medications (digoxin, Lasix, Viagra / sildenafil, Levitra / vardenafil, Cialis / tadalafil)
- Cardiac history --past myocardial infarction

Signs and Symptoms

- Respiratory distress, bilateral rales
- Apprehension, orthopnea
- Jugular vein distention
- Pink, frothy sputum
- Peripheral edema, diaphoresis
- Hypotension, shock
- Chest pain

Differential

- Myocardial infarction
- Congestive heart failure
- Asthma
- Anaphylaxis
- Aspiration
- COPD
- Pleural effusion
- Pneumonia
- Pulmonary embolus
- Pericardial tamponade
- Toxic Exposure

	Airway Protocol(s) AR 1, 2, 3 as indicated
	Chest Pain and STEMI Protocol AC 4 if indicated
B	12 Lead ECG Procedure
	Nitroglycerin 0.3 / 0.4 mg Sublingual Repeat every 5 minutes x 3 if prescribed to patient and (BP >100)
P	Cardiac Monitor
	IV / IO Procedure

Assess Symptom Severity

MILD
Normal Heart Rate
Elevated or Normal BP

MODERATE / SEVERE
Elevated Heart Rate
Elevated BP

CARDIOGENIC SHOCK
Tachycardia followed by bradycardia
Hypertension followed by hypotension

A	Nitroglycerin 0.3 / 0.4 mg SL Repeat every 5 minutes
	Nitroglycerin Paste SBP > 100 1 inch SBP > 150 1.5 inch SBP > 200 2 inch

Improving

YES

NO

B	Airway NIPPV Procedure
A	Nitroglycerin 0.3 / 0.4 mg SL Repeat every 5 minutes
	Nitroglycerin Paste SBP > 100 1 inch SBP > 150 1.5 inch SBP > 200 2 inch
P	ACE-Inhibitor
	Consider Furosemide 40 mg IV ONLY IF Transport time > 30 minutes Known CHF / Daily Lasix Afebrile

B	Remove NIPPV if in place
	Adult Hypotension / Shock Protocol AM 5 if indicated

Notify Destination or Contact Medical Control

Adult Cardiac Protocol Section



CHF / Pulmonary Edema

Pearls

- **Recommended Exam: Mental Status, Skin, Neck, Lung, Heart, Abdomen, Back, Extremities, Neuro**
- **Items in Red Text are key performance measures used to evaluate protocol compliance and care**
- **Diuretics (furosemide) and opioids have NOT been shown to improve the outcomes of EMS patients with pulmonary edema. Even though this historically has been a mainstay of EMS treatment, it is no longer routinely recommended.**
- **Nitroglycerin:**
 - **Avoid Nitroglycerin in any patient who has used Viagra (sildenafil) or Levitra (vardenafil) in the past 24 hours or Cialis (tadalafil) in the past 36 hours due to potential severe hypotension.**
 - **Nitroglycerin may cause hypotension during any type myocardial infarction. It is NOT more likely to cause hypotension in an inferior MI and should NOT be avoided unless already hypotensive.**
- **Document the time of the 12-Lead ECG in the PCR as a Procedure along with the interpretation (Paramedic).**
- **Consider myocardial infarction in all these patients. Diabetics, geriatric and female patients often have atypical pain, or only generalized complaints.**
- **Cardiac related symptoms in men and women:**
 - **Pressure, squeezing, fullness, or pain in the chest.**
 - **Pain or discomfort in one or both arms, the back, neck, jaw, or stomach.**
 - **Shortness of breath with or without chest pain.**
 - **Sweating, nausea, weakness, and/or lightheadedness.**
 - **Women, diabetic patients, and the elderly often experience only weakness, shortness of breath, nausea/ vomiting, and back or jaw pain.**
- **If patient has taken nitroglycerin without relief, consider potency of the medication.**
- **Contraindications to opioids include severe COPD and respiratory distress. Monitor the patient closely.**
- **Monitor for hypotension after administration of nitroglycerin and opioids.**
- **Allow the patient to be in their position of comfort to maximize their breathing effort.**
- **EMT may administer Nitroglycerin to patients already prescribed medication. May give from EMS supply.**
- **Agency medical director may require Contact of Medical Control.**



Adult Tachycardia

NARROW (≤ 0.11 sec)

History

- Age
- Past medical history (MI, Angina, Diabetes, post menopausal)
- Recent physical exertion
- Palpitations, irregular heart beat
- Time (onset/duration / repetition)

Signs and Symptoms

- Chest pain, heart failure, dyspnea
- AMS
- Shock, poor perfusion, hypotension
- Pale, diaphoresis
- Shortness of breath
- Nausea, vomiting, dizziness

Differential

- Trauma vs. Medical
- Sinus Tachycardia vs. dysrhythmia
- Fever, sepsis, infection
- Pericarditis, pulmonary embolism
- Aortic dissection or aneurysm
- Overdose: Stimulants

Assess tachycardia in context of clinical condition
Identify and treat underlying cause of tachycardia

Unstable / Serious Signs and Symptoms
HR Typically ≥ 150
Hypotension, Acute AMS, Ischemic Chest Pain,
Acute CHF, Seizures, Syncope, or Shock
secondary to tachycardia

P	<p>Consider <u>Only if regular/narrow complex</u> Adenosine 6 mg IV / IO Rapid push with flush May repeat 12 mg IV / IO</p>
	<p>Cardioversion Procedure</p>
	<p>Consider Sedation Prior to Cardioversion Midazolam 2 – 2.5 mg IV / IO May repeat as needed Maximum 10 mg OR Lorazepam 1 mg IV/IO, 2 mg IM OR Diazepam 5 mg IV/IO</p>
	<p>Synchronized Narrow and Regular: 50 – 100J Narrow and Irregular: 120 – 200J <i>May repeat and increase dose with subsequent cardioversion attempts</i></p>

NO

B	12 Lead ECG Procedure
P	Cardiac Monitor
	IV or IO Access Protocol UP 6

Regular Rhythm?

YES

P	<p>Attempt Vagal Maneuvers Procedure</p> <p>Adenosine 6 mg IV / IO Rapid push with flush</p> <p>May repeat 12 mg IV / IO May repeat 12 mg IV / IO</p>
----------	---

NO

P	<p>Diltiazem 0.25 mg/kg IV / IO Over 2 – 3 minutes Maximum 25 mg</p> <p><u>If No Improvement in 20 minutes</u></p> <p>Diltiazem 0.35 mg/kg IV / IO Over 2-3 minutes Maximum 25 mg</p> <p><u>If No Improvement in 20 minutes</u> Diltiazem 5 – 15 mg/hr IV / IO</p> <p>Or</p> <p>Amiodarone 150 mg in 100 mL of D5W IV / IO Over 10 minutes</p> <p>Or</p> <p>Metoprolol 5 mg IV / IO Over 5 minutes</p>
	<p>Monitor and Reassess</p>
	<p> Notify Destination or Contact Medical Control </p>



Adult Tachycardia

NARROW (≤ 0.11 sec)


Amiodarone Drip: Preferred

Mix Amiodarone 150 mg in 100mL of D5W IV / IO Infuse over 10 minutes

OR

Amiodarone 150 mg **SLOW IV PUSH** diluted in 10cc NS flush
MUST be given over 10 minutes

WPW- short PR interval(<120 ms), wide QRS complex longer than 120ms with a slurred onset of the QRS waveform, termed a delta wave, in the early part of the QRS.



Pearls

- **Recommended Exam: Mental Status, Skin, Neck, Lung, Heart, Abdomen, Back, Extremities, Neuro**
- **Most important goal is to differentiate the type of tachycardia and if STABLE or UNSTABLE and SYMPTOMATIC.**
- **12-Lead ECG:**
12 Lead ECG not necessary to diagnose and treat
Obtain when patient is stable and/or following rhythm conversion.
- **Unstable condition**
Condition which acutely impairs vital organ function and cardiac arrest may be imminent.
If at any point patient becomes unstable move to unstable arm in algorithm.
- Search for underlying cause of tachycardia such as fever, sepsis, dyspnea, etc.
- Typical sinus tachycardia is in the range of 100 to (200 - patient's age) beats per minute.
- **Symptomatic condition**
Arrhythmia is causing symptoms such as palpitations, lightheadedness, or dyspnea, but cardiac arrest is not imminent.
Symptomatic tachycardia usually occurs at rates ≥ 150 beats per minute.
Patients symptomatic with heart rates < 150 likely have impaired cardiac function such as CHF.
- **Serious Signs / Symptoms:**
Hypotension. Acutely altered mental status. Signs of shock / poor perfusion. Chest pain with evidence of ischemia (STEMI, T wave inversions or depressions.) Acute CHF.
- **If patient has history or 12 Lead ECG reveals Wolfe Parkinson White (WPW):**
DO NOT administer a Calcium Channel Blocker (e.g. Diltiazem) or Beta Blockers.
Use caution with Adenosine and give only with defibrillator available.
- **Regular Narrow-Complex Tachycardia:**
Vagal maneuvers and adenosine are preferred. Vagal maneuvers may convert 19% to 54 % of SVT.
Using passive leg raise with Valsalva is more effective.
Adenosine should be pushed rapidly via proximal IV site followed by 20 mL Normal Saline rapid flush.
Adenosine should not be used in the post-cardiac transplant patient without **Contact of Medical Control**.
Agencies using both calcium channel blockers and beta blockers should choose one primarily. Giving the agents sequentially requires **Contact of Medical Control**. This may lead to profound bradycardia / hypotension.
- **Irregular Narrow-Complex Tachycardia:**
Rate control is more important in pre-hospital setting rather than focus on rhythm conversion.
- **Synchronized Cardioversion:**
Recommended to treat UNSTABLE Atrial Fibrillation, Atrial Flutter and SVT.
- Monitor for hypotension after administration of Calcium Channel Blockers or Beta Blockers.
- Document all rhythm changes with monitor strips and obtain monitor strips with each therapeutic intervention.



Adult Monomorphic Tachycardia

Wide Complex (≥ 0.12 sec)

History


- Age
- Past medical history (MI, Angina, Diabetes, post menopausal)
- Recent physical exertion
- Palpitations, irregular heart beat
- Time (onset /duration / repetition)

Signs and Symptoms

- Chest pain, heart failure, dyspnea
- AMS
- Shock, poor perfusion, hypotension
- Pale, diaphoresis
- Shortness of breath
- Nausea, vomiting, dizziness


Differential

- Trauma vs. Medical
- Sinus Tachycardia vs. dysrhythmia
- Fever, sepsis, infection
- Pericarditis, pulmonary embolism
- Aortic dissection or aneurysm
- Overdose: Stimulants



Cardiac Monitor
Cardioversion Procedure
Consider Sedation Prior to Cardioversion Midazolam 2 – 2.5 mg IV / IO OR Lorazepam 1mg IV/IO, 2 mg IM OR Diazepam 5mg IV/IO
Wide: Regular and Irregular: 200 – 360J • Monomorphic QRS (Synchronized) • Polymorphic QRS (Not-Synchronized)
<i>May repeat and increase dose with subsequent cardioversion attempts</i>

B	12 Lead ECG Procedure
P	Cardiac Monitor
	IV or IO Access Protocol UP 6
P	Consider consultation with medical control



P	Attempt Vagal Maneuvers Procedure <i>Only if regular monomorphic complex</i>
P	Consider Only if regular monomorphic complex Adenosine 6 mg IV / IO Rapid push with flush May repeat 12 mg IV / IO

P	Amiodarone 150 mg in 100 mL of D5W IV / IO Infuse over 10 minutes	
	May repeat if wide complex tachycardia recurs	
	Amiodarone 450 mg in 250 mL of D5W 1 mg/min (33 mL/hr) IV / IO	
	Or Procainamide 20 – 50 mg / min IV / IO	
	Procainamide 1 – 4 mg / min	
	Maximum 17 mg / kg	
	Monitor and Reassess	
	Notify Destination or Contact Medical Control	

Monomorphic QRS:

- All QRS complexes in a single lead are similar in shape.



Adult Monomorphic Tachycardia

Wide Complex (≥ 0.12 sec)

Amiodarone Drip (Preferred):

Mix Amiodarone 150mg in 100mL of D5W IV / IO and infuse over 10 minutes

OR

Amiodarone 150mg **SLOW IV PUSH** diluted in 10cc NS flush
MUST be given over 10 minutes

Pearls

- **Recommended Exam: Mental Status, Skin, Neck, Lung, Heart, Abdomen, Extremities, Neuro**
- **Most important goal is to differentiate the type of tachycardia and if STABLE or UNSTABLE and if SYMPTOMATIC.**
- **12-Lead ECG:**
12-Lead ECG is not necessary to diagnose and treat arrhythmia. A single lead ECG is often all that is needed. Obtain 12-Lead when patient is stable and/ or following a rhythm conversion.
- **Monomorphic QRS:**
All QRS complexes in a single lead are similar in shape.
- **Polymorphic QRS:**
QRS complexes in a single lead will change shape from complex to complex.
- **Rhythm should be interpreted in the context of symptoms and pharmacological or electrical treatment given only when symptomatic, otherwise monitor and reassess.**
- **Unstable condition**
Condition which acutely impairs vital organ function and cardiac arrest may be impending.
If at any point patient becomes unstable move to unstable arm in algorithm.
- **Symptomatic condition**
Arrhythmia is causing symptoms such as palpitations, lightheadedness, or dyspnea but cardiac arrest is not impending.
Symptomatic tachycardia usually occurs at rates ≥ 150 beats per minute. Patients symptomatic with heart rates < 150 likely have impaired cardiac function such as CHF.
- **Serious Signs/ Symptoms:**
Hypotension. Acutely altered mental status. Signs of shock/ poor perfusion. Chest pain with evidence of ischemia (STEMI, T wave inversions or depressions.) Acute congestive heart failure.
- Search for underlying cause of tachycardia such as fever, sepsis, dyspnea, etc.
- Typical sinus tachycardia is in the range of 100 to (220 – patients age) beats per minute.
- If patient has history or 12-Lead ECG reveals Wolfe Parkinson White (WPW), DO NOT administer a Calcium Channel Blocker (e.g., Diltiazem) or Beta Blockers. Use caution with Adenosine and give only with defibrillator available.
- **Regular Wide-Complex Tachycardia:**
Unstable condition:
Immediate defibrillation if pulseless and begin CPR.
Stable condition:
Typically VT or SVT with aberrancy. Adenosine may be given if regular and monomorphic and if defibrillator available.
Verapamil contraindicated in wide-complex tachycardias.
Agencies using Amiodarone, Procainamide, and Lidocaine need to choose one agent primarily. Giving multiple anti-arrhythmics requires contact of Medical Control.
Atrial arrhythmias with WPW should be treated with Amiodarone or Procainamide
- **Irregular Tachycardia:**
Wide-complex, irregular tachycardia: Do not administer calcium channel, beta blockers, or adenosine as this may cause paradoxical increase in ventricular rate. This will usually require cardioversion. Contact Medical Control.
- Document all rhythm changes with monitor strips and obtain monitor strips with each therapeutic intervention.



Adult Polymorphic Tachycardia WIDE (≥ 0.12 sec) Torsades de pointes

History

- Age
- Past medical history (MI, Angina, Diabetes, post menopausal)
- Recent physical exertion
- Palpitations, irregular heart beat
- Time (onset/duration / repetition)

Signs and Symptoms

- Chest pain, heart failure, dyspnea
- AMS
- Shock, poor perfusion, hypotension
- Pale, diaphoresis
- Shortness of breath
- Nausea, vomiting, dizziness

Differential

- Cardiac arrest
- Sinus Tachycardia vs. dysrhythmia
- Fever, sepsis, infection
- Pericarditis, pulmonary embolism
- Aortic dissection or aneurysm
- Overdose

**Assess tachycardia in context of clinical condition
Identify and treat underlying cause of tachycardia**

Unstable / Serious Signs and Symptoms
HR Typically ≥ 150
Hypotension, Acute AMS, Ischemic Chest Pain,
Acute CHF, Seizures, Syncope, or Shock
secondary to tachycardia

P	Defibrillation Procedure
	Consider Sedation Prior to Defibrillation Midazolam 2 – 2.5 mg IV / IO May repeat as needed Maximum 10 mg OR Lorazepam 1 mg IV/IO, 2 mg IM OR Diazepam 5 mg IV/IO
	<u>Wide and Irregular: 200 – 360J</u> Polymorphic QRS (Not-Synchronized) <i>May repeat and increase dose with subsequent cardioversion attempts</i>

B	12 Lead ECG Procedure
P	Cardiac Monitor
	IV or IO Access Protocol UP 6

Pulse Present?

Consider consultation with medical control

NO
Exit to
Cardiac Arrest
Protocol AC 3

QT Interval < 500 msec

QT Interval > 500 msec

P	Amiodarone 150 mg in 100 mL of D5W IV / IO Infuse over 10 minutes May repeat if tachycardia recurs or persists
	Amiodarone 450 mg in 250 mL of D5W 1 mg/min (33 mL/hr) Or Lidocaine 1 – 1.5 mg/kg IV / IO May repeat if refractory Lidocaine 0.75 mg/kg IV / IO Maximum 3 mg/kg
	Monitor and Reassess

P	<i>Consider</i> Magnesium 2 g IV / IO May repeat Maximum 4 g
	Monitor and Reassess

Polymorphic QRS:
• QRS complexes in a single lead will change shape from complex to complex.

Notify Destination or Contact Medical Control



Adult Polymorphic Tachycardia WIDE (≥ 0.12 sec) Torsades de pointes

Amiodarone Drip: Preferred

Mix Amiodarone 150 mg in 100mL of D5W IV / IO Infuse over 10 minutes

OR

Amiodarone 150 mg **SLOW IV PUSH** diluted in 10cc NS flush
MUST be given over 10 minutes

Pearls

- **Recommended Exam: Mental Status, Skin, Neck, Lung, Heart, Abdomen, Back, Extremities, Neuro**
 - **Most important goal is to differentiate the type of tachycardia and if STABLE or UNSTABLE and SYMPTOMATIC.**
 - **12-Lead ECG:**
 - 12 Lead ECG not necessary to diagnose and treat
 - Obtain when patient is stable and/or following rhythm conversion.
 - **Monomorphic QRS:**
 - All QRS complexes in a single lead are similar in shape.
 - **Polymorphic QRS:**
 - QRS complexes in a single lead will change shape from complex to complex.
 - **Rhythm should be interpreted in the context of symptoms and pharmacological or electrical treatment given only when symptomatic, otherwise monitor and reassess.**
 - **Unstable condition**
 - Condition which acutely impairs vital organ function and cardiac arrest may be imminent.
 - If at any point patient becomes unstable move to unstable arm in algorithm.
 - **Symptomatic condition**
 - Arrhythmia is causing symptoms such as palpitations, lightheadedness, or dyspnea, but cardiac arrest is not imminent.
 - Symptomatic tachycardia usually occurs at rates ≥ 150 beats per minute. Patients symptomatic with heart rates < 150 likely have impaired cardiac function such as CHF.
 - **Serious Signs / Symptoms:**
 - Hypotension. Acutely altered mental status. Signs of shock / poor perfusion. Chest pain with evidence of ischemia (STEMI, T wave inversions or depressions.) Acute congestive heart failure.
 - Search for underlying cause of tachycardia such as fever, sepsis, dyspnea, etc.
 - Typical sinus tachycardia is in the range of 100 to (220 – patients age) beats per minute.
 - If patient has history or 12 Lead ECG reveals Wolfe Parkinson White (WPW), DO NOT administer a Calcium Channel Blocker (e.g., Diltiazem) or Beta Blockers. Use caution with Adenosine and give only with defibrillator available.
 - **Polymorphic / Irregular Tachycardia:**
 - This situation is usually unstable and immediate defibrillation is warranted.
 - If QT length is known, use for decision-making. Prolonged QT length defined as > 500 msec.
 - QT length < 500 msec:
 - Arrhythmia more likely related to ischemia or infarction and Magnesium not likely helpful.
 - May quickly deteriorate into Ventricular Fibrillation.
 - Even when terminated by defibrillation, may recur, so follow with medication therapy.
 - QT prolongation > 500 msec:
 - Magnesium more likely to be helpful.
- Document all rhythm changes with monitor strips and obtain monitor strips with each therapeutic intervention.



Ventricular Fibrillation Pulseless Ventricular Tachycardia

Cardiac Arrest Protocol AC 3

	<p>Begin Continuous CPR Compressions Push Hard (≥ 2 inches) Push Fast (100 - 120 / min) Change Compressors every 2 minutes <i>(sooner if fatigued)</i> <i>(Limit changes / pulse checks ≤ 10 seconds)</i></p> <p>Ventilate 1 breath every 6 seconds 30:2 Compression:Ventilation if no Advanced Airway Monitor EtCO2 <i>if available</i></p> <p>AED Procedure <i>if available</i></p>
A	Defibrillation Procedure
	IV / IO Access Protocol UP 6
A	<p>Epinephrine (1:10,000) 1 mg IV / IO Repeat every 3 to 5 minutes <i>If VF / VT refractory to defibrillation, delay Epinephrine administration until after 2d defibrillation</i></p>
	Search for Reversible Causes
	<p>Continue CPR Compressions Push Hard (≥ 2 inches) Push Fast (100 - 120 / min) Change Compressors every 2 minutes <i>(sooner if fatigued)</i> <i>(Limit changes / pulse checks ≤ 10 seconds)</i></p> <p>If Rhythm Refractory Continue CPR and give Agency specific Anti-arrhythmics and Epinephrine Continue CPR up to point where you are ready to defibrillate with device charged. Repeat pattern during resuscitation.</p>
P	<p>Amiodarone 300 mg IV / IO</p> <p>May repeat if refractory Amiodarone 150 mg IV / IO</p> <p>Or</p> <p>Lidocaine 1.0 – 1.5 mg/kg IV / IO</p> <p>May repeat if refractory Lidocaine 0.75 mg/kg IV / IO</p> <p>Maximum 3 mg/kg</p> <p>Refractory Magnesium 2 gm IV / IO</p>
	<p>Defibrillation Procedure <i>If VF / VT refractory after 3 shocks consider changing vector of defibrillation pads</i></p>

Reversible Causes
Hypovolemia
Hypoxia
Hydrogen ion (acidosis)
Hypothermia
Hypo / Hyperkalemia
Tension pneumothorax
Tamponade; cardiac
Toxins
Thrombosis; pulmonary (PE)
Thrombosis; coronary (MI)

AT ANY TIME

Return of Spontaneous Circulation



Go to Post Resuscitation Protocol AC 10

 **Notify Destination or Contact Medical Control** 



Ventricular Fibrillation Pulseless Ventricular Tachycardia

Pearls

- **Team Focused Approach / Pit-Crew Approach recommended; assigning responders to predetermined tasks. Refer to optional Team Focused CPR Protocol AC 11 or development of local agency protocol.**
- **Efforts should be directed at high quality and continuous compressions with limited interruptions and early defibrillation when indicated.**
- **DO NOT HYPERVENTILATE: If no advanced airway (BIAD, ETT) compression to ventilation ratio is 30:2. If advanced airway in place, ventilate 10 breaths per minute with continuous, uninterrupted compressions.**
- **Do not interrupt compressions to place endotracheal tube. Consider BIAD first to limit interruptions.**
- **Passive oxygenation optional in agencies practicing Team Focused Approach / Pit-Crew Approach.**
- **Reassess and document BIAD and / or endotracheal tube placement and EtCO₂ frequently, after every move, and at transfer of care.**
- **IV / IO access and drug delivery is secondary to high-quality chest compressions and early defibrillation.**
- **IV access is preferred route. Follow IV or IO Access Protocol UP 6.**
- **Defibrillation:**
 - Follow manufacture's recommendations concerning defibrillation / cardioversion energy when specified.
 - Charge defibrillator during chest compressions, near the end of 2-minute cycle, to decrease peri-shock pause.
 - Following defibrillation, provider should immediately restart chest compressions with no pulse check until end of next cycle.
- **End Tidal CO₂ (EtCO₂)**
 - If EtCO₂ is < 10 mmHg, improve chest compressions. Goal is ≥ 20 mmHg.
 - If EtCO₂ spikes, typically > 40 mmHg, consider Return of Spontaneous Circulation (ROSC)
- **Special Considerations**
 - Maternal Arrest** - Treat mother per appropriate protocol with immediate notification to Medical Control and rapid transport preferably to obstetrical center if available and proximate. Place mother supine and perform Manual Left Uterine Displacement moving uterus to the patient's left side. IV/IO access preferably above diaphragm. Defibrillation is safe at all energy levels.
 - Renal Dialysis / Renal Failure** - Refer to Dialysis / Renal Failure Protocol AM 3 caveats when faced with dialysis / renal failure patient experiencing cardiac arrest.
 - Opioid Overdose** - If suspected, administer Naloxone per Overdose / Toxic Ingestion Protocol TE 7 while ensuring airway, oxygenation, ventilations, and high-quality chest compressions.
 - Drowning / Suffocation / Asphyxiation / Hanging / Lightning Strike** – Hypoxic associated cardiac arrest and prompt attention to airway and ventilation is priority followed by high-quality and continuous chest compressions and early defibrillation.
- **Magnesium Sulfate is not routinely recommended during cardiac arrest, but may help with Torsades de points, prolonged QT, low Magnesium States (malnourished / alcoholic), and suspected digitalis toxicity**
- Return of spontaneous circulation: Heart rate should be > 60 when initiating anti-arrhythmic infusions.
- Success is based on proper planning and execution. Procedures require space and patient access. Make room to work.
- Discussion with Medical Control can be a valuable tool in developing a differential diagnosis and identifying possible treatment options.



Post Resuscitation

Return of Spontaneous Circulation

Transport Destination Decision
 Post-resuscitation patient is medically complex.
 Consider facility capabilities:

- 24-hour cardiac catheterization laboratory
- Medical ICU service
- Cardiology service
- Neurology service
- Pulmonology service
- Targeted Temperature Management

Repeat Primary Assessment	
	Optimize Ventilation and Oxygenation <ul style="list-style-type: none"> • Remove Impedance Threshold Device • Respiratory Rate 10 / minute • Maintain SpO2 92 – 98% • DO NOT HYPERVENTILATE
B	<ul style="list-style-type: none"> • ETCO2 ideally 35 – 45 mm Hg
	Airway Protocol(s) AR 1, 2, 3, 4 as indicated
B	12 Lead ECG Procedure
	IV or IO Access Protocol UP 6
P	Cardiac Monitor
	Monitor Vital Signs / Reassess
	Search for reversible causes

Reversible Causes

Hypovolemia
 Hypoxia
 Hydrogen ion (acidosis)
 Hypothermia
 Hypo / Hyperkalemia

Tension pneumothorax
 Tamponade; cardiac
 Toxins
 Thrombosis; pulmonary (PE)
 Thrombosis; coronary (MI)

	Chest Pain and STEMI Protocol AC 4 if indicated
	Hypotension / Shock Protocol AM 5 as indicated
A	Optimize Systolic BP and Mean Arterial BP <ul style="list-style-type: none"> • Systolic BP > 90 mmHg • Mean Arterial BP > 65 mmHg
P	
	Appropriate Arrhythmia Protocol(s) AC 2, 6, 7 as indicated
	Seizure Protocol UP 13 as indicated
	Post Intubation BIAD Management Protocol AR 8
	Targeted Temperature Management Protocol AC 13 if available

Arrhythmias are common and usually self limiting after ROSC

If Arrhythmia Persists follow Rhythm Appropriate Protocol

	Notify Destination or Contact Medical Control	
--	--	--



Post Resuscitation

Don't hurry!! In this case haste makes waste.

Priority 1: Blood Pressure

Priority 2: 12 Lead

Priority 3: Reduce O2 amount. Target range is 94-98% SpO2

All post arrest patients are to be transported Novant New Hanover Regional Medical Center or Grand Strand Regional Medical Center.


*In the event of re-arrest during transport, do not delay transport. The LUCAS should be utilized, if appropriate, and continue transport to the original facility transporting to.

Pearls

- **Recommended Exam: Mental Status, Neck, Skin, Lungs, Heart, Abdomen, Extremities, Neuro**
- **Continue to search for potential cause of cardiac arrest during post-resuscitation care.**
- **Hyperventilation is a significant cause of hypotension and recurrence of cardiac arrest in the post resuscitation phase and must be avoided. Titrate FiO₂ to maintain SpO₂ of 92 - 98%.**
- **Pain/sedation:**
Patients requiring advanced airways and ventilation commonly experience pain and anxiety. Unrelieved pain can lead to increased catecholamine release, ischemia, immunosuppression, and prolonged hospitalization.
Ventilated patients cannot communicate pain / anxiety and providers are poor at recognizing pain / anxiety.
Vital signs such as tachycardia and / or hypertension can provide clues to inadequate sedation, however they both are not always reliable indicators of patient's lack of adequate sedation.
Pain must be addressed first, before anxiety. Opioids are typically the first line agents before benzodiazepines. Ketamine is also a reasonable first choice agent.
- **Ventilator / Ventilation strategies:**
Tailored to individual patient presentations. Medical Control can indicate different strategies above.
In general ventilation with BVM should cause chest rise. With mechanical ventilation a reasonable tidal volume should be about 6 mL/kg and peak pressures should be < 30 cmH₂O.
Continuous pulse oximetry and capnography should be maintained during transport for monitoring.
Head of bed should be maintained at least 10 – 20 degrees of elevation when possible to decrease aspiration risk.
- **EtCO₂ Monitoring:**
Initial End tidal CO₂ may be elevated immediately post-resuscitation, but will usually normalize.
Goal is 35 – 45 mmHg but avoid hyperventilation to achieve.
- Titrate fluid resuscitation and vasopressor administration to maintain SBP of 90 – 100 mmHg or Mean Arterial Pressure (MAP) of 65 – 80 mmHg.
- **STEMI (ST-Elevation Myocardial Infarction)**
Consider placing 2 IV sites in the left arm: Many PCI centers use the right radial artery for intervention.
Consider placing defibrillator pads on patient as a precaution.
Document and time-stamp facility STEMI notification and make notification as soon as possible.
Document the time of the 12-Lead ECG in the PCR as a Procedure along with the interpretation (Paramedic).
- Consider transport to facility capable of managing the post-arrest patient including hypothermia therapy, cardiology / cardiac catheterization, intensive care service, and neurology services.
- **Targeted Temperature Management (optional):**
Maintain core temperature between 32 - 36°C.
Infusion of cold saline is NOT recommended in the prehospital setting.
No evidence suggests improved survival with prehospital cooling.
- The condition of post-resuscitation patients fluctuates rapidly and continuously, and they require close monitoring. Appropriate post-resuscitation management may best be planned in consultation with Medical Control.



Team Focused CPR



AT ANY TIME

Return of Spontaneous Circulation



Go to Post Resuscitation Protocol AC 10

BLS

Third Arriving Responder
BLS or ALS

ALS

Establish Team Leader

(Hierarchy)
Fire Department or Squad Officer
EMT
First Arriving Responder

Rotate with Compressor

To prevent Fatigue and effect high quality compressions
Take direction from Team Leader

Fourth / Subsequent Arriving Responders

Take direction from Team Leader

Continue Cardiac Arrest Protocol AC 3

Establish Team Leader

(Hierarchy)
EMS ALS Personnel
Fire Department or Squad Officer
EMT
First Arriving Responder

A

Initiate Defibrillation Automated Procedure
Establish IV / IO Protocol UP 6
Administer Appropriate Medications
Establish Airway with BIAD if not in place

P

Initiate Defibrillation Manual Procedure
Continuous Cardiac Monitoring
Establish IV / IO Protocol UP 6
Administer Appropriate Medications
Establish Airway with BIAD if not in place

Continue Cardiac Arrest Protocol AC 3

Team Leader

ALS Personnel
Responsible for patient care
Responsible for briefing / counseling family

Incident Commander

Fire Department / First Responder Officer
Team Leader until ALS arrival
Manages Scene / Bystanders
Ensures high-quality compressions
Ensures frequent compressor change
Responsible for briefing family prior to ALS arrival




Team Focused CPR

Pearls

- **This protocol is optional and given only as an example. Agencies may and are encouraged to develop their own.**
- **Team Focused Approach / Pit-Crew Approach recommended; assigning responders to predetermined tasks. Refer to optional protocol or development of local agency protocol.**
- **Efforts should be directed at high quality and continuous compressions with limited interruptions and early defibrillation when indicated.**
- **DO NOT HYPERVENTILATE: If no advanced airway (BIAD, ETT), compression to ventilation ratio is 30:2. If advanced airway in place, ventilate 10 breaths per minute with continuous, uninterrupted compressions.**
- **Do not interrupt compressions to place endotracheal tube. Consider BIAD first to limit interruptions.**
- **Passive oxygenation optional in agencies practicing Team Focused Approach / Pit-Crew Approach.**
- Reassess and document BIAD and / or endotracheal tube placement and EtCO₂ frequently, after every move, and at transfer of care.
- **IV / IO access and drug delivery are secondary to high-quality chest compressions and early defibrillation.**
- **IV access is preferred route. Follow IV or IO Access Protocol UP 6.**
- **Defibrillation:** Follow manufacture's recommendations concerning defibrillation / cardioversion energy when specified.
 - Charge defibrillator during chest compressions, near the end of 2-minute cycle, to decrease peri-shock pause.
 - Following defibrillation, provider should immediately restart chest compressions with no pulse check until end of next cycle.
- Success is based on proper planning and execution. Procedures require space and patient access. Make room to work.
- Discussion with Medical Control can be a valuable tool in developing a differential diagnosis and identifying possible treatment options.



On Scene Resuscitation Termination of CPR (Optional)



Adult Cardiac Protocol Section



On Scene Resuscitation / Termination of CPR

- *** Unit MUST be in county to utilize this protocol if unit is transporting.
1. If out of county: Continue resuscitation until arrival at destination.

Pearls

- **General approach:**
 1. Determine if a terminal disease is involved?
 2. Is there an advanced directive such as a DNR / MOST form?
 3. Did the patient express to your historian any desires regarding resuscitation and if so what measures?
 4. Remember a living will is not a DNR.
- Obtain a history while resuscitation efforts are ongoing. Determine the most legitimate person on scene as your information source such as a spouse, child, or sibling or Durable Health Care Power of Attorney.
- Basic and Advanced Life Support may use for treatment decisions.



Mechanical Circulatory Support LVAD, RVAD, and Bi-VAD

History

- SAMPLE
- Bridge to transplant
- Destination therapy
- Estimated downtime
- LVAD, RVAD, Bi-Vad, TAH
- DNR, MOST, or Living Will
- Contact with LVAD coordinator

Signs and Symptoms


- Unconsciousness
- Pulseless
- Apneic
- Poor capillary refill / skin color
- AMS or decreased mental status
- No electrical activity on ECG
- No heart tones on auscultation

Differential

- See Reversible Causes below
- Infection/Sepsis
- Hypovolemia
- Cardiac arrest
- Hemorrhage

Contact VAD coordinator:

- As quickly as possible for troubleshooting and treatment advice, but do not delay emergency treatment
- Follow patient specific emergency plan if present



Adult Cardiac Protocol Section



Mechanical Circulatory Support LVAD, RVAD, and Bi-VAD

History


- SAMPLE
- Bridge to transplant
- Destination therapy
- Estimated downtime
- LVAD, RVAD, Bi-Vad, TAH
- DNR, MOST, or Living Will
- Contact with LVAD coordinator

Signs and Symptoms

- Unconsciousness
- Pulseless
- Apneic
- Poor capillary refill / skin color
- AMS or decreased mental status
- No electrical activity on ECG
- No heart tones on auscultation


Differential

- See Reversible Causes below
- Infection/Sepsis
- Hypovolemia
- Cardiac arrest
- Hemorrhage





Mechanical Circulatory Support LVAD, RVAD, and Bi-VAD



Reversible Causes
 Hypovolemia
 Hypoxia
 Hydrogen ion (acidosis)
 Hypothermia
 Hypo / Hyperkalemia
 Tension pneumothorax
 Tamponade; cardiac
 Toxins
 Thrombosis; pulmonary (PE)
 Thrombosis; coronary (MI)

Suspected Opioid Overdose
 Administer Naloxone per
 Overdose / Toxic Ingestion
 Protocol TE 7



Left Ventricular Assist Device LVAD Unresponsive or AMS

Transport to NHRMC or GSRMC.

Pearls

- **Recommended exam: Mental status, skin color, capillary refill, peripheral pulses, blood pressure.**
- **Assessment of blood flow and perfusion status:**
 - Optimal BP attained by manual BP and Doppler.
 - Automated BP devices can measure a BP in about 50% of attempts and is not reliable to assess perfusion
 - A MAP of ≥ 60 mmHg is adequate for most LVAD patients.
 - Skin color, skin temperature, capillary refill
- **Mechanical Circulatory Support devices:**
 - LVAD – Left Ventricular Assist Device
 - RVAD – Right Ventricular Assist Device
 - BiVAD – Biventricular Ventricular Assist Device
 - TAH – Total Artificial Heart
- **Reasons for use:**
 - Bridge therapy – patients awaiting transplant or anticipated recovery.
 - Destination therapy – advanced heart failure, not candidate for transplant, and will live rest of life with device.
- **Pump type and assessing pulses:**
 - Pulsatile flow pumps – older units, not commonly in use now, but generate blood flow with a pulsatile flow and patient will have a palpable pulse.
 - Continuous flow pumps – majority of pumps now used and create blood flow in a continuous stream, no pulsatile flow, so patient will not have a palpable pulse.
 - Most devices are implanted inside the chest and have an internal pump, a driveline connected from the pump to the controller unit, and a power source consisting of batteries and electrical cord for receptacles.
- **Common complications:**
 - Disconnection of power supply, either battery disconnect, or electrical cord to receptacle disconnection.
 - Driveline failure or disconnection from controller unit.
 - Controller failure
 - Blood clot formation, acute stroke, and bleeding (mucosal and gastrointestinal most common sites)
 - Infection
- **Abnormal heart rhythm:**
 - Pseudo-PEA: Normal cardiac electrical activity in a patient who is alert and well perfused with no palpable pulse.
 - Tachyarrhythmias are usually well tolerated.
- **End Tidal CO₂ (EtCO₂)**
 - If EtCO₂ is < 10 mmHg, improve chest compressions. Goal is ≥ 20 mmHg.
 - If EtCO₂ spikes, typically > 40 mmHg, consider Return of Spontaneous Circulation (ROSC)
- **Transcutaneous Pacing:**
 - Pacing is NOT effective in cardiac arrest and pacing in cardiac arrest does NOT increase chance of survival



Total Artificial Heart

History

- SAMPLE
- Bridge to transplant
- Destination therapy
- Estimated downtime
- LVAD, RVAD, Bi-Vad, TAH
- DNR, MOST, or Living Will
- Contact with LVAD coordinator

Signs and Symptoms

- Unconsciousness
- Pulseless
- Apneic
- Poor capillary refill / skin color
- AMS or decreased mental status
- No electrical activity on ECG
- No heart tones on auscultation

Differential

- See Reversible Causes below
- Infection/Sepsis
- Hypovolemia
- Cardiac arrest
- Hemorrhage

Rapid assessment
Check for signs of life
Assess for adequate perfusion

DO NOT USE ECG MONITOR

- Total Artificial Heart does not generate ECG

Criteria for Death / No Resuscitation
Review DNR / MOST Form

YES

Decomposition, Rigor mortis, Dependent lividity, Blunt force trauma
Injury incompatible with life
Extended downtime
Do not begin resuscitation
Follow Deceased Subjects Policy

NO

Pulse Present?

NO

Go to Page 2

YES

- Airway Protocol(s) AR 1, 2, 3
if indicated
- Respiratory Distress Protocol AR 4
if indicated
- Altered Mental Status Protocol UP 4
if indicated

Check Blood Pressure

Systolic BP
≥ 150 mmHg

Systolic BP < 150 mmHg
And
≥ 90 mmHg

Systolic BP
< 90 mmHg

A
Nitroglycerin 0.3 / 0.4 mg SL
Repeat every 5 minutes
as needed

Maintain SBP ≥ 90 mmHg

P
Furosemide 40 mg IV / IO
if available

May assist patient taking their antihypertensive medication

Maintain SBP ≥ 90 mmHg

DO NOT USE:

- Manual or mechanical chest compressions
- ECG/Defibrillation/Pacing/AED devices
- Vasopressor medications
- Antiarrhythmic medications

IV / IO Access Protocol
UP 6

A
Normal Saline Bolus 500 mL
IV / IO
May repeat as needed
Maximum 1 L

Notify Destination or
Contact Medical Control



Total Artificial Heart

Entering from Page 1
No Pulse Present

DO NOT USE ECG MONITOR
• Total Artificial Heart does not generate ECG

Attempt to restart Total Artificial Heart
Expose Total Artificial Heart control module
Expose Total Artificial Heart power source
Expose Total Artificial Heart driveline

Driveline:
• Ensure it is connected firmly to control module
• Ensure lines are not kinked or bent

Power source:
• Ensure all connections firmly to control module (batteries or main power source)

Battery charge:
• Check by pressing charge indicator
• Replace battery(s) if indicated

Back up Driver
• Change to back up driver *if available*

Pulse Present? — YES —> Go to Page 1

NO

A Normal Saline Bolus 500 mL IV / IO
May repeat as needed
Maximum 1 L

Pulse Present? — YES —> Go to Page 1

NO

Follow Deceased Subjects Policy



Total Artificial Heart

Transport to NHRMC or GSRMC.

Pearls

- **Recommended exam: Mental status, skin color, capillary refill, peripheral pulses, blood pressure.**
- **Assessment of blood flow and perfusion status:**
Manual and automated BP devices can measure a BP.
Skin color, skin temperature, capillary refill
- **ECG and telemetry monitoring:**
The artificial heart does not produce an ECG wave form or tracing.
Do not use the 12-Lead ECG or ECG monitoring as it will only show asystole.
- **Total Artificial Heart:**
Different than Ventricular Assist Device (LVAD, RVAD, or Bi-VAD)
The patient's left and right ventricles are removed and the artificial heart is connected to the right and left atria.
The patient is totally dependent on the artificial heart for circulatory support – the native heart is removed.
There are both a right and left side pump, driven by air, and each side driven by a separate driveline.
The drivelines are not electric, they are driven by air, so kinking can disrupt the pumping action.
Artificial heart produces a pulsatile wave form so the patient will have a palpable pulse when operational.
- **Reasons for use:**
Bridge therapy – patients awaiting transplant or anticipated recovery.
Destination therapy – advanced heart failure, not candidate for transplant, and will live rest of life with device.
- **Common complications:**
Most common is kinking or bending of the driveline(s) which stops air from moving and stops pumping action.
Disconnection of power supply, either battery disconnect, or electrical cord to receptacle disconnection.
Driveline failure or disconnection from controller unit.
Controller failure
Blood clot formation, acute stroke, and bleeding (mucosal and gastrointestinal most common sites)
Infection
- **Blood pressure:**
Optimal SBP is < 130 mmHg and > 90 mmHg.
Hypertension puts great strain on the pump and can cause blood to back up into the lungs and cause pulmonary edema and respiratory failure.
Epinephrine and vasopressors are ineffective, can cause hypertension, and may worsen the patient's condition.
- **Manual or mechanical chest compressions:**
Do not use
- **End Tidal CO₂ (EtCO₂)**
Helpful in monitoring adequate perfusion status.
- **Defibrillation/Cardioversion:**
Do not use.
- **Transcutaneous Pacing:**
Do not use.



Wearable Cardioverter Defibrillator Vest

History


- SAMPLE
- Known risk for Sudden Cardiac Death
- Risk for life-threatening arrhythmia
- No implanted defibrillator
- Heart failure – cardiomyopathy
- Decreased ejection fraction

Signs and Symptoms

- Chest pain, dyspnea
- Palpitations
- Received shock from vest
- Poor capillary refill / skin color
- AMS or decreased mental status

Differential


- See Reversible Causes below
- Arrhythmia
- Infection/Sepsis
- Hypovolemia
- Cardiac arrest
- Hemorrhage





Wearable Cardioverter Defibrillator Vest

Transport to GSRMC or NHRMC.



1. Garment


- Worn under your normal clothing, directly against skin
- Includes the electrode belt

2. Electrode Belt

- Designed to detect dangerous heart rhythms and deliver a treatment shock

3. Monitor

- Worn around waist or with shoulder strap
- Continuously records heart rate



Battery Removal



Response button

Pearls

- Recommended exam: Mental status, skin color, capillary refill, peripheral pulses, blood pressure.
- Wearable Cardioverter-Defibrillator Vest:**
 - Device is preparing to deliver a shock to the patient:**
 - Before device delivers a shock, it tests to see if patient is conscious – voice prompt instructs patient to press the “response” button (see diagram above).
 - Only the patient should press the “response” button.
 - Once a treatable arrhythmia is detected it takes between 25 and 60 seconds to deliver the shock.
- Audible and tactile warning system:**
 - The device will provide a vibration, a siren tone, and voice prompts to check if the patient is conscious and give them an opportunity to press the “response” button to abort a shock.
 - See audible warning system above.
- Reasons for use:**
 - Currently only device on the market is the Zoll LifeVest.
 - Worn by patients at risk of sudden cardiac arrest or risk of abnormal and/or lethal arrhythmia.
- Blue gel on the patient’s skin from the device:**
 - Electrode pads release a blue gel prior to treatment to improve shock conduction and reduce burning. Do not remove the gel if the vest is left in place during treatment.
 - Remove gel if vest is removed for prehospital care.
- Shock to providers:**
 - Do not touch the patient when the device is instructing you that a shock will be delivered.
 - Providers can be shocked by the device during energy delivery if provider is touching the patient.
- Removing the device for prehospital care:**
 - The device should only be removed when ECG monitor and defibrillator is available.
 - Continuous ECG monitoring and electrode pads should be in place when vest is removed.
- Defibrillation/cardioversion with vest in place:**
 - Disconnect the device from the vest before you deliver a cardioversion or defibrillation
- Transcutaneous Pacing:**
 - May be utilized with vest in place – disconnect the device from the vest before you perform transcutaneous pacing.



Allergic Reaction/ Anaphylaxis

History

- Onset and location
- Insect sting or bite
- Food allergy / exposure
- Medication allergy / exposure
- New clothing, soap, detergent
- Past history of reactions
- Past medical history
- Medication history

Signs and Symptoms

- Itching or hives
- Coughing / wheezing or respiratory distress
- Chest or throat constriction
- Difficulty swallowing
- Hypotension or shock
- Edema
- N/V

Differential

- Urticaria (rash only)
- Anaphylaxis (systemic effect)
- Shock (vascular effect)
- Angioedema (drug induced)
- Aspiration / Airway obstruction
- Vasovagal event
- Asthma or COPD
- CHF

Assess Symptom Severity / Suspected Exposure to Allergen

MILD
Skin Only

	Diphenhydramine 25 - 50 mg PO
	IV or IO Access Protocol UP 6 <i>if indicated</i>
A	Diphenhydramine 25 - 50 mg PO / IV / IM / IO
	Famotidine 20mg IV/PO (PO swallow ≥ 12 yrs)

B Monitor and Reassess
Monitor for Worsening
Signs and Symptoms

MODERATE
2 + Body Systems

	Epinephrine 1:1000 IM 0.3 – 0.5 mg Repeat every 5 minutes if no improvement
	Diphenhydramine 25 - 50 mg PO <i>See Pearls</i>
B	Albuterol Nebulizer 2.5 – 5 mg Repeat as needed x 3 <i>if indicated</i>
A	Epinephrine 1:1000 0.3 – 0.5 mg IM Repeat every 5 minutes if no improvement
	Diphenhydramine 25 - 50 mg IV / IM / IO <i>if not given PO (See Pearls)</i>

SEVERE
2 + Body Systems + hypotension
Or Isolated Hypotension

	Epinephrine 1:1000 IM 0.3 – 0.5 mg Repeat every 5 minutes if no improvement
B	Albuterol 2.5 – 5 mg Nebulizer Repeat as needed x 3 <i>if indicated</i>
A	Epinephrine 1:1000 0.3 – 0.5 mg IM Repeat every 5 minutes if no improvement
	Airway Protocol(s) AR 1 - 4 <i>if indicated</i>
	Hypotension/ Shock Protocol AM 5 <i>if indicated</i>

	IV or IO Access Protocol UP 6
A	Albuterol Nebulizer 2.5 – 5 mg +/- Ipratropium 0.5 mg (DuoNeb) Repeat as needed x 3 <i>if indicated</i>
	Famotidine 20mg IV/PO (PO swallow ≥ 12 yrs)
	Normal Saline Bolus 500 mL IV / IO Repeat as needed Maximum 2 Liters
	Methylprednisolone 125 mg IV/IM or Decadron 4 mg IV/IM
P	No improvement w/ IM Epinephrine Epinephrine IV / IO <i>See Pearls for dosing regimen</i>
	Notify Destination or Contact Medical Control

Adult Medical Protocol Section



Allergic Reaction/ Anaphylaxis

For Epi IV with no improvement after repeated doses of Epi IM

Epinephrine Calculation:

Mix 1 mg (1:1) of Epi into 500cc bag of NS
This gives you a concentration of 1 mcg per 1/2 cc
Using a 10ggt set: 10 gtts = 1cc = 2 mcg
Dose range is 1 -10 mcg, titrate to SBP >90
5 gtts per minute = 1 mcg per min
10 gtts per minute = 2 mcg per min
15 gtts per minute = 3 mcg per min
20 gtts per minute = 4 mcg per min
etc.

Pearls

- **Recommended Exam: Mental Status, Skin, Heart, Lungs, Abdominal**
- **Anaphylaxis is an acute and potentially lethal multisystem allergic reaction.**
- **Epinephrine and administration:**
Drug of choice and the FIRST drug that should be administered in acute anaphylaxis (Moderate / Severe Symptoms.)
IM Epinephrine should be administered in priority before or during attempts at IV or IO access.
- **Diphenhydramine and steroid administration:**
Diphenhydramine/ steroids have no proven benefit in Moderate/ Severe anaphylaxis.
Diphenhydramine/ steroids should NOT delay initial or repeat Epinephrine administration.
In Moderate and Severe anaphylaxis, Diphenhydramine may decrease mental status.
Diphenhydramine should NOT be given to a patient with decreased mental status and/ or a hypotensive patient as this may cause nausea, vomiting, and/ or worsening mental status.
- **Anaphylaxis unresponsive to repeat doses of IM epinephrine may require IV epinephrine administration by IV push or epinephrine infusion. Contact Medical Control for appropriate dosing.**
- **Symptom Severity Classification:**
Mild symptoms:
Flushing, hives, itching, erythema with normal blood pressure and perfusion.
Moderate symptoms:
Flushing, hives, itching, erythema plus respiratory (wheezing, dyspnea, hypoxia) or gastrointestinal symptoms (nausea, vomiting, abdominal pain) with normal blood pressure and perfusion.
Severe symptoms:
Flushing, hives, itching, erythema plus respiratory (wheezing, dyspnea, hypoxia) or gastrointestinal symptoms (nausea, vomiting, abdominal pain) with hypotension/ poor perfusion or isolated hypotension.
- **Allergic reactions may occur with only respiratory and gastrointestinal symptoms and have no rash/ skin involvement.**
- **Angioedema** is seen in moderate to severe reactions and is swelling involving the face, lips or airway structures. This can also be seen in patients taking blood pressure medications like Prinivil / Zestril (lisinopril)-typically end in -il.
- **Hereditary Angioedema** involves swelling of the face, lips, airway structures, extremities, and may cause moderate to severe abdominal pain. Some patients are prescribed specific medications to aid in reversal of swelling.
Paramedic may assist or administer this medication per patient/ package instructions.
- **Patients with moderate and severe reactions should receive a 12 lead ECG and should be continually monitored, but this should NOT delay administration of epinephrine.**
- **EMR/ EMT:**
The use of Epinephrine IM is limited to the treatment of anaphylaxis and may be given by autoinjector or manual draw up.
Administration of diphenhydramine is limited to the oral route only.
- EMT administration of beta-agonist (Albuterol) can be administered to patients currently prescribed the medication and those that are not currently prescribed the medication.
- The shorter the onset from exposure to symptoms the more severe the reaction.



Diabetic; Adult

History

- Past medical history
- Medications
- Recent blood glucose check
- Last meal

Signs and Symptoms


- Altered mental status
- Combative / irritable
- Diaphoresis
- Seizures
- Abdominal pain
- Nausea / vomiting
- Weakness
- Dehydration
- Deep / rapid breathing

Differential

- Alcohol / drug use
- Toxic ingestion
- Trauma; head injury
- Seizure
- CVA
- Altered baseline mental status

	Blood Glucose Analysis Procedure
B	12 Lead ECG Procedure <i>if indicated</i>
	IV or IO Access Protocol UP 6
P	Cardiac Monitor
	Altered Mental Status Protocol UP 4 <i>if indicated</i>
	Hypotension/ Shock Protocol AM 5 <i>if indicated</i>
	Suspected Stroke Protocol AM 7 <i>if indicated</i>
	Seizure Protocol UP 13 <i>if indicated</i>

B	<p>Blood Glucose ≤ 69 mg / dl and symptomatic No venous access Glucagon 1 – 2 mg IM Repeat in 15 minutes if needed</p>
----------	--





Diabetic; Adult

Pearls

- **Recommended exam: Mental Status, Skin, Respirations and effort, Neuro.**
- **Patients with prolonged hypoglycemia or those who are malnourished may not respond to glucagon.**
- **Do not administer oral glucose to patients who are not able to swallow or protect their airway.**
- **Quality control checks should be maintained per manufacturers recommendation for all glucometers.**
- **Patient's refusing transport to medical facility after treatment of hypoglycemia:**
 - Blood sugar must be ≥ 80 , patient has ability to eat and availability of food with responders on scene.
 - Patient must have known history of diabetes and not taking any oral diabetic agents.
 - Patient returns to normal mental status and has a normal neurological exam with no new neurological deficits.
 - Must demonstrate capacity to make informed health care decisions. See Universal Patient Care Protocol UP-1.
 - Otherwise contact medical control.
- **Hypoglycemia with Oral Agents:**
 - Patient's taking oral diabetic medications should be encouraged to allow transportation to a medical facility.
 - They are at risk of recurrent hypoglycemia that can be delayed for hours and require close monitoring even after normal blood glucose is established.
 - Not all oral agents have prolonged action so Contact Medical Control or NC Poison Control Center for advice.
 - Patient's who meet criteria to refuse care should be instructed to contact their physician immediately and consume a meal.
- **Hypoglycemia with Insulin Agents:**
 - Many forms of insulin now exist. Longer acting insulin places the patient at risk of recurrent hypoglycemia even after a normal blood glucose is established.
 - Not all insulins have prolonged action so Contact Medical Control for advice.
 - Patient's who meet criteria to refuse care should be instructed to contact their physician immediately and consume a meal.
- **Congestive Heart Failure patients who have Blood Glucose > 250:**
 - Limit fluid boluses unless patient has signs of volume depletion such as, dehydration, poor perfusion, hypotension, and/ or shock.
- In extreme circumstances with no IV / IO access and no response to glucagon, D50 can be administered rectally, Contact Medical Control for advice.



Dialysis/ Renal Failure

History


- Peritoneal or Hemodialysis
- Anemia
- Catheter access noted
- Shunt access noted
- Hyperkalemia

Signs and Symptoms

- Hypotension
- Bleeding
- Fever
- Electrolyte imbalance
- Nausea and / or vomiting
- Altered Mental Status
- Seizure
- Arrhythmia

Differential

- Congestive heart failure
- Pericarditis
- Diabetic emergency
- Sepsis
- Cardiac tamponade



Adult Medical Protocol Section



Dialysis/ Renal Failure

All dialysis patients regardless of complaint or date of dialysis **WILL** be transported to a dialysis facility such as:

1. Novant Health New Hanover Regional Medical Center
2. Grand Strand Regional Medical Center
3. McLeod Seacoast offers on site dialysis every other day. If your patient is in need of emergent dialysis and requests this facility, make sure and advise them for further direction.

Pearls

- **Recommended exam: Mental status. Neurological. Lungs. Heart. Skin.**
- **Preferably transport to a medical facility capable of providing dialysis treatment.**
- **Do not take Blood Pressure or start IV / IO in extremity which has a shunt/ fistula in place.**
- **Access of shunt indicated in the dead or near-dead patient only with no IV or IO access.**
- **If hemorrhage cannot be controlled with firm, uninterrupted direct pressure, application of tourniquet with uncontrolled dialysis fistula bleeding is indicated.**
- **Hemodialysis:**
Process which removes waste from the blood stream and occurs about three times each week.
Some patients do perform hemodialysis at home.
- **Peritoneal dialysis:**
If patient complains of fever, abdominal pain, and/ or back pain, bring the Peritoneal Dialysis fluid bag, which has drained from the abdomen, to the hospital.

Complications of Dialysis Treatment:

Hypotension:

Typically responds to small fluid bolus of 250 mL Normal Saline.
May result in angina, AMS, seizure or arrhythmia.

Filtration and decreased blood levels of some medications like some seizure medications:

Disequilibrium syndrome:

Shift of metabolic waste and electrolytes causing weakness, dizziness, nausea and/ or vomiting and seizures.

Equipment malfunction:

Air embolism.
Bleeding.
Electrolyte imbalance.
Fever.

- **Fever:**
Consider sepsis in a dialysis patient with any catheter extending outside the body.
- Always consider Hyperkalemia in all dialysis or renal failure patients.
- Sodium Bicarbonate and Calcium Chloride/ Gluconate should not be mixed. Ideally give in separate lines.
- Renal dialysis patients have numerous medical problems typically. Hypertension and cardiac disease are prevalent.



Hypertension

History

- Documented Hypertension
- Related diseases: Diabetes; CVA; Renal Failure; Cardiac Problems
- Medications for Hypertension
- Compliance with Hypertensive Medications
- Erectile Dysfunction medications
- Pregnancy

Signs and Symptoms

One of these

- Systolic BP 220 or greater
- Diastolic BP 120 or greater


AND at least one of these

- Headache
- Chest Pain
- Dyspnea
- Altered Mental Status
- Seizure

Differential

- Hypertensive encephalopathy
- Primary CNS Injury
 - Cushing's Response with Bradycardia and Hypertension
- Myocardial Infarction
- Aortic Dissection / Aneurysm
- Pre-eclampsia / Eclampsia

Hypertension is not uncommon especially in an emergency setting. Hypertension is usually transient and in response to stress and/ or pain. A hypertensive emergency is based on blood pressure along with symptoms which suggest an organ is suffering damage such as MI, CVA or renal failure. This is very difficult to determine in the pre-hospital setting in most cases. Aggressive treatment of hypertension can result in harm. Most patients, even with significant elevation in blood pressure, need only supportive care. Specific complaints such as chest pain, dyspnea, pulmonary edema or altered mental status should be treated based on specific protocols and consultation with Medical Control.



Pearls

- **Recommended Exam: Mental Status, Skin, Neck, Lung, Heart, Abdomen, Back, Extremities, Neuro**
- Elevated blood pressure is based on two to three sets of vital signs.
- Symptomatic hypertension is typically revealed through end organ dysfunction to the cardiac, CNS, or renal systems.
- All symptomatic patients with hypertension should be transported with their head elevated at 30 degrees.
- Ensure appropriate size blood pressure cuff utilized for body habitus.



Hypotension/ Shock

History

- Blood loss - vaginal or gastrointestinal bleeding, AAA, ectopic
- Fluid loss - vomiting, diarrhea, fever
- Infection
- Cardiac ischemia (MI, CHF)
- Medications
- Allergic reaction
- Pregnancy
- History of poor oral intake

Signs and Symptoms


- Restlessness, confusion
- Weakness, dizziness
- Weak, rapid pulse
- Pale, cool, clammy skin
- Delayed capillary refill
- Hypotension
- Coffee-ground emesis
- Tarry stools

Differential

- Ectopic pregnancy
- Dysrhythmias
- Pulmonary embolus
- Tension pneumothorax
- Medication effect / overdose
- Vasovagal
- Physiologic (pregnancy)
- Sepsis

	Blood Glucose Analysis Procedure
B	12 Lead ECG Procedure
	IV or IO Access Protocol UP 6
P	Cardiac Monitor
	Airway Protocol(s) <i>if indicated</i>
	Diabetic Protocol AM 2 <i>if indicated</i>

History and Exam Suggest Type of Shock



Adult Medical Protocol Section



Hypotension/ Shock

Epinephrine Calculation:

Mix 1 mg (1:1) of Epi into 500cc bag of NS
 This gives you a concentration of 1 mcg per 1/2 cc
 Using a 10gtt set: 10 gtts = 1cc = 2 mcg
 Dose range is 1 -10 mcg, titrate to SBP >90
 5 gtts per minute = 1 mcg per min
 10 gtts per minute = 2 mcg per min
 15 gtts per minute = 3 mcg per min
 20 gtts per minute = 4 mcg per min
 etc.

Pearls

- **Recommended Exam: Mental Status, Skin, Heart, Lungs, Abdomen, Back, Extremities, Neuro**
- Hypotension is defined as a systolic blood pressure less than 90. This is not always reliable and should be interpreted in context and consider patient's typical BP if known.
- Shock may be present with a normal blood pressure initially or even elevated blood pressure.
- Shock is often present with normal vital signs and may develop insidiously. Tachycardia may be the first and only sign.
- Consider all possible causes of shock and treat per appropriate protocol.
- **Hypovolemic Shock:**
 Hemorrhage, trauma, GI bleeding, ruptured aortic aneurysm or pregnancy-related bleeding.

Tranexamic Acid (TXA):

Agencies utilizing TXA must submit letters from the their receiving trauma centers for approval by the OEMS Medical Director.

Receiving trauma centers must agree to continue TXA therapy with repeat dosing.

TXA is NOT indicated and should NOT be administered where trauma occurred > 3 hours prior to EMS arrival.

Cardiogenic Shock:

Heart failure: MI, Cardiomyopathy, Myocardial contusion, Ruptured ventricular / septum / valve / toxins.

- **Distributive Shock:**
Sepsis/ Anaphylactic/ Neurogenic/ Toxins
 Hallmark is warm, dry, pink skin with normal capillary refill time and typically alert.
- **Obstructive Shock:**
 Pericardial tamponade. Pulmonary embolus. Tension pneumothorax.
 Signs may include hypotension with distended neck veins, tachycardia, unilateral decreased breath sounds or muffled heart sounds.
- **Acute Adrenal Insufficiency or Congenital Adrenal Hyperplasia:**
 Body cannot produce enough steroids (glucocorticoids/ mineralocorticoids.)
 May have primary or secondary adrenal disease, congenital adrenal hyperplasia, or more commonly have stopped a steroid like prednisone. Injury or illness may precipitate.
 Usually hypotensive with nausea, vomiting, dehydration and/ or abdominal pain.
If suspected, Paramedic should give Methylprednisolone 125 mg IM / IV / IO or Dexamethasone 10 mg IM / IV / IO. Use steroid agent specific to your drug list.
May administer prescribed steroid carried by patient IM / IV / IO. Patient may have Hydrocortisone (Cortef or Solu-Cortef). Dose: < 1y.o. give 25 mg, 1-12 y.o. give 50 mg, and > 12 y.o. give 100 mg or dose specified by patient's physician.



Medical Blood Administration

History

- Signs of massive hemorrhage
- Suspected dissecting / rupturing aneurysm (abdominal or thoracic)
- GI bleeding
- Vaginal bleeding
- Signs of intra-abdominal bleeding

Signs and Symptoms

- Restlessness, confusion
- Weakness, dizziness
- Weak, rapid pulse
- Pale, cool, clammy skin
- Delayed capillary refill
- Hypotension
- Coffee-ground emesis
- Tarry stools

Differential

- Ectopic pregnancy
- Dysrhythmias
- Pulmonary embolus
- Tension pneumothorax
- Medication effect / overdose
- Vasovagal
- Physiologic (pregnancy)
- Sepsis


If patient with **History + Criteria:**
Follow Blood Administration checklist

Whole Blood Criteria

- SBP < 70
- SBP < 90 AND HR > 110
- Age > 65: SBP < 100 AND HR > 100
- Witnessed traumatic arrest < 10 min PTA w/ continuous CPR throughout downtime
- Shock Index Criteria > 1 (SI = HR/SBP)

Checklist:

- Patient fully exposed
- Patient airway secure
- Hypoxia has been corrected
- External bleeding has been stopped
- Patient covered with blanket



Adult Medical Protocol Section



Medical Blood Administration

For any reaction, **STOP the infusion**, remove all tubing and product from the patient and save all equipment. Flush the IV line and begin fluid bolus.

Document time of reaction onset, blood volume administered, s/s and treatment provided.

Pearls

- Care should be taken to prevent hypothermia.
- Monitor patients for signs and symptoms of transfusion reaction and adverse effects, including temperature at time of infusion and 15 minutes after start.
- Consider any fluid overload issues such as CHF or patient weight (pediatrics), and monitor for signs and symptoms appropriately.
- **Allergic Reaction:** onset <15 minutes, s/s- mild skin itching or hives <25% body, Temp 100.4°F or change of >2°F from the pre-transfusion value, chills, and hives/rash >25% body. Follow AM1.
- **Febrile Transfusion Reaction:** Temp 100.4°F or change of >1.8°F from pre-transfusion value, chills, headache, facial flushing, palpitations, cough, chest tightness, increased pulse rate and/or flank pain.
- **Hemolytic Transfusion Reaction:** Immediate lysis of transfused blood can result in fever and/or tachycardia. Other symptoms can include chills, back/flank pain, nausea/vomiting, dyspnea, flushing, bleeding, and/or hypotension. Begin aggressive NS 0.9%/ LR treatment.
- **Dilutional Thrombocytopenia:** This is generally not seen with infusion of 1-2 units, unless the patient has pre-existing thrombocytopenia or disseminated intravascular coagulation.
- **Potassium intoxication (hyperkalemia):** Symptoms can include flaccidity, muscle twitching, bradycardia, EKG changes (tall peaked T waves, prolonged P-R interval, absent P waves, prolonged QRS) and/or cardiac arrest.
- **Hypocalcemia:** (from citrate toxicity that binds Ca) Symptoms can include arrhythmias, hypotension, muscle cramping, nausea, vomiting, seizure activity, and/or tingling sensation in the fingers. Patient with acute or chronic hepatic insufficiency are at relatively higher risk of citrate toxicity. To avoid, administer WB at a minimum rate of 1 Unit > 5 minutes. Treatment with Calcium Gluconate 3 gm infused slowly in a different IV/IO line if possible.
- **Transfusion Related Acute Lung Injury (TRALI):** Caused by the activation of immune cells in the lungs, leading to fluid leakage and difficulty breathing. Symptoms can include rapid onset shortness of breath and tachypnea, hypoxemia, fever/chills, cyanosis, hypotension, crackles or wheezing.
- **Transfusion- Associated Circulatory Overload (TACO):** Occurs when the volume of blood exceeds the circulatory system's capacity. Symptoms include breathing difficulty, pulmonary edema, frothy sputum, tachycardia and/or hypoxemia. Treatment with CPAP/BiPAP if applicable.



Childbirth/ Labor

History


- Due date
- Time contractions started / how often
- Rupture of membranes
- Time / amount of any vaginal bleeding
- Sensation of fetal activity
- Past medical and delivery history
- Medications
- Gravida / Para Status
- High Risk pregnancy

Signs and Symptoms

- Spasmodic pain
- Vaginal discharge or bleeding
- Crowning or urge to push
- Meconium

Differential

- Abnormal presentation
 - Buttock
 - Foot
 - Hand
- Prolapsed cord
- Placenta previa
- Abruptio placenta



Adult Obstetrical Protocol Section

Field Delivery

Go to Newly Born Protocol AO 2

Field Delivery

Oxytocin
10 units IM
Administer to mother
Maximum 10 units

See Pearls



Childbirth/ Labor

Apgar score			
	Score 2	Score 1	Score 0
A	 Pink	 Extremities blue	 Pale or blue
P	> 100 bpm	< 100 bpm	No pulse
G	Cries and pulls away	Grimaces or weak cry	No response to stimulation
A	 Active movement	 Arms, legs flexed	 No movement
R	Strong cry	Slow, irregular	No breathing

Pearls

- **Recommended Exam (of Mother): Mental Status, Heart, Lungs, Abdomen, Neuro**
- **Record APGAR at 1 minute and 5 minutes after birth. Do not delay resuscitation to obtain APGAR.**
- **If neonate requiring resuscitation, move quickly to AO 2 Newly Born Protocol**
- **After delivery, massaging the uterus (lower abdomen) will promote uterine contraction and help to control post-partum bleeding (apply uterine massage only after placenta delivery).**
- **Postpartum hemorrhage:**
 - **Pitocin (Oxytocin):**
Following field delivery, where available, administer 10 IU IM to promote uterine contraction and decrease postpartum hemorrhage.
Agencies may administer via IV or IO route per local agency medical director.
 - **Tranexamic Acid (TXA):**
Administer when postpartum hemorrhage is associated with signs and symptoms of shock.
CONTRAINDICATED where birth occurs > 3 hours prior to EMS arrival.
- **Transport or Delivery?**
Decision to transport versus remain and deliver is multifactorial and difficult. Generally it is preferable to transport.
Factors that will impact decision include: number of previous deliveries; length of previous labors; frequency of contractions; urge to push; and presence of crowning.
- **Maternal positioning for labor:**
Supine with head flat or elevated per mother's choice. Maintain flexion of both knees and hips. Elevated buttocks slightly with towel. If delivery not imminent, place mother in the left, lateral recumbent position with right side up about 10 – 20°.
- **Umbilical cord clamping and cutting:**
Place first clamp about 10 cm from infant's abdomen and second clamp about 5 cm away from first clamp.
- **Multiple Births:**
Twins occur about 1/90 births. Typically manage the same as single gestation. If imminent delivery call for additional resources, if needed. Most twins deliver at about 34 weeks so lower birth weight and hypothermia are common. Twins may share a placenta so clamp and cut umbilical cord after first delivery. Notify receiving facility immediately.
- Document all times (Contraction onset, contraction duration and frequency, delivery, APGAR 1 and 2, and placenta delivery).
- If maternal seizures occur, refer to the Obstetrical Emergencies Protocol.
- Some perineal bleeding is normal with any childbirth. Large quantities of blood or free bleeding are abnormal.



Newly Born

History

- Due date and gestational age
- Multiple gestation (twins etc.)
- Meconium / Delivery difficulties
- Congenital disease
- Medications (maternal)
- Maternal risk factors such as substance abuse or smoking


Signs and Symptoms

- Respiratory distress
- Peripheral cyanosis or mottling (normal)
- Central cyanosis (abnormal)
- Altered level of responsiveness
- Bradycardia

Differential

- Airway failure, Secretions, or Respiratory drive
- Infection
- Maternal medication effect
- Hypovolemia, Hypoglycemia, Hypothermia
- Congenital heart disease

In a non-vigorous infant whose respirations are not improving after warming, drying, and stimulating within 30 seconds, move quickly to Positive Pressure Ventilation with BVM



Adult Obstetrical Protocol Section



Newly Born

Pearls

- Recommended Exam: Quality of Cry, Muscle tone, Respirations, Heart Rate, Pulse Oximetry, and Gestational Age**
- Majority of newborns do not require resuscitation, only warming, drying, stimulating, and cord clamping.**
 - With term gestation, strong cry/ breathing, and good muscle tone, generally will not need resuscitation.
 - If no resuscitation needed, skin-to-skin contact with the mother is best way to maintain warmth of infant.
 - Maintain warmth of infant following delivery adjuncts; cap/ hat, plastic wrap, thermal mattress, radiant heat.
 - Most important vital signs in the newly born are heart rate, respirations, and respiratory effort.
 - About 10% of newborns need assistance to help them start breathing after birth.
 - About 1% of newborns require intensive resuscitation to restore/ support cardiorespiratory functions.
- Airway:**
 - Positive Pressure Ventilations with BVM is the most important treatment in a newborn with poor respirations and/ or persistent bradycardia (HR < 100 BPM).**
 - When BVM is needed, ventilation rate is 40 – 60 breaths per minute.
 - Adequacy of ventilation/ is measured mainly by increase in heart rate as well as chest rise.
 - If heart rate or respirations are not improving after 30 to 60 seconds of resuscitation, place BIAD or endotracheal tube.
 - Routine suctioning is no longer recommended, bulb suction only if needed.
- Breathing:**
 - Oxygen is not necessary initially, but if infant is not responding with increased heart rate or adequate breathing, add oxygen to the BVM.
- Circulation/ Compressions:**
 - Heart rate is critical during first few moments of life and is best monitored by 3 or 4 lead ECG, as pulse assessment is difficult in the neonate. Heart Rate is best tool for gauging resuscitation success.
 - If heart rate remains < 60 BPM after 30 to 60 seconds of BVM/ resuscitation, begin compressions.
 - With BIAD or ETT in place, compressions and ventilation should be coordinated with compression, compression, then ventilation. (3:1 ratio with all events totaling 120 per minute)
 - 2-thumbs encircling chest and supporting the back is recommended. Limit interruptions of chest compressions.
- If infant not responding to BVM, compressions, and/ or epinephrine, consider hypovolemia, pneumothorax, and/ or hypoglycemia (< 40 mg/dL).
- Document 1 and 5 minute APGAR in PCR or ePCR. DO NOT delay or interrupt resuscitation to obtain an APGAR score.
- Meconium staining:**
 - Infant born through meconium staining who is NOT vigorous:**
 - Bulb suction mouth and nose and provide positive pressure ventilation.
 - Direct endotracheal suctioning is no longer recommended.
- Expected Pulse Oximetry readings following birth:**

(Accurate only in infant NOT requiring resuscitation)

1 minute	60 – 65%
2 minutes	65 – 70%
3 minutes	70 – 75%
4 minutes	75 – 80%
5 minutes	80 – 85%
10 minutes	85 – 95%
- Pulse oximetry should be applied to the right upper arm, wrist, or palm.
- Cord clamping:**
 - Recommended to delay for 1 minute, unless infant requires resuscitation.
- Maternal sedation or narcotics will sedate infant (Naloxone NO LONGER recommended, use supportive care only).
- D10 = D50 diluted (1 ml of D50 with 4 ml of Normal Saline) or **D10 solution at 2 mL/kg IV / IO.**
- In the NEONATE, D10 is administered at 2 mL/kg. (NOT 5 mL/kg in the pediatric patient after the first month of life.)**

Apgar score

	Score 2	Score 1	Score 0
A ppearance	 Pink	 Extremities blue	 Pale or blue
P ulse	> 100 bpm	< 100 bpm	No pulse
G rimace	Cries and pulls away	Grimaces or weak cry	No response to stimulation
A ctivity	 Active movement	 Arms, legs flexed	 No movement
R espiration	Strong cry	Slow, irregular	No breathing



OB-GYN Emergency

History


- Past medical history
- Hypertension meds
- Prenatal care
- Prior pregnancies / births
- Gravida / Para

Signs and Symptoms

- Vaginal bleeding
- Abdominal pain
- Seizures
- Hypertension
- Severe headache
- Visual changes
- Edema of hands and face

Differential

- Pre-eclampsia / Eclampsia
- Placenta previa
- Placenta abruptio
- Spontaneous abortion





OB-GYN Emergency

Pearls

- **Recommended Exam: Mental Status, Abdomen, Heart, Lungs, Neuro**
- **With active seizure activity, benzodiazepine is a priority over magnesium sulfate.**
- **Midazolam 5 – 10 mg IM is effective in termination of seizures. Do not delay IM administration with difficult or no IV or IO access.**
- **Magnesium Sulfate should be administered as quickly as possible. May cause hypotension and decreased respiratory drive, but more likely in doses higher than 6 gm.**
- **Any pregnant patient involved in a MVC should be seen immediately by a physician for evaluation. Greater than 20 weeks generally require 4 to 6 hours of fetal monitoring. DO NOT suggest the patient needs an ultrasound but emphasize patient needs 4 to 6 hours of fetal monitoring.**
- **After delivery, massaging the uterus (lower abdomen) will promote uterine contraction and help to control post-partum bleeding (apply uterine massage only after placenta delivery).**
- **Postpartum or Vaginal hemorrhage:**
 - **Pitocin (Oxytocin):**

Following field delivery, where available, administer 10 IU IM to promote uterine contraction and decrease postpartum hemorrhage.

Agencies may administer via IV or IO route per local agency medical director.
 - **Tranexamic Acid (TXA):**

Administer when postpartum hemorrhage is associated with signs and symptoms of shock. **CONTRAINDICATED** where birth occurs > 3 hours prior to EMS arrival.

Vaginal hemorrhage unrelated to pregnancy, administer with signs and symptoms of shock.
- **Ectopic pregnancy:**

Implantation of fertilized egg outside the uterus, commonly in or on the fallopian tube. As fetus grows, rupture may occur. Vaginal bleeding may or may not be present. Many women with ectopic pregnancy do not know they are pregnant. Usually occurs within 5 to 10 weeks of implantation. Maintain high index of suspicion with women of childbearing age experiencing abdominal pain.
- **Preeclampsia:**

Occurs in about 6% of pregnancies. Defined by hypertension and protein in the urine. RUQ pain, epigastric pain, N/V, visual disturbances, headache, and hyperreflexia are common symptoms.

In the setting of pregnancy, hypertension is defined as a BP > 140 systolic or > 90 diastolic mmHg, or a relative increase of 30 systolic and 20 diastolic from the patient's normal (pre-pregnancy) blood pressure.

Risk factors: < 20 years of age, first pregnancy, multi-gestational pregnancy, gestational diabetes, obesity, personal or family history of gestational hypertension.
- **Eclampsia:**

Seizures occurring in the context of preeclampsia. Remember, women may not have been diagnosed with preeclampsia.
- **Maintain patient in a left lateral position, right side up 10 - 20° to minimize risk of supine hypotensive syndrome.**
- **Ask patient to quantify bleeding - number of pads used per hour.**



Blast Injury/ Incident

History

- Type of exposure (heat, gas, chemical)
- Inhalation injury
- Time of Injury
- Past medical history/ Medications
- Other trauma
- Loss of Consciousness
- Tetanus/Immunization status

Signs and Symptoms

- Hearing loss (TM rupture)
- Ocular burns/vision changes
- Multiple trauma/ penetrating trauma
- Hypotension/ shock
- Airway compromise/distress could be indicated by hoarseness/ wheezing
- Pneumothorax/ hemothorax
- Traumatic amputation (tourniquet)

Differential

- Thermal / Chemical / Electrical Burn Injury
 - Superficial (1st Degree) red – painful (Don't include in TBSA)
 - Partial Thickness (2nd Degree) blistering
 - Full Thickness (3rd Degree) painless/charred or leathery skin
- Radiation injury

Nature of Device: Agent/ Amount. Industrial Explosion. Terrorist Incident. Improvised Explosive Device.

Method of Delivery: Incendiary/ Explosive

Nature of Environment: Open / Closed.

Distance from Device: Intervening protective barrier. Other environmental hazards,

Evaluate for: Blunt Trauma/ Crush Injury/ Compartment Syndrome/ Traumatic Brain Injury/ Concussion/ Tympanic Membrane Rupture/ Abdominal hemorrhage or Evisceration, Blast Lung Injury and Penetrating Trauma.

Scene Safety/ Quantify number and Triage Patients/ Load and Go with Assessment/ Treatment Enroute

Call for help/ additional resources
Stage until scene safe

Accidental/ Intentional Explosions
(See Pearls)

	Triage Protocol UP 2 as indicated
	Age Appropriate Airway Protocol(s) AR 1, 2, 3, 5, 6 as indicated
	Multiple Trauma Protocol TB 6 if indicated
	IV and IO Access Protocol UP 6 if indicated
P	Cardiac Monitor if indicated
	Thermal Burn Protocol TB 9 Chemical and Electrical Burn Protocol TB 2 if indicated
	Crush Injury Protocol TB 3 if indicated
	Radiation Incident Protocol TB 7 if indicated
	Decontamination Procedure USP 2 if indicated
	Pain Control Protocol UP 11 if indicated

Blast Lung Injury

YES

Age Appropriate
Airway Protocol(s) AR 4, 7
as indicated

NO

Rapid Transport to appropriate destination using
Trauma and Burn:
EMS Triage and Destination Plan

Notify Destination or Contact Medical Control

Trauma and Burn Protocol Section



Blast Injury/ Incident

Pearls

- **Types of Blast Injury:**

- Primary Blast Injury: From the blast pressure (air) wave.
 - Secondary Blast Injury: Impaled objects. Debris which becomes missiles/ shrapnel.
 - Tertiary Blast Injury: Patient falling or being thrown/ pinned by debris.
 - Most Common Cause of Death: Secondary Blast Injuries.

- **Triage of Blast Injury patients:**

- Blast Injury patients with burn injuries should be triaged using the Thermal Burn/ Chemical and Electrical Burn Protocol Guidelines for Critical/ Serious/ Minor Trauma and Burns and the Trauma and Burn: EMS Triage and Destination Plan.

- Patients may be hard of hearing due to tympanic membrane rupture.

- **Care of Blast Injury Patients:**

- Patients may suffer multi-system injuries including blunt and penetrating trauma, shrapnel, barotrauma, burns, and toxic chemical exposure.

- Consider airway burns, which should prompt early and aggressive airway management as indicated.

- Cover open chest wounds with semi-occlusive dressing or commercial chest seal product.

- Use Lactated Ringers (if available) for all Critical or Serious Burns.

- Minimize IV fluids resuscitation in patients with no signs of shock or poor perfusion.

- **Blast Lung Injury:**

- Blast Lung Injury is characterized by respiratory difficulty and hypoxia. Can occur (rarely) in patients without external thoracic trauma. More likely to occur in an enclosed space or in close proximity to explosion.

- Symptoms: Dyspnea, hemoptysis, cough, chest pain, wheezing, and hemodynamic instability.

- Signs: Apnea, tachypnea, hypopnea, hypoxia, cyanosis, and diminished breath sounds.

- Air embolism should be considered and patient transported in left-lateral decubitus position.

- Blast Lung Injury patients may require early intubation but positive pressure ventilation may worsen the injury, avoid hyperventilation, which can cause further injury.

- Air transport may worsen lung injury, monitor oxygenation and ventilation closely. Tension pneumothorax may occur requiring chest decompression. Be judicious with fluids as volume overload may worsen lung injury.

- **Accidental Explosions or Intentional Explosions:**

- All explosions or blasts should be considered intentional until determined otherwise.**

- Greatest concern is potential threat for a secondary device.**

- Attempt to determine the source of the blast to include any potential threat for aerosolization of hazardous materials.

- Evaluate scene safety including the source of the blast, which may continue to spill explosive liquids or gases.

- Consider structural collapse, environmental hazard, and fire.

- Conditions that led to the initial explosion may reoccur and lead to a second explosion.

- Patients who physically able, typically will attempt to move as far away from the explosive source.

- Evaluate surroundings for suspicious items; unattended back packs or packages, or unattended vehicles.

- If patient(s) is unconscious or there is fatalities and you are evaluating patient(s) for signs of life:**

- Before moving, note if there are wires coming from the patient(s), or if it appears the patient(s) is lying on a package/ pack, or bulky item. If so, do not move the patient(s), quickly back away and immediately notify a law enforcement officer.**

- If there are no indications the patient is connected to a triggering mechanism for a secondary device, expeditiously remove the patient(s) from the scene and begin transport to the hospital.

- Protect the airway and cervical spine, however beyond the primary survey, care and a more detailed assessment should be deferred until rapid transport begins.

- If there are signs the patient was carrying the source of the blast, notify law enforcement immediately, and most likely a law enforcement officer will accompany your patient to the hospital.



Chemical and Electrical Burn

History

- Type of exposure (heat, gas, chemical)
- Inhalation injury
- Time of Injury
- Past medical history/ Medications
- Other trauma
- Loss of Consciousness
- Tetanus/Immunization status

Signs and Symptoms

- Burns, pain, swelling
- Ocular burns/ vision changes
- Loss of consciousness
- Hypotension/ shock
- Compartment syndrome
- Airway compromise/ distress could be indicated by hoarseness/ wheezing
- Electrical burn may be misleading with small contact/ external burn and major internal injury – burn/ trauma center transport is recommended

Differential

- Thermal / Chemical / Electrical Burn Injury
 - Superficial
 - (1st Degree) red – painful (Don't include in TBSA)
 - Partial Thickness
 - (2nd Degree) blistering
 - Full Thickness
 - (3rd Degree) painless/charred or leathery skin
- Radiation injury
- Blast injury

**Assure Chemical Source is NOT Hazardous to Responders.
Assure Electrical Source is NO longer in contact with patient before touching patient.**

Assess Burn/ Concomitant Injury Severity

< 5% TBSA 2nd/3rd Degree Burn
No inhalation injury, Not Intubated,
Normotensive
GCS 14 or Greater
Minor Burn

5-15% TBSA 2nd/3rd Degree Burn
Suspected inhalation injury or requiring intubation for airway stabilization
Hypotension or GCS 13 or Less
(When reasonably accessible, transport to a Burn Center)
Serious Burn

>15% TBSA 2nd/3rd Degree Burn
Burns with Multiple Trauma
Burns with definitive airway compromise
(When reasonably accessible, transport to a Burn Center)
Critical Burn

	Age Appropriate Airway Protocol(s) AR 1, 2, 3, 4, 5, 6, 7 if indicated
	IV or IO Access Protocol UP 6 Consider 2 IV sites if ≥ 15 % TBSA
	Thermal Burn Protocol TB 9
	Pain Control Protocol UP 11 if indicated
	Identify Contact Points
	Eye Involvement Irrigate Involved Eye(s) with Normal Saline + for 30 minutes Continue irrigation during transport
	Chemical Exposure/ Burn Flush Contact Area with Normal Saline for 15 minutes Continue irrigation during transport
	Decontamination Procedure USP 2 if indicated
	Age Appropriate Cardiac Protocol(s) if indicated
Rapid Transport to appropriate destination using Trauma and Burn: EMS Triage and Destination Plan	
	Notify Destination or Contact Medical Control

Trauma and Burn Protocol Section



Chemical and Electrical Burn

Pearls

- **Recommended Exam: Mental Status, HEENT, Neck, Heart, Lungs, Abdomen, Extremities, Back, and Neuro**
- **Green, Yellow, and Red in burn severity do not apply to the Start/ JumpStart Triage System.**
- **Refer to Rule of Nines.**
- **Transport and Destination:**
 - **In general, chemical and electrical burns should be transported to a burn center.**
 - **Burn center should be initial destination choice unless EMS system access is limited by time and/ or distance.**
 - **When EMS transport to burn center is limited, transport to and stabilization at local center is appropriate.**
- **Chemical Burns:**
 - Refer to Decontamination Procedure.
 - With dry powders/ substances, gently brush or wipe off prior to irrigation. Do not aerosolize by brushing too vigorously.
 - Normal Saline or Sterile Water is preferred, however if not available, do not delay irrigation and use tap water. Other water sources may be used based on availability.
 - Flush the area as soon as possible with the cleanest, most readily available water or saline solution and use copious amounts of fluids.
 - Flush contact area for a minimum of 15 minutes and continue until arrival at receiving facility.
 - **Hydrofluoric acid burns:**
 - Monitor ECG for peaked T waves, which can be sign of hypocalcemia.
 - **Eye involvement:**
 - Irrigation is recommended for a minimum of 30 minutes and continue until arrival at receiving facility.
- **Electrical Burns:**
 - **Remember the extent of the obvious external burn from an electrical source does not always reflect more extensive internal damage. Small external injury may have large internal injury.**
 - **Do not refer to wounds as an entry and exit wound.**
 - **DO NOT contact patient until you are certain the source of the electrical shock is disconnected.**
 - Attempt to locate contact points (generally there will be two or more.) A point where the patient contacted the source and a point(s) where the patient is grounded.
 - Sites will generally be full thickness (3rd).
 - Cardiac Monitor: Anticipate ventricular or atrial irregularity including VT, VF, atrial fibrillation, and/ or heart blocks.
 - Attempt to identify the nature of the electrical source (AC or DC), the amount of voltage, and the amperage the patient may have been exposed to during the electrical shock.
 - **Lightning strike:**
 - Lightning strike victims are amenable to airway, breathing, cardiac compressions, as well as early defibrillation.
 - **Use concept of reverse triage with multiple casualties. Resuscitate lightning strikes as the priority.**
 - Lightning strike victims found alive do not often deteriorate quickly.



Crush Syndrome Trauma

History

- Entrapped and crushed under heavy load > 30 minutes
- Extremity / body crushed
- Building collapse, trench collapse, industrial accident, pinned under heavy equipment

Signs and Symptoms

- Hypotension
- Hypothermia
- Abnormal ECG findings
- Pain
- Anxiety

Differential

- Entrapment without crush syndrome
- Vascular injury with perfusion deficit
- Compartment syndrome
- Altered mental status

	Age Appropriate Airway Protocol(s) AR 1 - 7 as indicated
B	12 Lead ECG Procedure
	IV or IO Access Protocol UP 6
P	Cardiac Monitor
	Multiple Trauma Protocol TB 6 if indicated
	Thermal Burn Protocol TB 9 Chemical and Electrical Burn Protocol TB 2 if indicated
	Pain Control Protocol UP 11 as indicated

Age Specific Blood Pressure indicating possible shock

Age 0 – 28 days: SBP < 60
 Ages ≥ 1 month: SBP < 70
 Age 1 – 9: SBP < 70 + (2x Age)

Ages 10 – 64: SBP < 90
 Ages ≥ 65: SBP < 100


All ages:
 Shock Index HR > SBP

P	Consider Midazolam 0.5 – 2 mg IV / IO Midazolam 1 – 2 mg IN Maximum 5 mg Pediatric: 0.1 – 0.2 mg / kg IV / IO / IN Pediatric Maximum 2 mg Over 2 – 3 minutes as needed	OR Lorazepam 1 mg/kg IV/IO/IM Pedi: 0.1 mg/kg IV/IO/PR Max: 2 mg OR Diazepam 5 mg/kg IV/IO/PR Pedi: 0.1 mg/kg IV/IO
----------	--	---

A	Normal Saline Infusion 1 Liter per hour IV / IO Pediatric: 3 x maintenance fluid rate
----------	--

A	Decrease Normal Saline Infusion 500 mL per hour IV / IO Pediatric: Maintenance fluid rate
----------	--

P	Calcium Gluconate 2 g IV / IO Or (Calcium Chloride 1 g IV / IO) Pediatric: 20 mg / kg IV / IO Over 2- 3 minutes
	Albuterol Nebulizer 2.5 – 5 mg May repeat x 3
	Sodium Bicarbonate 50 mEq IV / IO Pediatric: 1 mEq / kg IV / IO



Rapid Transport to appropriate destination using **Trauma and Burn: EMS Triage and Destination Plan**

Notify Destination or Contact Medical Control

Exit to Age Appropriate Cardiac Arrest Protocol AC 3 / PC 4 Arrhythmia Protocol(s) **if indicated**

Trauma and Burn Protocol Section



Crush Syndrome Trauma

Pearls

- **Recommended exam: Mental Status, Musculoskeletal, Neuro**
 - **Scene safety is of paramount importance as typical scenes may pose hazards to rescuers. Call for appropriate resources.**
 - **Crush Injury is a localized crush injury with systemic signs and symptoms causing muscle breakdown and release of potentially toxic muscle cell components and electrolytes into the circulation.**
 - **Crush syndrome typically manifests after 1 – 4 hours of crush injury.**
 - **Fluid resuscitation strategy:**
 - **If possible, administer IV / IO fluids prior to release of crushed body part, especially with crush > 1 hour. If access to patient and initiation of IV / IO fluids occurs after 2 hours, give 2 liters of IV fluids in adults and 20 mL/kg of IV fluids in pediatrics, and then begin > 2 hour dosing regimen.**
 - **If not able to perform IV / IO fluid resuscitation immediately, place tourniquet on crushed limb until IV / IO fluids can be initiated (even if tourniquet is not being used for hemorrhage control).**
 - **Pediatric IV Fluid maintenance rate:**
 - **4 mL for the first 10 kg of weight +**
 - **2 mL for the second 10 kg of weight +**
 - **1 mL for every additional kg in weight after 20 kg**
- Example: 28 kg pediatric**

First 10 kg:	4 mL/kg/hr = 40 mL/hr
Second 10 kg:	2 mL/kg/hr = 20 mL/hr
Final 8 Kg:	1 mL/kg/hr = 8 mL/hr
Total: 68 mL/hr rate	
- **Consider all possible causes of shock and treat per appropriate protocol.**
 - **Majority of decompensation in pediatrics is airway or respiratory related.**
 - **Decreasing heart rate and hypotension occur late in children and are signs of impending cardiac arrest.**
 - **Shock may be present with a normal blood pressure initially or even elevated.**
 - **Shock often is present with normal vital signs and may develop insidiously. Tachycardia may be the only sign.**
 - **Patients may become hypothermic even in warm environments. Maintain warmth.**
 - **Hyperkalemia from crush syndrome can produce ECG changes described in protocol, but may also be a bizarre, wide complex rhythm. Wide complex rhythms should also be treated using the VF/ Pulseless VT Protocol if indicated (AC 9 VF Pulseless VT Protocol and/ or PC 7 Pediatric VF Pulseless VT Protocol).**



Extremity Trauma

History


- Type of injury
- Mechanism: crush/ penetrating/ amputation
- Time of injury
- Open vs. closed wound / fracture
- Wound contamination
- Medical history
- Medications

Signs and Symptoms

- Pain and/ or swelling
- Deformity
- Altered sensation/ motor function
- Diminished pulse/ capillary refill
- Decreased extremity temperature

Differential

- Abrasion
- Contusion
- Laceration
- Sprain
- Dislocation
- Fracture
- Amputation



Open Fracture
or
Amputated Part with Bone Fracture

- Best outcomes in patients who receive antibiotics within 60 minutes of injury

Trauma and Burn Protocol Section

Pearls

- **Recommended Exam: Mental Status, Extremity, Neuro, Perfusion**
- Peripheral neurovascular status is important to assess and document, as well as time of assessment.
- In amputations, time is critical. Transport and notify medical control immediately, so that the appropriate destination can be determined.
- Hip dislocations as well as knee and elbow fracture/ dislocations have a high incidence of vascular compromise.
- Urgently transport any injury with neurological or vascular compromise.
- Blood loss may be concealed or not apparent with extremity injuries.
- Lacerations optimally should be evaluated for repair within 6 hours from the time of injury.
- **Multiple casualty incident: Tourniquet Procedure may be considered first instead of direct pressure.**



Head Trauma

History

- Time of injury
- Mechanism (blunt vs. penetrating)
- Loss of consciousness
- Bleeding
- Past medical history
- Medications
- Evidence for multi-trauma

Signs and Symptoms

- Pain, swelling, bleeding
- Altered mental status
- Unconscious
- Respiratory distress/ failure
- Vomiting
- Major traumatic mechanism of injury
- Seizure

Differential

- Skull fracture
- Brain injury (Concussion, Contusion, Hemorrhage)
- Epidural hematoma
- Subdural hematoma
- Subarachnoid hemorrhage
- Spinal injury
- Abuse

Prevent hypoxia, hypotension, and hyperventilation

A single episode of hypoxia, hypotension, and hyperventilation increases mortality

	Age Appropriate Airway Protocol(s) AR 1, 2, 3, 5, 6 <i>if indicated</i>
	Obtain and Record GCS
	All patients Titrate target SpO2 100% Monitor HR, BP and O2 every 3-5 minutes Blood Glucose Analysis Procedure
B	Maintain EtCO2 35 – 45 mmHg
A	IV or IO Access - UP 6 <i>if indicated</i>
P	Cardiac Monitor
	Altered Mental Status - UP 4 <i>if indicated</i>
	Multiple Trauma - TB 6 <i>if indicated</i>
	Age Appropriate Hypotension/ Shock - AM 5/ PM 3 <i>if indicated</i>
	Seizure - UP 13 <i>if indicated</i>
	Spinal Motion Restriction Protocol TB 8 Procedure WTP 2 <i>if indicated</i>
	Pain Control - UP 11 <i>if indicated</i>
	Monitor and Reassess

Hyperventilation:
Hyperventilation is **NOT** recommended in patients who require BVM, BIAD, or ETT.
Maintain ventilation rate to target EtCO2 of 35 – 45 mmHg
See Pearls

Age Specific Blood Pressure indicating possible shock

Age 0 – 28 days: SBP < 60
Ages ≥ 1 month: SBP < 70
Age 1 – 9: SBP < 70 + (2x Age)

Ages 10 – 64: SBP < 90
Ages ≥ 65: SBP < 110

All ages Shock Index:
SI = HR ÷ SBP

Use Shock Index, Pediatric Adjusted (SIPA) for children <12 (see pearls)

Rapid Transport to appropriate destination using
**Trauma and Burn:
EMS Triage and Destination Plan**

Notify Destination or Contact Medical Control



Head Trauma

Eye Opening Response	Verbal Response	Motor Response
4 = Spontaneous	5 = Oriented	6 = Obeys commands
3 = To verbal stimuli	4 = Confused	5 = Localizes pain
2 = To pain	3 = Inappropriate words	4 = Withdraws from pain
1 = None	2 = Incoherent	3 = Flexion to pain or decorticate
	1 = None	2 = Extension to pain or decerebrate
		1 = None

Age	HR	SBP	SIPA cutoff value
1–3 years	70–110	90–110	1.2
4–6 years	65–110	90–110	1.2
7–12 years	60–100	100–120	1.0
> 12 years	55–90	100–135	0.9

SIPA, shock index, pediatric age-adjusted; HR, heart rate; SBP, systolic blood pressure.

Pearls

- **Recommended Exam: Mental Status, HEENT, Heart, Lungs, Abdomen, Extremities, Back, Neuro**
- **Hyoxia:**
Single episode of hypoxia can worsen head injury and double mortality.
Titrate SpO₂ as close to 100% as possible.
- **Hyperventilation in head injury requiring advanced airway:**
Hyperventilation lowers CO₂ and causes vasoconstriction leading to increased intracranial pressure (ICP).
Hyperventilation is not recommended and can worsen the brain injury.
In patients requiring BVM, BIAD, or endotracheal tube, titrate ventilation rate to EtCO₂ between 35 - 45 mmHg.
Recommended ventilation rates with advanced airways:
Infant/ Toddler: 25 breaths / minute
Children: 20 Breaths / minute
Adolescents/ Adults: 10 – 12 Breaths / minute
- **Hypotension:**
Episodes of hypotension can worsen head injury and increase mortality:
In adults, minimal SBP is at least 90 - 100 mmHg.
In pediatrics, minimal SBP is at least > 70 + (2 x the age in years).
Usually indicates shock unrelated to the head injury and should be aggressively treated, otherwise limit fluid administration.
- **GCS**
Key performance measure used in the EMS Acute Trauma Care Toolkit.
Serial assessments of GCS with ongoing assessments should be performed.
- Do not place in Trendelenburg position as this may increase ICP and worsen blood pressure.
- Poorly fitted cervical collars may also increase ICP when applied too tightly.
- In areas with short transport times, Drug Assisted Airway protocol is not recommended for patients who are spontaneously breathing and who have oxygen saturations of ≥ 90% with supplemental oxygen including BIAD/ BVM.
- Increased intracranial pressure (ICP) may cause hypertension and bradycardia (Cushing's Response).
- Consider Restraints if necessary for patient's and/ or personnel's protection per the Restraints: Physical Procedure USP 5.
- **Concussions:**
Traumatic brain injuries involving any of a number of symptoms including confusion, loss of consciousness, vomiting, or headache.
Any prolonged confusion or mental status abnormality which does not return to normal within 15 minutes or any documented loss of consciousness should be evaluated by a physician ASAP.
EMS Providers should not make return-to-play decisions when evaluating an athlete with suspected concussion. This is outside the scope of practice.



Multiple Trauma

History

- Time and mechanism of injury
- Damage to structure or vehicle
- Location in structure or vehicle
- Others injured or dead
- Speed and details of MVC
- Restraints/ protective equipment
- Past medical history
- Medications

Signs and Symptoms

- Pain, swelling
- Deformity, lesions, bleeding
- Altered mental status or unconscious
- Hypotension or shock
- Arrest

Differential (Life threatening)

- Uncontrolled hemorrhage
- Airway obstruction/ deformity
- Chest:
 - Tension pneumothorax
 - Flail chest/ Open chest wound
 - Pericardial tamponade/ Hemothorax
- Head Trauma Protocol TB 5
- Intra-abdominal bleeding
- Pelvis/ Femur/ Extremity fracture
- Spine fracture/ Cord injury
- Hypothermia

	Age Appropriate Airway Protocol(s) AR 1 - 7 as indicated
P	Chest Decompression Procedure WTP 1 if indicated
	Control External Hemorrhage Procedure(s) WTP 4, 5, 7 Place Pelvic Binding WTP 9 Splint Fractures Procedure WTP 3
	IV or IO Access Protocol UP 6
	Spinal Motion Restriction Procedure WTP 2 Spinal Motion Restriction Protocol TB 8 if indicated
	Obtain and Record GCS

TXA Indicators:
V/S parameters
for blunt or penetrating trauma:

Adult:

- SBP \leq 70 mmHg
or
- SBP \leq 90 mmHg + HR \geq 110
- Age \geq 65
SBP $<$ 100 mmHg + HR $>$ 100

VS / Perfusion Abnormal / Shock?

YES

NO

	Age Appropriate Hypotension/ Shock Protocol AM 5/ PM 3 if indicated
	Adults: TXA 1 - 2 g over 10 minutes IV / IO
P	Blood Products if indicated go to AM 6 or PM4

	Head Injury Protocol TB 5 if indicated
	Altered Mental Status Protocol UP 4 if indicated
	Pain Control Protocol UP 11 if indicated
	Extremity Trauma Protocol TB 4 if indicated
	Crush Syndrome Protocol TB 3 if indicated
	Repeat Assessment Adult Procedure
	Monitor and Reassess

Rapid Transport to appropriate destination using
Trauma and Burn:
EMS Triage and Destination Plan
Limit Scene Time \leq 15 minutes
Provide Early Notification

	Notify Destination or Contact Medical Control	
--	--	--

Trauma and Burn Protocol Section



Multiple Trauma

Pearls

- **Recommended Exam:** Mental Status, Skin, HEENT, Heart, Lung, Abdomen, Extremities, Back, Neuro
- **Items in Red Text** are key performance measures used in the EMS Acute Trauma Care Toolkit
- **Scene time** should not be delayed for procedures and all should be performed during rapid transport of unstable patients.
- **Ask all patients** if they are taking any anticoagulants and report during facility transition of care.
- **Airway:**
 - BVM and BIAD are acceptable for airway management to maintain SpO₂ of 92 – 98%.
 - Endotracheal intubation, if performed, should be completed during transport and should not delay scene time.
- **Breathing:**
 - Consider Chest Decompression with signs of shock and/ or injury to torso with evidence of tension pneumothorax.
- **Circulation:**
 - Control external hemorrhage and prevent hypothermia by keeping patient warm.
 - IV or IO access should be established during rapid transport of unstable patients.
- **Head Injury with multiple trauma (Refer to Head Trauma Protocol TB 5):**
 - Higher SBP targets are needed to maintain cerebral perfusion pressure.
 - Single episodes of Hypotension and/ or hypoxia are associated with worse outcomes in head injured patients.
 - Adult SBP target is ≥ 100 mmHg.
 - Pediatric SPB target is $\geq 70 + 2(\text{Age})$ mmHg.
- **Trauma Triad of Death:**
 - Metabolic acidosis/ Coagulopathy/ Hypothermia
 - Address by appropriate resuscitation measures and keeping patient warm, regardless of ambient temperature, which helps to treat metabolic acidosis, coagulopathy, and hypothermia.
- **Tranexamic Acid (TXA):**
 - Agencies utilizing TXA must submit letters from the their receiving trauma centers for approval by the OEMS Medical Director.
 - Receiving trauma centers must agree to continue TXA therapy with repeat dosing.
 - TXA is NOT indicated and should NOT be administered where trauma occurred > 3 hours prior to EMS arrival.
- **Trauma in Pregnancy:**
 - Providing optimal care for the mother = optimal care for the fetus.
 - After 20 weeks gestation (fundus at or above umbilicus) transport patient on left side with 10 – 20° of elevation.
- **Geriatric Trauma:**
 - Age ≥ 65 : SBP < 110 mmHg or HR > SBP may indicate shock.
 - Evaluate with a high index of suspicion, occult injuries difficult to recognize and with unexpected patient decompensation.
 - Risk of death with trauma increases after age 55.
 - Low impact mechanisms, such as ground level falls might result in severe injury especially in age over 65.
- See Regional Trauma Guidelines when declaring Trauma Activation.
- Maintain high-index of suspicion for domestic violence or abuse, pediatric non-accidental trauma, or geriatric abuse.
- **Refer to your Regional Trauma Guidelines when declaring Trauma Activation.**
- Severe bleeding from an extremity, not rapidly controlled with direct pressure, needs application of a tourniquet.
- Maintain high-index of suspicion for domestic violence or abuse, pediatric non-accidental trauma, or geriatric abuse.



Radiation Incident

History

- Type of exposure (heat, gas, chemical)
- Inhalation injury
- Time of Injury
- Past medical history/ Medications
- Other trauma
- Loss of Consciousness
- Tetanus/Immunization status


Signs and Symptoms

- Burns, pain, swelling
- Dizziness
- Loss of consciousness
- Hypotension/shock
- Airway compromise/ distress could be indicated by hoarseness/ wheezing
- Hypotension
- Thermal or Chemical Injury

Differential

- Thermal / Chemical / Electrical Burn Injury
 - Superficial (1st Degree) red – painful (Don't include in TBSA)
 - Partial Thickness (2nd Degree) blistering
 - Full Thickness (3rd Degree) painless/charred or leathery skin

Scene Safety / Quantify number and Triage Patients/ Load and Go with Assessment/ Treatment Enroute



Trauma and Burn Protocol Section


Collateral Injury: Most all injuries immediately seen will be a result of collateral injury, such as heat from the blast, trauma from concussion, treat collateral injury based on typical care for the type of injury displayed.

Qualify: Determine exposure type; external irradiation, external contamination with radioactive material, internal contamination with radioactive material.

Quantify: Determine exposure (generally measured in Grays/Gy). Information may be available from those on site who have monitoring equipment, do not delay transport to acquire this information.



Radiation Incident



Time Phases of Radiation Injury
(Exposure Dose vs Clinical Outcome)

Exposure Dose (Gy)	Prodrome Severity	Manifest Illness - Symptom Severity			Prognosis
		Hematologic	Gastrointestinal	Neurologic	
0.5 to 1.0	+	+	0	0	Survival almost certain
1.0 to 2.0	+/++	+	0	0	Survival >90 percent
2.0 to 3.5	++	++	0	0	Probable survival
3.5 to 5.5	+++	+++	+	0	Death in 50% at 3.5 to 6 wks
5.5 to 7.5	+++	+++	++	0	Death probable in 2-3 wks
7.5 to 10	+++	+++	+++	0*	Death probable in 1-2.5 wks
10 to 20	+++	+++	+++	+++	Death certain in 5-12 days
> 20	+++	+++	+++	+++ ^{AA}	Death certain in 2-5 days

Abbreviations: Gy: dose in Grey;
0: no effects; +: mild; ++: moderate; +++: severe or marked

* Hypotension
** Also cardiovascular collapse, fever, shock

Modified from: Waselenko, JK, MacVittie, TJ, Blakely, WF, et al. Medical management of the acute radiation syndrome: Recommendations of the strategic national stockpile radiation working group. Ann Int Med 2004; 140:1039.

Pearls


- The three primary methods of protection from radiation sources:**

Limiting time of exposure
Distance from
Shielding from the source

- Dealing with a patient with a radiation exposure can be a frightening experience. Do not ignore the ABC's, a dead but decontaminated patient is not a good outcome. Refer to the Decontamination Procedure USP 2 for more information.
- Normal Saline or Sterile Water is preferred, however if not available, do not delay irrigation using tap water. Other water sources may be used based on availability. Flush the area as soon as possible with the cleanest, most readily available water or saline solution using copious amounts of fluids.
- Three methods of exposure:**
 - External irradiation
 - External contamination
 - Internal contamination
- Two classes of radiation:**
 - Ionizing radiation (greater energy) is the most dangerous and is generally in one of three states:
 - Alpha Particles, Beta Particles and Gamma Rays.
 - Non-ionizing (lower energy) examples include microwaves, radios, lasers and visible light.
- Radiation burns with early presentation are unlikely, it is more likely this is a combination event with either thermal or chemical burn being presented as well as a radiation exposure. When the burn is from a radiation source, it indicates the patient has been exposed to a significant source, (> 250 rem).
- Patients experiencing radiation poisoning are not contagious. Cross contamination is only a threat with external and internal contamination.
- Typical ionizing radiation sources in the civilian setting include soil density probes used with roadway builders and medical uses such as x-ray sources as well as radiation therapy. Sources used in the production of nuclear energy and spent fuel are rarely exposure threats as is military sources used in weaponry. Nevertheless, these sources are generally highly radioactive and in the unlikely event they are the source, consequences could be significant and the patient's outcome could be grave.
- Dirty bomb ingredients generally include previously used radioactive material and are usually combined with a conventional explosive device to spread and distribute the contaminated material.
- Refer to Decontamination Procedure USP 2/ WMD and Nerve Agent Protocol TE 8 for contamination events.
- If there is a time lag between the time of exposure and the encounter with EMS, key clinical symptom evaluation includes: nausea/ vomiting, hypothermia/ hyperthermia, diarrhea, neurological/ cognitive deficits, headache, and hypotension.
- This event may require an activation of the National Radiation Injury Treatment Network (RITN). UNC Hospitals, Atrium Health Wake Forest Baptist and Duke are the RITN hospitals, with burns managed at UNC and Wake Forest.



Selective Spinal Motion Restriction



Pearls

- **Recommended Exam: Mental Status, Skin, Neck, Heart, Lungs, Abdomen, Back, Extremities, Neuro**
- **Patients meeting all the above criteria do not require spinal motion restriction. However, patients who fail one or more criteria above require spinal motion restriction, but does NOT require use of the long spine board for immobilization.**
- **Long spine boards are NOT considered standard of care in most cases of potential spinal injury. Spinal motion restriction with cervical collar, and securing patient to cot, while padding all void areas is appropriate.**
- **True spinal immobilization is not possible. Spine protection and spinal motion restriction do not equal long spine board.**
- **Spinal motion restriction is always utilized in at-risk patients. This includes cervical collar, securing to stretcher, minimizing movement/ transfers. and maintenance of in-line spine stabilization during any necessary movement/ transfers. This includes the elderly, or others with body or spine habitus preventing them from lying flat.**
- **Consider spinal motion restriction in patients with arthritis, cancer, dialysis, and underlying spine or bone disease.**
- **Range of motion (ROM) is tested by touching chin to chest (look down), extending neck (look up), and turning head from side to side (chin to each shoulder) only in patients without posterior cervical mid-line pain. ROM should NOT be assessed if patient has midline spinal tenderness. Patient's range of motion should not be assisted, they must be able to complete alone.**
- **EMR may participate in spinal motion restriction per Agency Medical Director.**
- **Immobilization on a long spine board is not necessary where:**
 - **Penetrating trauma to the head, neck or torso with no signs and/ or symptoms of spinal injury.**
- **Concerning mechanisms that may result in spinal column injury:**
 - **Fall from ≥ 3 feet and/ or ≥ 5 stairs or steps. Ground level falls in patients ≥ 65 years of age.**
 - **MVC ≥ 30 mph, rollover, and/or ejection**
 - **Motorcycle, bicycle, other mobile device, or pedestrian-vehicle crash**
 - **Diving or axial load to spine**
 - **Electric shock**



Thermal Burn

History


- Type of exposure (heat, gas, chemical)
- Inhalation injury
- Time of Injury
- Past medical history/ Medications
- Other trauma
- Loss of Consciousness
- Tetanus/Immunization status

Signs and Symptoms

- Burns, pain, swelling
- Dizziness
- Loss of consciousness
- Hypotension/shock
- Airway compromise/ distress could be indicated by hoarseness/ wheezing

Differential


- Thermal / Chemical / Electrical Burn Injury
 - Superficial (1st Degree) red – painful (Don't include in TBSA)
 - Partial Thickness (2nd Degree) blistering
 - Full Thickness (3rd Degree) painless/charred or leathery skin
- Radiation injury
- Blast injury



1. Lactated Ringers preferred over Normal Saline. Use if available, if not change over once available.



Thermal Burn




Relative percentage of body surface area (% BSA) affected by growth

Body Part	Age				
	0 yr	1 yr	5 yr	10 yr	15 yr
a = 1/2 of head	9 1/2	8 1/2	6 1/2	5 1/2	4 1/2
b = 1/2 of 1 thigh	2 3/4	3 1/4	4	4 1/4	4 1/2
c = 1/2 of 1 lower leg	2 1/2	2 1/2	2 3/4	3	3 1/4

Rule of Nines

- Rarely find a complete isolated body part that is injured as described in the Rule of Nines.
- More likely, it will be portions of one area, portions of another, and an approximation will be needed.
- For the purpose of determining the extent of serious injury, differentiate the area with minimal or 1st degree burn (superficial) from those of partial (2nd) or full (3rd) thickness burns.
- **For the purpose of determining Total Body Surface Area (TBSA) of burn, include only Partial (2nd) and Full Thickness (3rd) burns.** Report the observation of other superficial (1st degree) burns but do not include those burns in your TBSA estimate.
- Some texts will refer to 4th, 5th and 6th degree burns. There is significant debate regarding the actual value of identifying a burn injury beyond that of the superficial, partial and full thickness burn at least at the level of emergent and primary care. For our work, all are included in Full Thickness burns

Estimate spotty areas of burn by using the size of the patient's palm as 1 %




Palmer Method:
 Patient's entire palm, including fingers = 1%
 Patient's palm only = 0.5%

Pearls

- **Recommended Exam: Mental Status, HEENT, Neck, Heart, Lungs, Abdomen, Extremities, Back, and Neuro**
- **Green, Yellow, and Red In burn severity do not apply to the Start/ JumpStart Triage System.**
- **Airway considerations:**
 For systems performing RSI, Rocuronium is preferred agent (succinylcholine can be used in the first 24-hours)
 Singed nasal hairs, facial burns, and/ or carbonaceous sputum are NOT absolute indications for intubation in a burn patient.
 Utilizing non-rebreather face mask as well as NIPPV procedure are acceptable as tolerated.
- **Critical or Serious Burns:**
 > 5-15% total body surface area (TBSA) 2nd or 3rd degree burns
 3rd (full thickness) degree burns > 5% TBSA for any age group
 Circumferential burns of extremities
 Electrical or lightning injuries
 Suspicion of abuse or neglect
 Inhalation injury
 Chemical burns
 Burns of face, hands, perineum, or feet
 Require direct transport to a Burn Center. Local facility should be utilized only if distance to Burn Center is excessive or critical interventions such as airway management are not available in the field.
- Burn patients are trauma patients, evaluate for multisystem trauma.
- Assure whatever has caused the burn is no longer contacting the injury. (Stop the burning process!)
- Circumferential burns to extremities are dangerous due to potential vascular compromise secondary to soft tissue swelling.
- Burn patients are prone to hypothermia - never apply ice or cool the burn, must maintain normal body temperature.
- Evaluate the possibility of geriatric abuse with burn injuries in the elderly.
- Do not administer IM pain injections to a burn patient. IM dosing is variable in burn patients and may result in over or under dose.

Traumatic Arrest



Traumatic Arrest

Do not use the LUCAS Device.
Perform manual compressions.

Pearls

- **Recommended Exam: Mental Status, Skin, HEENT, Heart, Lung, Abdomen, Extremities, Back, Neuro**
- **Withholding resuscitative efforts with blunt and penetrating trauma victims who meet criteria is appropriate.**
- **If transport time to Trauma Center is < 15 minutes use of ECG monitor may delay resuscitation.**
- **Rhythm determination is more helpful in rural settings or where transport to nearest facility is > 15 minutes. Omit from algorithm where appropriate.**
- **Organized rhythms for the purposes of this protocol include Ventricular Tachycardia, Ventricular Fibrillation and PEA.**
- **Wide, bizarre rhythms such as Idioventricular and severely bradycardic rhythms < 40 BPM are not organized rhythms.**
- **First arriving EMS personnel should make the assessment concerning agonal respirations, pulselessness, asystole or PEA < 40, pupillary reflexes and spontaneous body movements.**
- **Efforts should be directed at high quality and continuous compressions with limited interruptions and early defibrillation when indicated. Consider early IO placement if available and difficult IV anticipated.**
- **DO NOT HYPERVENTILATE: If no advanced airway (BIAD, ETT) compressions to ventilations are 30:2. If advanced airway in place ventilate 8 – 10 breaths per minute.**
- **ALS procedures should optimally be performed during rapid transport.**
- **Time considerations:**
 - **From the time cardiac arrest is identified, if CPR is performed \geq 15 minutes with no ROSC consider termination of resuscitation.**
 - **From the time cardiac arrest is identified, if transport time to closest Trauma Center is > 15 minutes consider termination of resuscitation.**
- **Lightning strike, drowning or in situations causing hypothermia resuscitation should be initiated.**
- **Where multiple lightning strike victims are found used Reverse Triage: Begin CPR where apneic / pulseless**
- **Agencies utilizing Targeted Temperature Management Protocol should not cool the trauma patient, but rather make every effort to maintain warmth.**



Pediatric Asystole / PEA

History

- Events leading to arrest
- Estimated downtime
- SAMPLE
- Existence of terminal illness
- Airway obstruction
- Hypothermia
- Suspected abuse

Signs and Symptoms

- Pulseless
- Apneic
- No electrical activity on ECG
- No heart tones on auscultation

Differential

- Respiratory failure
- Foreign body
- Infection (croup, epiglottitis)
- Congenital heart disease
- See Reversible Causes below

Pediatric Pulseless Arrest Protocol

Criteria for Death / No Resuscitation Review DNR / MOST Form

YES

NO

Decomposition
Rigor mortis
Dependent lividity
Blunt force trauma
Injury incompatible with life
Extended downtime with asystole

Do not begin resuscitation

Follow Deceased Subjects Policy

AT ANY TIME

Return of Spontaneous Circulation



Go to Post Resuscitation Protocol

	<p>Begin Continuous CPR Compressions Push Hard ($\geq 1/3$ AP Diameter of Chest) (1.5 inches Infant / 2 inches in Children) Push Fast (100 - 120 / min) Change Compressors every 2 minutes (sooner if fatigued) (Limit changes / pulse checks ≤ 10 seconds)</p> <p>Ventilation rate: 1 breath every 2 seconds when age < 1 1 breathe every 3 seconds when age ≥ 1 15:2 Compression:Ventilation if no Advanced Airway</p>
	AED Procedure <i>if available</i>
P	Cardiac Monitor
	IV or IO Access Protocol UP 6
A	<p>Epinephrine 1:10,000 0.01 mg/kg IV / IO Maximum Single Dose 1mg Or Epinephrine 1:1000 0.1 mg / kg ETT Maximum 2.5 mg Repeat every 3 – 5 minutes</p>
	<p>Normal Saline Bolus 20 mL/kg IV / IO May repeat as needed Maximum 60 mL/kg</p>
	Search for Reversible Causes
	Blood Glucose Analysis Procedure <i>if applicable</i>

Reversible Causes

Hypovolemia
Hypoxia
Hydrogen ion (acidosis)
Hypothermia
Hypo / Hyperkalemia

Tension pneumothorax
Tamponade; cardiac
Toxins
Thrombosis; pulmonary (PE)
Thrombosis; coronary (MI)

 **Notify Destination or Contact Medical Control** 

Pediatric Cardiac Protocol Section



Pediatric Asystole / PEA

Pearls

- **Team Focused Approach / Pit-Crew Approach recommended; assigning responders to predetermined tasks.**
- **Refer to optional protocol AC 11 or development of local agency protocol.**
- **Efforts should be directed at high quality and continuous compressions with limited interruptions and early defibrillation when indicated. Compress $\geq 1/3$ anterior-posterior diameter of chest, in infants 1.5 inches and in children 2 inches.**
- **Majority of pediatric arrests stem from a respiratory insult or hypoxic event. Compressions should be coupled with ventilations.**
- **When advanced airway not in place perform 15 compressions with 2 ventilations.**
- **Use Handtevy App or Handtevy length-based pediatric resuscitation tape for medication, equipment, cardioversion, and defibrillation guidance. Pediatric paddles should be used in children < 10 kg.**
- **DO NOT HYPERVENTILATE:**
If advanced airway in place ventilate:
 - **Age < 1 year: 1 breath every 2 seconds with continuous, uninterrupted compressions.**
 - **Age ≥ 1 year: 1 breath every 3 seconds with continuous, uninterrupted compressions.**
- **Airway is a more important intervention in pediatric arrests. This should be accomplished quickly with BVM or BIAD.**
- **Patient survival is often dependent on proper ventilation and oxygenation / airway interventions.**
- **Do not interrupt compressions to place endotracheal tube. Consider BIAD first to limit interruptions.**
- **High-Quality CPR:**
 - Make sure chest compressions are being delivered at 100 – 120 / min.
 - Make sure chest compressions are adequate depth for age and body habitus.
 - Make sure you allow full chest recoil with each compression to provide maximum perfusion.
 - Minimize all interruptions in chest compressions to < 10 seconds.
 - Use AED or apply ECG monitor / defibrillator as soon as available.
- **Defibrillation:** Follow manufacture's recommendations concerning defibrillation / cardioversion energy when specified.
- **End Tidal CO₂ (EtCO₂)**
 - If EtCO₂ is < 10 mmHg, improve chest compressions. Goal is ≥ 20 mmHg.
 - If EtCO₂ spikes, typically > 40 mmHg, consider Return of Spontaneous Circulation (ROSC)
- **IV / IO access and drug delivery are secondary to high-quality chest compressions and early defibrillation.**
- **IV access is preferred route. Follow IV or IO Access Protocol UP 6.**
- **Special Considerations**
 - **Maternal Arrest** - Treat mother per appropriate protocol with immediate notification to Medical Control and rapid transport preferably to obstetrical center if available and proximate. Place mother supine and perform Manual Left Uterine Displacement moving uterus to the patient's left side. IV/IO access preferably above diaphragm. Defibrillation is safe at all energy levels.
 - **Renal Dialysis / Renal Failure** - Refer to Dialysis / Renal Failure Protocol AM 3 caveats when faced with dialysis / renal failure patient experiencing cardiac arrest.
 - **Opioid Overdose** - If suspected, administer Naloxone per Overdose / Toxic Ingestion Protocol UP 7 while ensuring airway, oxygenation, ventilations, and high-quality chest compressions.
 - **Drowning / Suffocation / Asphyxiation / Hanging / Lightning Strike** – Hypoxic associated cardiac arrest and prompt attention to airway and ventilation is priority followed by high-quality and continuous chest compressions and early defibrillation.
- **Success is based on proper planning and execution. Procedures require space and patient access. Make room to work.**



Pediatric Bradycardia With a Pulse

History

- Past medical history
- Foreign body exposure
- Respiratory distress or arrest
- Apnea
- Possible toxic or poison exposure
- Congenital disease
- Medication (maternal or infant)

Signs and Symptoms

- Decreased heart rate
- Delayed capillary refill or cyanosis
- Mottled, cool skin
- Hypotension or arrest
- Altered level of consciousness

Differential

- Respiratory failure, Foreign body, Secretions, Infection (croup, epiglottitis)
- Hypovolemia (dehydration)
- Congenital heart disease
- Trauma
- Tension pneumothorax
- Hypothermia
- Toxin or medication
- Hypoglycemia
- Acidosis

Bradycardia
Typically HR < 60/min
Hypotension / AMS / Poor Perfusion / Shock

	Pediatric Airway Protocol(s) AR 5, 6 as indicated
	Identify underlying cause Search for reversible causes
P	Cardiac Monitor
	IV or IO Protocol UP 6

Heart Rate < 60/min
Persists despite oxygenation and ventilation

Exit to
Pediatric Cardiac Arrest
Protocol(s) PC 1, 4, 7

NO

	Identify underlying cause Search for reversible causes
	Blood Glucose Analysis Procedure
	IV or IO Protocol UP 6
A	Normal Saline Bolus 20 ml / kg IV / IO Repeat as needed x 3 Maximum 60 mL / kg
P	Epinephrine 1:10,000 0.01 mg/kg IV / IO Maximum Single Dose 1mg Or Epinephrine 1:1000 0.1 mg / kg ETT Maximum 2.5 mg Repeat every 5 minutes
P	Atropine 0.02 mg / kg IV / IO May repeat x 1 Minimum single dose 0.1 mg Maximum single dose 0.5 mg
P	If no improvement Consider Transcutaneous Pacing Procedure

Reversible Causes

Hypovolemia
Hypoxia
Hydrogen ion (acidosis)
Hypothermia
Hypo / Hyperkalemia
Hypoglycemia

Tension pneumothorax
Tamponade; cardiac
Toxins
Thrombosis; pulmonary (PE)
Thrombosis; coronary (MI)

Suspected Beta-Blocker or Calcium Channel Blocker

Follow Pediatric Toxicology Protocol

Notify Destination or Contact Medical Control



Pediatric Bradycardia With Poor Perfusion

Pearls

- **Recommended Exam:** Mental Status, HEENT, Skin, Heart, Lungs, Abdomen, Back, Extremities, Neuro
- Bradycardia is often associated with hypoxia so insure patent airway, breathing, and circulation as needed.
- Begin CPR immediately with persistent bradycardia and poor perfusion despite adequate oxygenation and ventilation.
- Use Handtevy App or Handtevy length-based pediatric resuscitation tape for medication, equipment, cardioversion, and defibrillation guidance. Pediatric paddles should be used in children < 10 kg.
- Rhythm should be interpreted in the context of symptoms and pharmacological treatment given only when symptomatic, otherwise monitor and reassess.
- Consider hyperkalemia with wide complex, bizarre appearance of QRS complex, and bradycardia.
- **12-Lead ECG:**
 - 12 Lead ECG not necessary to diagnose and treat
 - Obtain when patient is stable and/or following rhythm conversion.
- **Unstable condition**
 - Condition which acutely impairs vital organ function and cardiac arrest may be imminent.
 - If at any point patient becomes unstable move to unstable arm in algorithm
- **Epinephrine** is first drug choice for persistent, symptomatic bradycardia.
- **Atropine:**
 - Second choice, unless there is evidence of increased vagal tone or a primary AV conduction block, then give atropine first.
 - Ineffective and potentially harmful in cardiac transplantation. May cause paradoxical bradycardia.
- **Symptomatic Bradycardia Causing Shock or Peri-arrest Condition:**
 - If no IV or IO access immediately available, start Transcutaneous Pacing, establish IV / IO access, and then administer epinephrine.
 - Epinephrine should be administered followed Atropine if no response.
- **Symptomatic Condition:**
 - Arrhythmia is causing symptoms such as palpitations, lightheadedness, or dyspnea, but cardiac arrest is not imminent.
 - Symptomatic bradycardia usually occurs at rates < 50 beats per minute.
 - Search for underlying causes such as hypoxia or impending respiratory failure.
- **Serious Signs / Symptoms:**
 - Hypotension. Acutely altered mental status. Signs of shock / poor perfusion. Chest pain with evidence of ischemia (STEMI, T wave inversions or depressions.) Acute CHF.
- **Transcutaneous Pacing Procedure (TCP):**
 - Indicated with unstable bradycardia unresponsive to medical therapy.
 - If time allows transport to specialty center because transcutaneous pacing is a temporizing measure.
 - Transvenous / permanent pacemaker will probably be needed.
 - Immediate TCP with high-degree AV block (2d or 3d degree) with no IV / IO access.
- Most maternal medications pass through breast milk to the infant so maintain high-index of suspicion for OD-toxins.
- Hypoglycemia, severe dehydration and narcotic effects may produce bradycardia. Many other agents a child ingests can cause bradycardia, often is a single dose.



Pediatric Pulmonary Edema / CHF

History


- Congenital Heart Disease
- Chronic Lung Disease
- Congestive heart failure
- Past medical history

Signs/Symptoms

- Infant: Respiratory distress, poor feeding, lethargy, weight gain, +/- cyanosis
- Child/Adolescent: Respiratory distress, bilateral rales, apprehension, orthopnea, jugular vein distention (rare), pink, frothy sputum, peripheral edema, diaphoresis, chest pain
- Hypotension, shock

Differential

- Congestive heart failure
- Asthma
- Anaphylaxis
- Aspiration
- Pleural effusion
- Pneumonia
- Pulmonary embolus
- Pericardial tamponade
- Toxic Exposure



Pearls

- **Recommended exam: Mental status, Respiratory, Cardiac, Skin, Neuro**
- **Contact Medical Control early in the care of the pediatric cardiac patient.**
- **Most children with CHF have a congenital heart defect, obtain a precise past medical history.**
- **Congenital heart disease varies by age:**
 - < 1 month: Tetralogy of Fallot, Transposition of the great arteries, Coarctation of the aorta.
 - 2 – 6 months: Ventricular septal defects (VSD), Atrioseptal defects (ASD).
 - Any age: Myocarditis, Pericarditis, SVT, heart blocks.
- **Treatment of Congestive Heart Failure / Pulmonary edema may vary depending on the underlying cause and may include the following with consultation by Medical Control:**
 - Morphine Sulfate: 0.1 mg/kg IV / IO. Max single dose 5mg/dose**
 - Fentanyl: 1 mcg/kg IV / IO. Max single dose 50 mcg.**
 - Nitroglycerin: Dose determined after consultation of Medical Control.**
 - Lasix 1 mg/kg IV / IO.**
 - Agency specific vasopressor.**
- Do not assume all wheezing is pulmonary, especially in a cardiac child: avoid albuterol unless strong history of recurrent wheezing secondary to pulmonary etiology (discuss with Medical Control)



Pediatric Cardiac Arrest

History


- Time of arrest
- Medical history
- Medications
- Possibility of foreign body
- Hypothermia

Signs and Symptoms

- Unresponsive
- Cardiac arrest

Differential

- Respiratory failure: Foreign body, Secretions, Infection (croup, epiglottitis)
- Hypovolemia (dehydration)
- Congenital heart disease
- Trauma
- Tension pneumothorax, cardiac tamponade, pulmonary embolism
- Hypothermia
- Toxin or medication
- Electrolyte abnormalities (Glucose, K)
- Acidosis





Pediatric Cardiac Arrest

Pearls

- **Team Focused Approach / Pit-Crew Approach recommended; assigning responders to predetermined tasks. Refer to optional protocol or development of local agency protocol.**
- **Efforts should be directed at high quality and continuous compressions with limited interruptions and early defibrillation when indicated. Compress $\geq 1/3$ anterior-posterior diameter of chest, in infants 1.5 inches and in children 2 inches.**
- **Majority of pediatric arrests stem from a respiratory insult or hypoxic event. Compressions should be coupled with ventilations.**
- **When advanced airway not in place perform 15 compressions with 2 ventilations.**
- **Use Handtevy App or length-based pediatric resuscitation system for medication, equipment, cardioversion, and defibrillation guidance. Pediatric paddles should be used in children < 10 kg.**
- **DO NOT HYPERVENTILATE:**
If advanced airway in place ventilate:
 - **Age < 1 year: 1 breath every 2 seconds with continuous, uninterrupted compressions.**
 - **Age ≥ 1 year: 1 breath every 3 seconds with continuous, uninterrupted compressions.**
- **Patient survival is often dependent on proper ventilation and oxygenation / airway Interventions.**
- **Do not interrupt compressions to place endotracheal tube. Consider BIAD first to limit interruptions.**
- **High-Quality CPR:**
 - Make sure chest compressions are being delivered at 100 – 120 / min.
 - Make sure chest compressions are adequate depth for age and body habitus.
 - Make sure you allow full chest recoil with each compression to provide maximum perfusion.
 - Minimize all interruptions in chest compressions to < 10 seconds.
 - Use AED or apply ECG monitor / defibrillator as soon as available.
- **Defibrillation:**
 - Follow manufacture's recommendations concerning defibrillation / cardioversion energy when specified.
 - Charge defibrillator during chest compressions, near the end of 2-minute cycle, to decrease peri-shock pause.
 - Following defibrillation, provider should immediately restart chest compressions with no pulse check until end of next cycle.
- **End Tidal CO₂ (EtCO₂)**
 - If EtCO₂ is < 10 mmHg, improve chest compressions. Goal is ≥ 20 mmHg.
 - If EtCO₂ spikes, typically > 40 mmHg, consider Return of Spontaneous Circulation (ROSC)
- **IV / IO access and drug delivery are secondary to high-quality chest compressions and early defibrillation.**
- **IV access is preferred route. Follow IV or IO Access Protocol UP 6.**
- **Special Considerations**
 - **Maternal Arrest** - Treat mother per appropriate protocol with immediate notification to Medical Control and rapid transport preferably to obstetrical center if available and proximate. Place mother supine and perform Manual Left Uterine Displacement moving uterus to the patient's left side. IV/IO access preferably above diaphragm. Defibrillation is safe at all energy levels.
 - **Renal Dialysis / Renal Failure** - Refer to Dialysis / Renal Failure Protocol AM 3 caveats when faced with dialysis / renal failure patient experiencing cardiac arrest.
 - **Opioid Overdose** - If suspected, administer Naloxone per Overdose / Toxic Ingestion Protocol UP 7 while ensuring airway, oxygenation, ventilations, and high-quality chest compressions.
 - **Drowning / Suffocation / Asphyxiation / Hanging / Lightning Strike** – Hypoxic associated cardiac arrest and prompt attention to airway and ventilation is priority followed by high-quality and continuous chest compressions and early defibrillation.
- **Success is based on proper planning and execution. Procedures require space and patient access. Make room to work.**



Pediatric Tachycardia

Narrow Complex (≤ 0.09 sec)

History


- Past medical history
- Medications or Toxic Ingestion (Aminophylline, Diet pills, Thyroid supplements, Decongestants, Digoxin)
- Drugs (nicotine, cocaine)
- Congenital Heart Disease
- Respiratory Distress
- Syncope or Near Syncope

Signs and Symptoms

- Heart Rate: Child > 180 /bpm
Infant > 220 /bpm
- Pale or Cyanosis
- Diaphoresis
- Tachypnea
- Vomiting
- Hypotension
- Altered Level of Consciousness
- Pulmonary Congestion
- Syncope

Differential

- Heart disease (Congenital)
- Hypo / Hyperthermia
- Hypovolemia or Anemia
- Electrolyte imbalance
- Anxiety / Pain / Emotional stress
- Fever / Infection / Sepsis
- Hypoxia, Hypoglycemia
- Medication / Toxin / Drugs (see HX)
- Pulmonary embolus
- Trauma, Tension Pneumothorax





Pediatric Tachycardia

Narrow Complex (≤ 0.09 sec)

Pearls

- **Recommended Exam: Mental Status, Skin, Neck, Lung, Heart, Abdomen, Back, Extremities, Neuro**
- **Monomorphic QRS:**
All QRS complexes in a single lead are similar in shape.
- **Polymorphic QRS:**
QRS complexes in a single lead will change from complex to complex.
- **Use length-based or weight-based pediatric resuscitation system for medication, equipment, cardioversion, and defibrillation guidance. Pediatric paddles should be used in children < 10 kg.**
- **Rhythm should be interpreted in the context of symptoms and pharmacological or electrical treatment given only when symptomatic, otherwise monitor and reassess.**
- **12-Lead ECG:**
12-Lead ECG not necessary to diagnose and treat.
Obtain when patient is stable and/or following rhythm conversion.
When administering adenosine, obtaining a continuous 12-Lead can be helpful to physicians.
- **Unstable condition:**
Condition which acutely impairs vital organ function and cardiac arrest may be imminent.
If at any point patient becomes unstable move to unstable arm in algorithm
If IV or IO access is in place, may administer adenosine and repeat, prior to synchronized cardioversion.
- **Document all rhythm changes with monitor strips and obtain monitor strips with each therapeutic intervention.**
- **Serious Signs and Symptoms:**
Respiratory distress / failure.
Signs of shock / poor perfusion with or without hypotension.
AMS
Sudden collapse with rapid, weak pulse
- **Narrow Complex Tachycardia (≤ 0.09 seconds):**
Sinus tachycardia: P waves present. Variable R-R waves. Infants usually < 220 beats / minute. Children usually < 180 beats / minute.
SVT: > 90 % of children with SVT will have a narrow QRS (≤ 0.09 seconds.) P waves absent or abnormal. R-R waves not variable. Usually abrupt onset. Infants usually > 220 beats / minute. Children usually > 180 beats / minute.
Atrial Flutter / Fibrillation
- **Vagal Maneuvers:**
Breath holding. Blowing a glove into a balloon. Have child blow out "birthday candles" or through an obstructed straw. Infants: May put a bag of ice water over the upper half of the face careful not to occlude the airway.
- **Separating the child from the caregiver may worsen the child's clinical condition.**
- **Monitor for respiratory depression and hypotension associated if Diazepam, Lorazepam, or Midazolam is used.**
- **Continuous pulse oximetry is required for all SVT Patients if available.**



Pediatric Tachycardia

Wide Complex (> 0.09 sec)

History


- Past medical history
- Medications or Toxic Ingestion (Aminophylline, Diet pills, Thyroid supplements, Decongestants, Digoxin)
- Drugs (nicotine, cocaine)
- Congenital Heart Disease
- Respiratory Distress
- Syncope or Near Syncope

Signs and Symptoms

- Heart Rate: Child > 180/bpm
Infant > 220/bpm
- Pale or Cyanosis
- Diaphoresis
- Tachypnea
- Vomiting
- Hypotension
- Altered Level of Consciousness
- Pulmonary Congestion
- Syncope

Differential

- Heart disease (Congenital)
- Hypothermia/ Hyperthermia
- Hypovolemia or Anemia
- Electrolyte imbalance
- Anxiety/ Pain/ Emotional stress
- Fever/ Infection/ Sepsis
- Hypoxia, Hypoglycemia
- Medication/ Toxin/ Drugs (see HX)
- Pulmonary embolus
- Trauma, Tension Pneumothorax



Pediatric Cardiac Protocol Section



Pediatric Tachycardia

Wide Complex (> 0.09 sec)

Pearls

- **Recommended Exam: Mental Status, Skin, Neck, Lung, Heart, Abdomen, Neuro**
- **Monomorphic QRS:**
 - All QRS complexes in a single lead are similar in shape.
- **Polymorphic QRS:**
 - QRS complexes in a single lead will change from complex to complex.
- Use length-based or weight-based pediatric resuscitation system for medication, equipment, cardioversion, and defibrillation guidance. Pediatric paddles should be used in children < 10 kg.
- Rhythm should be interpreted in the context of symptoms and pharmacological or electrical treatment given only when symptomatic, otherwise monitor and reassess.
- **12-Lead ECG:**
 - 12-Lead ECG is not necessary to diagnose and treat arrhythmia. A single lead ECG is often all that is needed. Obtain 12-Lead when patient is stable and/ or following a rhythm conversion.
 - When administering adenosine, obtaining a continuous 12-Lead can be helpful later to physicians.
- **Unstable condition:**
 - Condition which acutely impairs vital organ function and cardiac arrest may be imminent.
 - If at any point patient becomes unstable move to unstable arm in algorithm
- Document all rhythm changes with monitor strips and obtain monitor strips with each therapeutic intervention.
- **Serious Signs and Symptoms:**
 - Respiratory distress/ failure.
 - Signs of shock/ poor perfusion with or without hypotension.
 - AMS
 - Sudden collapse with rapid, weak pulse
- **Serious Signs and Symptoms:**
 - Respiratory distress/ failure.
 - Signs of shock/ poor perfusion with or without hypotension.
 - AMS
 - Sudden collapse with rapid, weak pulse
- **Wide Complex Tachycardia (≥ 0.09 seconds):**
 - SVT with aberrancy.
 - VT: Uncommon in children. Rates may vary from near normal to > 200/ minute.
 - Most children with VT have underlying heart disease / cardiac surgery/ long QT syndrome/ cardiomyopathy.
 - **Amiodarone 5 mg / kg over 20 – 60 minutes or Procainamide 15 mg / kg over 30 – 60 minutes IV / IO** are recommended agents. They should not be administered together. Consultation with Medical Control is advised when these agents are considered.
- **Torsade's de Pointes/ Polymorphic (multiple shaped) Tachycardia:**
 - Rate is typically 150 to 250 beats/ minute.
 - Associated with long QT syndrome, hypomagnesaemia, hypokalemia, many cardiac drugs.
 - May quickly deteriorate to VT.
 - Separating the child from the caregiver may worsen the child's clinical condition.
- Monitor for respiratory depression and hypotension associated if Diazepam, Lorazepam, or Midazolam is used.
- Continuous pulse oximetry is required for all SVT patients if available.



Pediatric Ventricular Fibrillation Pulseless Ventricular Tachycardia

History

- Events leading to arrest
- Estimated downtime
- Past medical history
- Medications
- Existence of terminal illness
- Airway obstruction
- Hypothermia

Signs and Symptoms

- Unresponsive
- Cardiac Arrest

Differential

- Respiratory failure / Airway obstruction
- Hyper / hypokalemia, Hypovolemia
- Hypothermia, Hypoglycemia, Acidosis
- Tension pneumothorax, Tamponade
- Toxin or medication
- Thrombosis: Coronary / Pulmonary Embolism
- Congenital heart disease

 Pediatric Pulseless Arrest Protocol PC 4

	<p>Begin Continuous CPR Compressions Push Hard ($\geq 1/3$ AP Diameter of Chest) (1.5 inches Infant / 2 inches in Children) Push Fast (100 - 120 / min) Change Compressors every 2 minutes (sooner if fatigued) <i>(Limit changes / pulse checks ≤ 10 seconds)</i></p> <p>Ventilation rate: 1 breath every 2 seconds when age < 1 1 breathe every 3 seconds when age ≥ 1 15:2 Compression:Ventilation if no Advanced Airway</p>
	Automated Defibrillation Procedure
P	<p>Defibrillation Manual Procedure</p> <ul style="list-style-type: none"> • First shock: 2 J / Kg • Second shock: 4 J / Kg • Subsequent shocks ≥ 4 J / kg <p>Maximum 10 J / kg or adult dose</p>
	 IV / IO Protocol UP 6
A	<p>Epinephrine 1:10,000 0.01 mg/kg IV / IO Maximum 1mg Or Epinephrine 1:1000 0.1 mg / kg ETT Maximum 2.5 mg Repeat every 3 – 5 minutes</p>
	<p>If Rhythm Refractory to defibrillation</p> <ul style="list-style-type: none"> • Continue CPR and give Agency specific Anti-arrhythmic(s) in a drug-shock-drug-shock pattern. • Continue CPR up to point where you are ready to defibrillate with device charged. <p>Repeat pattern during resuscitation.</p>
P	<p>Fist Dose - Lidocaine 1 mg / kg IV / IO Second dose - Lidocaine 20-50 mcg / kg / min</p>

AT ANY TIME

Return of Spontaneous Circulation



Go to Post Resuscitation Protocol

Reversible Causes

Hypovolemia
 Hypoxia
 Hydrogen ion (acidosis)
 Hypothermia
 Hypo / Hyperkalemia
 Hypoglycemia

Tension pneumothorax
 Tamponade; cardiac
 Toxins
 Thrombosis; pulmonary (PE)
 Thrombosis; coronary (MI)

 **Notify Destination or Contact Medical Control** 

Pediatric Cardiac Protocol Section



Pediatric Ventricular Fibrillation Pulseless Ventricular Tachycardia

Pearls

- **Team Focused Approach / Pit-Crew Approach recommended; assigning responders to predetermined tasks. Refer to optional protocol or development of local agency protocol.**
- **Efforts should be directed at high quality and continuous compressions with limited interruptions and early defibrillation when indicated. Compress $\geq 1/3$ anterior-posterior diameter of chest, in infants 1.5 inches and in children 2 inches.**
- **Majority of pediatric arrests stem from a respiratory insult or hypoxic event. Compressions should be coupled with ventilations.**
- **When advanced airway not in place perform 15 compressions with 2 ventilations.**
- **Use Handtevy length-based pediatric resuscitation system for medication, equipment, cardioversion, and defibrillation guidance. Pediatric paddles should be used in children < 10 kg.**
- **DO NOT HYPERVENTILATE:**
If advanced airway in place ventilate:
 Age < 1 year: 1 breath every 2 seconds with continuous, uninterrupted compressions.
 Age ≥ 1 year: 1 breath every 3 seconds with continuous, uninterrupted compressions.
- **Patient survival is often dependent on proper ventilation and oxygenation / airway Interventions.**
- **Do not interrupt compressions to place endotracheal tube. Consider BIAD first to limit interruptions.**
- **High-Quality CPR:**
 Make sure chest compressions are being delivered at 100 – 120 / min.
 Make sure chest compressions are adequate depth for age and body habitus.
 Make sure you allow full chest recoil with each compression to provide maximum perfusion.
 Minimize all interruptions in chest compressions to < 10 seconds.
 Use AED or apply ECG monitor / defibrillator as soon as available.
- **Defibrillation:**
 Follow manufacture's recommendations concerning defibrillation / cardioversion energy when specified.
 Charge defibrillator during chest compressions, near the end of 2-minute cycle, to decrease peri-shock pause.
 Following defibrillation, provider should immediately restart chest compressions with no pulse check until end of next cycle.
- **End Tidal CO₂ (EtCO₂)**
 If EtCO₂ is < 10 mmHg, improve chest compressions. Goal is ≥ 20 mmHg.
 If EtCO₂ spikes, typically > 40 mmHg, consider Return of Spontaneous Circulation (ROSC)
- **IV / IO access and drug delivery are secondary to high-quality chest compressions and early defibrillation.**
- **IV access is preferred route. Follow IV or IO Access Protocol UP 6.**
- **Special Considerations**
 - **Maternal Arrest** - Treat mother per appropriate protocol with immediate notification to Medical Control and rapid transport preferably to obstetrical center if available and proximate. Place mother supine and perform Manual Left Uterine Displacement moving uterus to the patient's left side. IV/IO access preferably above diaphragm. Defibrillation is safe at all energy levels.
 - **Renal Dialysis / Renal Failure** - Refer to Dialysis / Renal Failure Protocol AM 3 caveats when faced with dialysis / renal failure patient experiencing cardiac arrest.
 - **Opioid Overdose** - If suspected, administer Naloxone per Overdose / Toxic Ingestion Protocol UP 7 while ensuring airway, oxygenation, ventilations, and high-quality chest compressions.
 - **Drowning / Suffocation / Asphyxiation / Hanging / Lightning Strike** – Hypoxic associated cardiac arrest and prompt attention to airway and ventilation is priority followed by high-quality and continuous chest compressions and early defibrillation.
- **Success is based on proper planning and execution. Procedures require space and patient access. Make room to work.**



Pediatric Post Resuscitation

History

- Respiratory arrest
- Cardiac arrest

Signs/Symptoms

- Return of pulse

Differential

- Continue to address specific differentials associated with the original dysrhythmia

Transport Destination Decision

Post-resuscitation patient is medically complex.

Consider facility capabilities:

- Pediatric ICU service
- Pediatric Cardiology service
- Pediatric Neurology service
- Targeted Temperature Management

	Pediatric Airway Protocol(s) AR 5 - 7 <i>as needed</i>
	Monitor Vital Signs / Reassess
	Blood Glucose Analysis Procedure
	Optimize Ventilation and Oxygenation <ul style="list-style-type: none"> • Maintain SpO2 ≥ 92 – 98% • Advanced airway if indicated • Age Appropriate Respiratory Rate • Remove Impedance Threshold Device DO NOT HYPERVENTILATE
	ETCO2 ideally 35 – 45 mm Hg
B	12 Lead ECG Procedure
	IV or IO Protocol UP 6
P	Cardiac Monitor
	Pediatric Diabetic Protocol PM 2 <i>if indicated</i>
	Pediatric Hypotension / Shock Protocol PM 3 <i>if indicated</i>
	Pediatric Bradycardia Protocol PC 2 <i>if indicated</i>
	Pediatric Tachycardia Protocol PC 5, 6 <i>as indicated</i>

Hypotension Age Based

0 – 31 Days
< 60 mmHg


1 Month to 1 Year
< 70 mmHg

> than 1 Year
< 70 + (2 x age) mmHg

Arrhythmias are common and usually self limiting after ROSC



If Arrhythmia Persists follow Rhythm Appropriate Protocol



	Continue Antiarrhythmic Utilized Refer to Appropriate Pediatric Arrhythmia Protocol
P	Agency Specific Antiarrhythmic

	Post-intubation / BIAD Management Protocol AR 8
--	---

	Notify Destination or Contact Medical Control	
--	--	--



Pediatric Post Resuscitation

Pearls

- **Recommended Exam: Mental Status, Neck, Skin, Lungs, Heart, Abdomen, Extremities, Neuro**
- **Goals of care are to preserve neurologic function, prevent secondary organ damage, treat the underlying cause of illness, and optimize prehospital care. Frequent reassessment is necessary.**
- **Hyperventilation is a significant cause of hypotension and recurrence of cardiac arrest in the post resuscitation phase and must be avoided. Titrate FiO_2 to maintain SpO_2 of 92 - 98%.**
- **Use length-based or weight-based pediatric resuscitation system for medication, equipment, cardioversion, and defibrillation guidance. Pediatric paddles should be used in children < 10 kg.**
- **Pain/sedation:**

Patients requiring advanced airways and ventilation commonly experience pain and anxiety. Unrelieved pain can lead to increased catecholamine release, ischemia, immunosuppression, and prolonged hospitalization.

Ventilated patients cannot communicate pain / anxiety and providers are poor at recognizing pain / anxiety.

Vital signs such as tachycardia and / or hypertension can provide clues to inadequate sedation, however they both are not always reliable indicators of patient's lack of adequate sedation.

Pain must be addressed first, before anxiety. Opioids are typically the first line agents before benzodiazepines. Ketamine is also a reasonable first choice agent.
- **Ventilator / Ventilation strategies:**

Tailored to individual patient presentations. Medical Control can indicate different strategies above.

In general ventilation with BVM should cause chest rise. With mechanical ventilation a reasonable tidal volume should be about 6 mL/kg and peak pressures should be < 30 cmH₂O.

Continuous pulse oximetry and capnography should be maintained during transport for monitoring.

Head of bed should be maintained at least 10 – 20 degrees of elevation when possible to decrease aspiration risk.
- **EtCO₂ Monitoring:**

Initial End tidal CO₂ may be elevated immediately post-resuscitation, but will usually normalize.

Goal is 35 – 45 mmHg but DO NOT hyperventilate to achieve.

EtCO₂ should be continually monitored with advanced airway in place.
- Administer resuscitation fluids and vasopressor agents to maintain SBP at targets listed on page 1. This table represents minimal SBP targets.
- Targeted Temperature Management is recommended in pediatrics, but prehospital use is not associated with improved outcomes. Transport to facility capable of intensive pediatric care.
- Consider transport to facility capable of managing the post-arrest patient including hypothermia therapy, cardiology / cardiac catheterization, intensive care service, and neurology services.
- The condition of post-resuscitation patients fluctuates rapidly and continuously, and they require close monitoring. Appropriate post-resuscitation management may best be planned in consultation with Medical Control.



Pediatric Allergic Reaction

History


- Onset and location
- Insect sting or bite
- Food allergy/ exposure
- Medication allergy/ exposure
- New clothing, soap, detergent
- Past medical history/ reactions
- Medication history

Signs and Symptoms

- Itching or hives
- Coughing/ wheezing or respiratory distress
- Chest or throat constriction
- Difficulty swallowing
- Hypotension or shock
- Edema

Differential

- Urticaria (rash only)
- Anaphylaxis (systemic effect)
- Shock (vascular effect)
- Angioedema (drug induced)
- Aspiration/ Airway obstruction
- Vasovagal event
- Asthma/ COPD /CHF





Pediatric Allergic Reaction

Known or suspected COVID:

Terbutaline can be administered alone, without Albuterol /Atrovent, if COVID is suspected or known. This decision is left up to the provider.

If Atrovent and/or Albuterol are administered a N95 mask or P100 mask **MUST** be worn by the Providers.

Terbutaline Dosage:

Pediatric: 0.01mg/kg (up to 0.25mg) SQ May repeat as needed every 15 minutes.

EPI Infusion:

Mix 1mg of **EPI 1:1mg** into a 500cc bag of NS. This concentration is 2mcg per 1mL. Use a 10 gtt set. Example for 10 kg child: $0.1 \text{ mcg/kg} \times 10 \text{ kg} = 1 \text{ mcg} - 0.5\text{cc} = 5 \text{ gtt}/\text{min}$

Pearls

- **Recommended Exam: Mental Status, Skin, Heart, Lungs, Abdomen**
- **Anaphylaxis is an acute and potentially lethal multisystem allergic reaction.**
- **Epinephrine administration:**
Drug of choice and the **FIRST** drug that should be administered in acute anaphylaxis (**Moderate/ Severe Symptoms.**)
IM Epinephrine should be administered in priority before or during attempts at IV or IO access.
- **Diphenhydramine and steroid administration:**
Diphenhydramine/ steroids have **no proven benefit in Moderate/ Severe anaphylaxis.**
Diphenhydramine/ steroids should **NOT** delay initial or repeat Epinephrine administration.
In **Moderate and Severe anaphylaxis, Diphenhydramine may decrease mental status.**
Diphenhydramine should **NOT** be given to a patient with decreased mental status and/ or a hypotensive patient as this may cause nausea, vomiting, and/ or worsening mental status.
- **Anaphylaxis unresponsive to repeat doses of IM epinephrine may require IV epinephrine administration by IV push or epinephrine infusion. Contact Medical Control for appropriate dosing.**
- **Symptom Severity Classification:**
Mild symptoms:
Flushing, hives, itching, erythema with normal blood pressure and perfusion.
Moderate symptoms:
Flushing, hives, itching, erythema plus respiratory (wheezing, dyspnea, hypoxia) or gastrointestinal symptoms (nausea, vomiting, abdominal pain) with normal blood pressure and perfusion.
Severe symptoms:
Flushing, hives, itching, erythema plus respiratory (wheezing, dyspnea, hypoxia) or gastrointestinal symptoms (nausea, vomiting, abdominal pain) with hypotension and poor perfusion.
- **Allergic reactions may occur with only respiratory and gastrointestinal symptoms and have no rash/ skin involvement.**
- **Angioedema** is seen in moderate to severe reactions and is swelling involving the face, lips or airway structures. This can also be seen in patients taking blood pressure medications like Prinivil / Zestril (lisinopril)-typically end in -il.
- **Hereditary Angioedema** involves swelling of the face, lips, airway structures, extremities, and may cause moderate to severe abdominal pain. Some patients are prescribed specific medications to aid in reversal of swelling. **Paramedic may assist or administer this medication per patient/ package instructions.**
- **Fluids and Medication titrated to maintain a SBP $>70 + (\text{age in years} \times 2)$ mmHg.**
- **Patients with moderate and severe reactions should receive a 12-Lead ECG and should be continually monitored, but this should NOT delay administration of epinephrine.**
- **EMR/ EMT:**

The use of Epinephrine IM is limited to the treatment of anaphylaxis and may be given by autoinjector or manual draw up.

Administration of diphenhydramine is limited to the oral route only.

- EMT administration of beta-agonist (Albuterol) may be administered to patients currently prescribed the medication and to patients who are not currently prescribed the medication.
- The shorter the onset from exposure to symptoms the more severe the reaction.



Pediatric Diabetic

History

- Past medical history
- Medications
- Recent blood glucose check
- Last meal

Signs and Symptoms


- Altered mental status
- Combative/ irritable
- Diaphoresis
- Seizures
- Abdominal pain
- Nausea/ vomiting
- Weakness
- Dehydration
- Deep/ rapid breathing

Differential

- Alcohol/ drug use
- Toxic ingestion
- Trauma; head injury
- Seizure
- CVA
- Altered baseline mental status.

	Blood Glucose Analysis Procedure
B	12 Lead ECG Procedure <i>if indicated</i>
	IV or IO Access Protocol UP 6
P	Cardiac Monitor
	Altered Mental Status Protocol UP 4 <i>if indicated</i>
	Hypotension/ Shock Protocol AM 5 <i>if indicated</i>
	Seizure Protocol UP 13 <i>if indicated</i>

B	<p>Blood glucose ≤ 69 mg/dl Symptomatic with NO IV / IO</p> <p>Access: Awake, alert and able to tolerate oral agent:</p> <p>Give oral glucose solution.</p> <p>If unable to tolerate oral: Glucagon 0.1 mg/kg IM (Maximum 1 mg)</p> <p>Repeat every 15 minutes as needed to keep Blood glucose > 60 mg / dl.</p>
----------	--



Pediatric Medical Protocol Section



Pediatric Diabetic

Pearls

- **Recommended Exam: Mental Status, HEENT, Skin, Respirations and effort, Abdomen, Neuro.**
- **Patients with prolonged hypoglycemia or those who are malnourished may not respond to glucagon.**
- **Do not administer oral glucose to patients that are not able to swallow or protect their airway.**
- **Quality control checks should be maintained per manufacturers recommendation for all glucometers.**
- **D10/ D25 Preparation:**
 - D10: Remove 10 mL of D50 from a D50 vial. Add 40 mL of NS with the 10 mL of D50 with a total volume of 50 mL.
 - D10: Alternative, Discard 40 mL from the D50 vial and draw up 40 mL of NS with a total volume of 50 mL.
 - D25: Remove 25 mL of D50 and draw up 25 mL of NS with a total volume of 50 mL.
- **Patient's refusing transport to medical facility after treatment of hypoglycemia:**
 - Adult caregiver must be present with pediatric patient.
 - Blood sugar must be ≥ 80 , patient has ability to eat and availability of food with responders on scene.
 - Patient must have known history of diabetes and not taking any oral diabetic agents.
 - Patient returns to normal mental status and has a normal neurological exam with no new neurological deficits.
 - Must demonstrate capacity to make informed health care decisions. See Universal Patient Care Protocol UP-1.
 - Otherwise contact medical control.
- **Hypoglycemia with Oral Agents:**
 - Patients taking oral diabetic medications should be strongly encouraged to allow transportation to a medical facility.
 - They are at risk of recurrent hypoglycemia that can be delayed for hours and require close monitoring even after normal blood glucose is established.
 - Not all oral agents have prolonged action so Contact Medical Control or NC Poison Control Center for advice.
 - Patients who meet criteria to refuse care should be instructed to contact their physician immediately and consume a meal.
- **Hypoglycemia with Insulin Agents:**
 - Many forms of insulin now exist. Longer acting insulin places the patient at risk of recurrent hypoglycemia even after a normal blood glucose is established.
 - Not all insulins have prolonged action so Contact Medical Control for advice. Patients who meet criteria to refuse care should be instructed to contact their physician immediately and consume a meal.
- In extreme circumstances with no IV and no response to glucagon, Dextrose 50 % can be administered rectally. Contact medical control for advice.



Pediatric Hypotension/ Shock

History

- Blood loss
- Fluid loss
- Vomiting
- Diarrhea
- Fever
- Infection

Signs and Symptoms

- Restlessness, confusion, weakness
- Dizziness
- Tachycardia
- Hypotension (Late sign)
- Pale, cool, clammy skin
- Delayed capillary refill
- Dark-tarry stools

Differential

- Shock
 - Hypovolemic
 - Cardiogenic
 - Septic
 - Neurogenic
 - Anaphylactic
- Trauma
- Infection
- Dehydration
- Congenital heart disease
- Medication or Toxin

	Blood Glucose Analysis Procedure
	IV or IO Access Protocol UP 6
P	Cardiac Monitor
	Pediatric Airway Protocol(s) <i>if indicated</i>
	Diabetic Protocol PM 2 <i>if indicated</i>


Age Specific Blood Pressure indicating possible shock

Age 0 – 28 days: SBP < 60
 Ages ≥ 1 month: SBP < 70
 Age 1 – 9: SBP < 70 + (2x Age)

Ages 10 – 64: SBP < 90

All ages:
 Shock Index Criteria >1:
 SI = HR / SBP

History and Exam Suggest Type of Shock





Hypotension/ Shock

Whole Blood Criteria:

- Age 0-28 days: SBP < 60
- Age 1-12 months: SBP < 70
- Age 1-10 Years: SBP <70 + (age in years x2)
- Age > 10 Years: SBP <90
OR
- Witnessed traumatic arrest <10min PTA w/ continuous CPR throughout downtime
- OR
Shock Index Criteria >1
SI = HR/SBP
*Monitor for Transfusion-associated circulatory overload (TACO) and allergic reactions.

Epi Infusion:

You will mix 1mg (1:1) of Epi into a 500cc NS.
This gives you a concentration of 1mcg per 1/2 cc.

Using a **10gtt set:** 1cc = 2 mcg

Example:

0.1 mcg/kg x 10kg = 1 mcg = 0.5cc = 5 gtts

0.1 mcg/kg x 20kg = 2 mcg = 1cc = 10 gtts

0.1 mcg/kg x 30kg = 3 mcg = 1.5cc = 15 gtts

Pearls

- **Recommended Exam: Mental Status, Skin, Heart, Lungs, Abdomen, Back, Extremities, Neuro**
- Consider all possible causes of shock and treat per appropriate protocol. Majority of decompensation in pediatrics is airway or respiratory related.
- Decreasing heart rate and hypotension occur late in children and are signs of impending cardiac arrest.
- Shock may be present with a normal blood pressure initially or even elevated.
- Shock often is present with normal vital signs and may develop insidiously. Tachycardia may be the first and only sign.
- Consider all possible causes of shock and treat per appropriate protocol.
- **Hypovolemic Shock:**
Hemorrhage, trauma, GI bleeding, or pregnancy-related bleeding.
Tranexamic Acid (TXA):
Agencies utilizing TXA must submit letters from their receiving trauma centers for approval by the OEMS Medical Director.
Receiving trauma centers must agree to continue TXA therapy with repeat dosing.
TXA is NOT indicated and should NOT be administered where trauma occurred > 3 hours prior to EMS arrival.
- **Cardiogenic Shock:**
Heart failure: MI, Cardiomyopathy, Myocardial contusion, Ruptured ventricle/ septum/ valve/ toxins.
- **Distributive Shock:**
Septic/ Anaphylactic/ Neurogenic/ Toxic
Hallmark is warm, dry, pink skin with normal capillary refill time and typically alert.
- **Obstructive Shock:**
Pericardial tamponade. Pulmonary embolus. Tension pneumothorax.
Signs may include hypotension with distended neck veins, tachycardia, unilateral decreased breath sounds or muffled heart sounds.
- **Acute Adrenal Insufficiency or Congenital Adrenal Hyperplasia:**
Body cannot produce enough steroids (glucocorticoids/ mineralocorticoids.)
May have primary or secondary adrenal disease, congenital adrenal hyperplasia, or more commonly have stopped a steroid like prednisone. Injury or illness may precipitate.
Usually hypotensive with nausea, vomiting, dehydration and/ or abdominal pain.
If suspected, Paramedic should give Methylprednisolone 125 mg IM / IV / IO or Dexamethasone 10 mg IM / IV / IO.
Use steroid agent specific to your drug list.
May administer prescribed steroid carried by patient IM / IV / IO. Patient may have Hydrocortisone (Cortef or Solu-Cortef). Dose: < 1y.o. give 25 mg, 1-12 y.o. give 50 mg, and > 12 y.o. give 100 mg or dose specified by patient's physician.



Pediatric Medical Blood Administration

History

- Signs of massive hemorrhage
- Low blood counts
- Anemia
- Recent chemotherapy
- Unstable GI bleeding

Signs and Symptoms

- Restlessness, confusion
- Weakness, dizziness
- Weak, rapid pulse
- Pale, cool, clammy skin
- Hypotension
- Delayed capillary refill
- Coffee ground emesis / tarry stools

Differential

- Trauma
- Infection
- Dehydration
- Congenital heart disease
- Medication /Toxin

If patient with **History + Criteria:**
Follow Blood Administration Checklist

Whole Blood Criteria

- Age 0-28 days: SBP < 60
- Ages ≥ 1 month: SBP < 70
- Ages 1-9: SBP < 70 +(2 x Age)
- Ages 10 - 64: SBP < 90
- All Ages Shock Index Criteria > 1 (SI = HR / SBP)

Checklist:

- Patient fully exposed
- Patient airway secured
- Hypoxia has been corrected
- External bleeding has been stopped
- Patient covered with blanket

Do NOT administer LR on same side as blood product

Do NOT delay blood for fluid bolus

A

**Initiate IV / IO
Normal Saline Bolus
20 mL / kg**

Titrate to age appropriate
SBP ≥ 70 +(2 x Age)

Maximum 60 mL / kg

P

**Whole Blood Criteria Met
Alert- Code Blood**

10 cc/kg

Calcium Gluconate

30 mg/kg IV/IO over 10 minutes

Continuously monitor patient
through all phases of
transport

**Monitor for
Transfusion Reaction**

STOP INFUSION!
Allergy Protocol PM1
if indicated



**Notify Destination or
Contact Medical Control**





Pediatric Medical Blood Administration

For any reaction, **STOP the infusion**, remove all tubing and product from the patient and save all equipment. Flush the IV line and begin fluid bolus.

Document time of reaction onset, blood volume administered, s/s and treatment provided.

Pearls

- Care should be taken to prevent hypothermia.
- Monitor patients for signs and symptoms of transfusion reaction and adverse effects, including temperature at time of infusion and 15 minutes after start.
- Consider any fluid overload issues such as CHF or patient weight (pediatrics), and monitor for signs and symptoms appropriately.
- Allergic Reaction: onset <15 minutes, s/s- mild skin itching or hives <25% body, Temp 100.4°F or change of >2°F from the pre-transfusion value, chills, and hives/rash >25% body. Follow AM1.
- Febrile Transfusion Reaction: Temp 100.4°F or change of >1.8°F from pre-transfusion value, chills, headache, facial flushing, palpitations, cough, chest tightness, increased pulse rate and/or flank pain.
- Hemolytic Transfusion Reaction: Immediate lysis of transfused blood can result in fever and/or tachycardia. Other symptoms can include chills, back/flank pain, nausea/vomiting, dyspnea, flushing, bleeding, and/or hypotension. Begin aggressive NS 0.9%/ LR treatment.
- Dilutional Thrombocytopenia: This is generally not seen with infusion of 1-2 units, unless the patient has pre-existing thrombocytopenia or disseminated intravascular coagulation.
- Potassium intoxication (hyperkalemia): Symptoms can include flaccidity, muscle twitching, bradycardia, EKG changes (tall peaked T waves, prolonged P-R interval, absent P waves, prolonged QRS) and/or cardiac arrest.
- Hypocalcemia: (from citrate toxicity that binds Ca) Symptoms can include arrhythmias, hypotension, muscle cramping, nausea, vomiting, seizure activity, and/or tingling sensation in the fingers. Patient with acute or chronic hepatic insufficiency are at relatively higher risk of citrate toxicity. To avoid, administer WB at a minimum rate of 1 Unit > 5 minutes. Treatment with Calcium Gluconate 30 mg/kg infused slowly in a different IV/IO line if possible.
- Transfusion Related Acute Lung Injury (TRALI): Caused by the activation of immune cells in the lungs, leading to fluid leakage and difficulty breathing. Symptoms can include rapid onset shortness of breath and tachypnea, hypoxemia, fever/chills, cyanosis, hypotension, crackles or wheezing.
- Transfusion- Associated Circulatory Overload (TACO): Occurs when the volume of blood exceeds the circulatory system's capacity. Symptoms include breathing difficulty, pulmonary edema, frothy sputum, tachycardia and/or hypoxemia. Treatment with CPAP/BiPAP if applicable.



Bites and Envenomations

History

- Type of bite/ sting
- Description/ photo for identification
- Time, location, size of bite/ sting
- Previous reaction to bite/ sting
- Domestic vs. Wild
- Tetanus and Rabies risk
- Immunocompromised patient

Signs and Symptoms

- Rash, skin break, wound
- Pain, soft tissue swelling, redness
- Blood oozing from the bite wound
- Evidence of infection
- Shortness of breath, wheezing
- Allergic reaction, hives, itching
- Hypotension or shock


Differential

- Animal bite
- Human bite
- Snake bite (poisonous)
- Spider bite (poisonous)
- Insect sting / bite (bee, wasp, ant, tick)
- Infection risk
- Rabies risk
- Tetanus risk

Call for help/ additional resources
Stage until scene safe

Contact
Carolinas Poison Control
1-800-222-1222
Or
Agency Specific Number

	General Wound Care Procedure
	Immobilize Injury
	Remove any constricting clothing/ bands/ jewelry
	IV or IO Access Protocol UP 6 <i>if indicated</i>
	Age Appropriate Trauma Protocol(s) TB 4, 5, 6 <i>if indicated</i>
	Age Appropriate Allergic Reaction/ Anaphylaxis Protocol AM 1/ PM 1 <i>if indicated</i>
	Age Appropriate Hypotension/ Shock Protocol AM 5 / PM 3 <i>if indicated</i>
	Pain Control Protocol UP 11 <i>if indicated</i>
	Extremity Trauma Protocol TB 4 <i>if indicated</i>





Bites and Envenomations

Muscle Spasm

Midazolam 2 – 2.5 mg IV / IO Over 2 to 3 minutes

Maximum 10 mg

Pediatrics: Midazolam 0.1 – 0.2 mg / kg IV / IO / IM / IN

Over 2 to 3 minutes Maximum 5 mg IM

OR

Lorazepam 1-2 mg IV/IO/IM

May repeat every 10 minutes as needed

Maximum 4 mg

Pediatrics: Lorazepam 0.05 - 0.1 mg/kg IV/IO/IM

Maximum 4 mg total

May repeat every 3-5 minutes as needed

OR

Diazepam 4 mg IV/IO

10 mg Per Rectum if no IV/IO access

May repeat 2mg every 3-5 minutes as needed

Maximum 10 mg total

Pediatric: Diazepam: 0.1 -0.3 mg/kg IV/IO

Maximum 4 mg

0.5 mg/kg Per Rectum every 3-5 minutes as needed

Maximum 10mg total

Pearls

- **Recommended Exam: Mental Status, Skin, Extremities (Location of injury), and a complete Neck, Lung, Heart, Abdomen, Back, and Neuro exam if systemic effects are noted**
- **Immunocompromised patients are at an increased risk for infection: diabetes, chemotherapy, transplant patients.**
- **Consider contacting the North Carolina Poison Control Center for guidance (1-800-222-1222).**
- **Do not put responders in danger attempting to capture an animal or insect for identification purposes.**
- **Evidence of infection: swelling, redness, drainage, fever, red streaks proximal to wound.**
- **Human bites:**
 - Human bites have higher infection rates than animal bites due to normal mouth bacteria.
 - Hand and foot bites have highest rates of infection.
- **Dog/ Cat/ Carnivore bites:**
 - Carnivore bites are much more likely to become infected and all have risk of Rabies exposure.
 - Cat bites may progress to infection rapidly due to a specific bacteria (*Pasteurella multocoda*).
- **Snake bites:**
 - Poisonous snakes in this area are generally of the pit viper family: rattlesnake and copperhead.
 - Coral snake bites are rare: Very little pain but very toxic. "Red on yellow - kill a fellow, red on black - venom lack."
 - Amount of envenomation is variable, generally worse with larger snakes and early in spring.
 - Snake bites are treated based on signs and symptoms and progression.
 - It is not important to attempt to identify the type of snake and attempts may endanger providers.**
 - Do not bring a snake to the facility for identification as accidental bites to providers may occur.**
- **Spider bites:**
 - Black Widow spider bites tend to be minimally painful, but over a few hours, muscular pain and severe abdominal pain may develop (spider is black with red hourglass on belly).
 - Brown Recluse spider bites are minimally painful to painless. Little reaction is noted initially but tissue necrosis at the site of the bite develops over the next few days (brown spider with fiddle shape on back).
- **Animal bite(s) in subjects declining transport to a medical facility for evaluation:**
 - NCGS 130A-196 requires that all animal bites be reported to the local health department even if the bite is by the owner's animal, and even if accidental.
 - Reporting requirements can be satisfied by reporting to local animal control official.



Carbon Monoxide/ Cyanide

History

- Smoke inhalation
- Ingestion of cyanide
- Eating large quantity of fruit pits
- Industrial exposure
- Trauma
- Reason: Suicide, criminal, accidental
- Past Medical History
- Time/ Duration of exposure

Signs and Symptoms


- AMS
- Malaise, weakness, flu like illness
- Dyspnea
- GI Symptoms; N/V; cramping
- Dizziness
- Seizures
- Syncope
- Reddened skin
- Chest pain

Differential

- Diabetic related
- Infection
- MI
- Anaphylaxis
- Renal failure/ dialysis problem
- Head injury/ trauma
- Co-ingestant or exposures

	Immediately Remove from Exposure
	Appropriate Airway Protocol(s) 1 - 7 as indicated
	High Flow Oxygen
	Blood Glucose Analysis Procedure
B	12 Lead ECG Procedure
	IV or IO Access Protocol UP 6
P	Cardiac Monitor/ CO Monitor
	Altered Mental Status Protocol UP 4 if indicated
	Age Appropriate Diabetic Protocol AM 2/ PM 2 if indicated
	Age Appropriate Multiple Trauma Protocol TB 6 Head Injury TB 5 if indicated
	Age Appropriate Hypotension/ Shock Protocol AM 5/ PM 3 if indicated

**Contact
Carolinas Poison Control
1-800-222-1222
Or
Agency Specific Number**



Pearls

- **Recommended exam: Neuro, Skin, Heart, Lungs, Abdomen, Extremities**
- **Scene safety is priority.**
- Consider CO and Cyanide with any product of combustion.
- Normal environmental CO level does not exclude CO poisoning.
- Symptoms present with lower CO levels in pregnancy, children, and the elderly.
- Continue high flow oxygen regardless of pulse ox readings.



Drowning

History


- Submersion in water regardless of depth
- Possible history of trauma
- Slammed into shore wave break
- Duration of submersion/ immersion
- Temperature of water or possibility of hypothermia

Signs and Symptoms

- Unresponsive
- Mental status changes
- Decreased or absent vital signs
- Foaming/ Vomiting
- Coughing, Wheezing, Rales, Rhonchi, Stridor
- Apnea

Differential

- Trauma
- Pre-existing medical problem
 - Hypoglycemia
 - Cardiac Dysrhythmia
- Pressure injury (SCUBA diving)
 - Barotrauma
 - Decompression sickness
- Post-immersion syndrome



Toxic-Environmental Protocol Section

Pearls

- **Recommended Exam: Respiratory, Mental status, Trauma Survey, Skin, Neuro**
- **Drowning is the process of experiencing respiratory impairment (any respiratory symptom) from submersion/ immersion in a liquid.**
- **Begin with BVM ventilations, if patient does not tolerate, then apply appropriate mode of supplemental oxygen.**
- **Ensure scene safety. Drowning is a leading cause of death among would-be rescuers.**
- **When feasible, only appropriately trained and certified rescuers should remove patients from areas of danger.**
- **Regardless of water temperature – resuscitate all patients with known submersion time of ≤ 25 minutes.**
- **Regardless of water temperature – If submersion time ≥ 1 hour consider moving to recovery phase instead of rescue.**
- **Foam is usually present in airway and may be copious, DO NOT waste time attempting to suction. Ventilate with BVM through foam (suction water and vomit only when present.)**
- **Cardiac arrest in drowning is caused by hypoxia, airway and ventilation are equally important to high-quality CPR.**
- **Encourage transport of all symptomatic patients (cough, foam, dyspnea, abnormal lung sounds, hypoxia) due to potential worsening over the next 6 hours.**
- Predicting prognosis in prehospital setting is difficult and does not correlate with mental status. Unless obvious death, transport.
- Hypothermia is often associated with drowning and submersion injuries even with warm ambient conditions.
- Drowning patient typically has <1 – 3 mL/kg of water in lungs (does not require suction) Primary treatment is reversal of hypoxia.
- Spinal motion restriction is usually unnecessary. When indicated it should not interrupt ventilation, oxygenation and/ or CPR.



Hyperthermia

History

- Age, very young and old
- Exposure to increased temperatures and / or humidity
- Past medical history / Medications
- Time and duration of exposure
- Poor PO intake, extreme exertion
- Fatigue and / or muscle cramping

Signs and Symptoms

- Altered mental status / coma
- Hot, dry or sweaty skin
- Hypotension or shock
- Seizures
- Nausea

Differential

- Fever (Infection)
- Dehydration
- Medications
- Hyperthyroidism (Thyroid Storm)
- Delirium tremens (DT's)
- Heat cramps, exhaustion, stroke
- CNS lesions or tumors

Temperature Measurement Procedure **if available**

Temperature Measurement should NOT delay treatment of hyperthermia

Remove from heat source to cool environment
Cooling measures
Remove tight clothing
Blood Glucose Analysis Procedure
Age Appropriate Diabetic Protocol AM 2/ PM 2 as indicated

Heat Stroke

Classic Heat Stroke

- Not common type
- Hot and Dry
- Altered Mental Status

Exertional Heat Stroke

- **Most common type**
- Wet with prior sweating
- Altered Mental Status

Assess Symptom Severity

HEAT CRAMPS

Normal to elevated body temperature
Warm, moist skin
Weakness, Muscle cramping

HEAT EXHAUSTION

Elevated body temperature
Cool, moist skin
Weakness, Anxious, Tachypnea

HEAT STROKE

Fever, usually > 104°F (40°C)
Hot, dry skin
Hypotension, AMS / Coma

PO Fluids as tolerated
Monitor and Reassess

Age Appropriate Airway Protocol(s) AR 1 - 7 as indicated
Altered Mental Status Protocol UP 4 as indicated
Active cooling measures Target Temp < 102.5° F (39°C)
B 12 Lead ECG Procedure
IV or IO Access Protocol UP 6
P Cardiac Monitor
A Normal Saline Bolus 500 mL IV / IO Repeat to effect SBP > 90 Maximum 2 L PED: Bolus 20 mL/kg IV / IO Repeat to effect Age appropriate SBP ≥ 70 + 2 x Age Maximum 60 mL/kg
Age Appropriate Hypotension/ Shock Protocol AM 5/ PM 3 as indicated
Monitor and Reassess

Notify Destination or Contact Medical Control

Toxic-Environmental Protocol Section



Hyperthermia

Pearls

- **Recommended Exam: Mental Status, Skin, HEENT, Heart, Lungs, Neuro**
- **Extremes of age are more prone to heat emergencies (i.e. very young and very old).**
- **Temperature measurement:**
 - **Obtain and document patient temperature if able.**
 - **Many thermometers and routes of measurement are available.**
 - **Order of preference for route of measurement: Rectal > oral > temporal > axillary.**
- Heat illness is predisposed by use of: tricyclic antidepressants, phenothiazines, anticholinergic medications, and alcohol.
- Cocaine, Amphetamines, and Salicylates may elevate body temperatures.
- Intense shivering may occur as patient is cooled.
- **Heat Cramps:**
 - Consists of benign muscle cramping secondary to dehydration and is not associated with an elevated temperature.
- **Heat Exhaustion:**
 - Consists of dehydration, salt depletion, dizziness, fever, mental status changes, headache, cramping, nausea and vomiting.
 - Vital signs usually consist of tachycardia, hypotension, and an elevated temperature.
- **Heat Stroke:**
 - Consists of dehydration, tachycardia, hypotension, temperature $\geq 104^{\circ}\text{F}$ (40°C), and an altered mental status.
 - Sweating generally disappears as body temperature rises above 104°F (40°C).
 - The young and elderly are more prone to be dry with no sweating.
 - **Exertional Heat Stroke:**
 - **In exertional heat stroke (athletes, hard labor), the patient may have sweated profusely and be wet on exam.**
 - **Rapid cooling takes precedence over transport as early cooling decreases morbidity and mortality.**
 - **If available, immerse in an ice water bath for 20 minutes. Monitor rectal temperature and remove patient when temperature reaches 102.5°F (39°C). Your goal is to decrease rectal temperature below 104°F (40°C) with target of 102.5°F (39°C) within 15 minutes. Stirring the water aids in cooling.**
 - **Nearly 66% of all exertional heat strokes occur in high school athletes during the month of August.**
 - **In NC, it is mandatory that all high school field houses have a dunk tank and available ice and water.**
 - **Other methods include cold wet towels below and above the body or spraying cold water over body continuously.**
- **Neuroleptic Malignant Syndrome (NMS):**
 - Neuroleptic Malignant Syndrome is a hyperthermic emergency which is not related to heat exposure.
 - It occurs after taking neuroleptic antipsychotic medications.
 - This is a rare but often lethal syndrome characterized by muscular rigidity, AMS, tachycardia and hyperthermia.
 - **Drugs Associated with Neuroleptic Malignant Syndrome:**
 - Prochlorperazine (Compazine), promethazine (Phenergan), clozapine (Clozaril), risperidone (Risperdal)
 - metoclopramide (Reglan), amoxapine (Ascendin), and lithium.
 - **Management of NMS:**
 - Supportive care with attention to hypotension and volume depletion.
 - Use benzodiazepines such as diazepam or midazolam for seizures and/ or muscular rigidity.



Hypothermia/ Frostbite

History

- Age, very young and old
- Exposure to decreased temperatures but may occur in normal temperatures
- Past medical history / Medications
- Drug use: Alcohol, barbituates
- Infections/ Sepsis
- Length of exposure/ Wetness/ Wind chill

Signs and Symptoms

- Altered mental status/ coma
- Cold, clammy
- Shivering
- Extremity pain or sensory abnormality
- Bradycardia
- Hypotension or shock


Differential

- Sepsis
- Environmental exposure
- Hypothyroidism
- Hypoglycemia
- CNS dysfunction
 - Stroke
 - Head injury
 - Spinal cord injury

Temperature Measurement Procedure **if available**

Temperature Measurement should NOT delay treatment of hypothermia

Remove wet clothing Dry/ Warm Patient
Passive warming measures
Blood Glucose Analysis Procedure
Age Appropriate Diabetic Protocol AM 2/ PM 2 as indicated



Toxic-Environmental Protocol Section



Hypothermia/ Frostbite

Pearls

- **Recommended Exam: Mental Status, Heart, Lungs, Abdomen, Extremities, Neuro**
- **NO PATIENT IS DEAD UNTIL WARM AND DEAD (Body temperature $\geq 93.2^\circ$ F, 32° C.)**
- **Temperature measurement:**
 - Obtain and document patient temperature if able.
 - Many thermometers and routes of measurement are available.
 - Order of preference for route of measurement: Rectal > oral > temporal > axillary.
 - Many thermometers do not register temperature below 93.2° F.
- **Hypothermia categories:**
 - Mild $90 - 95^\circ$ F ($32 - 35^\circ$ C)
 - Moderate $82 - 90^\circ$ F ($28 - 32^\circ$ C)
 - Severe $< 82^\circ$ F ($< 28^\circ$ C)
- **Mechanisms of hypothermia:**
 - Radiation: Heat loss to surrounding objects via infrared energy (60% of most heat loss.)
 - Convection: Direct transfer of heat to the surrounding air.
 - Conduction: Direct transfer of heat to direct contact with cooler objects (important in submersion.)
 - Evaporation: Vaporization of water from sweat or other body water losses.
- Contributing factors of hypothermia: Extremes of age, malnutrition, alcohol or other drug use.
- If the temperature is unable to be measured, treat the patient based on the suspected temperature.
- **CPR:**
 - Severe hypothermia may cause cardiac instability and rough handling of the patient theoretically can cause ventricular fibrillation. This has not been demonstrated or confirmed by current evidence. Intubation and CPR techniques should not be with-held due to this concern.
 - Intubation can cause ventricular fibrillation, so it should be done gently by the most experienced provider(s).
 - Below 86° F (30° C) antiarrhythmics may not work and if given, should be given at increased time intervals. Contact medical control for direction. Epinephrine can be administered.
 - Below 86° F (30° C) pacing should not utilized.
 - Consider withholding CPR if patient has organized rhythm or has other signs of life. Contact Medical Control.
 - If the patient is below 86° F (30° C) then defibrillate 1 time if defibrillation is required. Deferring further attempts until more warming occurs is controversial. Contact medical control for direction.
 - Hypothermia may produce severe bradycardia so take at least 60 seconds to palpate a pulse.
- **Active Warming:**
 - Remove from cold environment and into warm environment protected from wind and wet conditions.
 - Remove wet clothing and provide warm blankets/ warming blankets.
 - Hot packs can be activated and placed in the armpit and groin area if available. Care should be taken not to place the packs directly against the patient's skin.



Marine Envenomation/ Injury

History

- Type of bite/ sting
- Identification of organism
- Previous reaction to marine organism
- Immunocompromised
- Household pet

Signs and Symptoms

- Intense localized pain
- Increased oral secretions
- Nausea/ vomiting
- Abdominal cramping
- Allergic reaction / anaphylaxis


Differential

- Jellyfish sting
- Sea Urchin sting
- Sting ray barb
- Coral sting
- Swimmers itch
- Cone Shell sting
- Fish bite
- Lion Fish sting

Call for help/ additional resources
Stage until scene safe

	General Wound Care Procedure
	IV or IO Access Protocol UP 6 if indicated
P	Cardiac Monitor if indicated
	Drowning Protocol TE 3 if indicated
	Age Appropriate Allergy/ Anaphylaxis Protocol AM 1/ PM 1 if indicated
	Age Appropriate Hypotension/ Shock Protocol AM 5/ PM 3 if indicated
	Pain Control Protocol UP 11 if indicated

Contact Carolinas Poison Control
1-800-222-1222
Or
Agency Specific Number



Toxic-Environmental Protocol Section



Marine Envenomation/ Injury

Pearls

- **Ensure your safety: Avoid the organism or fragments of the organism as they may impart further sting or injury.**
- **Priority is removal of the patient from the water to prevent drowning.**
- **Coral:**
 - Coral is covered by various living organisms which are easily dislodged from the structure.
 - Victim may swim into coral causing small cuts and abrasions and the coral may enter into cuts, causing little if any symptoms initially, but later causing inflammation, pain and/ or infection.
 - The next 24 – 48 hours may reveal an inflammatory reaction with swelling, redness, itching, tenderness, and ulceration.
 - Treatment is flushing with large amounts of fresh water or soapy water then repeating.
- **Jelly Fish/ Anemone/ Man-O-War:**
 - Wash the area with fresh seawater to remove tentacles and nematocysts.
 - Do not apply fresh water or ice as this will cause nematocysts firing as well.
 - Recent evidence does not demonstrate a clear choice of any solution that neutralizes nematocysts.
 - Vinegar (immersion for 30 seconds), 50:50 mixture of Baking Soda and Seawater, and even meat tenderizer may have similar effects.
 - Immersion in warm water for 20 minutes, 110 – 114°F (43 - 46°C), is effective in pain control.
 - Shaving cream may be useful in removing the tentacles and nematocysts with a sharp edge (card).
 - Stimulation of the nematocysts by pressure or rubbing cause the nematocyst to fire even if detached from the jellyfish.
 - Lift away tentacles as scrapping or rubbing will cause nematocysts firing.
 - Typically symptoms are immediate stinging sensation on contact, intensity increases over 10 minutes.
 - Redness and itching usually occur.
 - Papules, vesicles and pustules may be noted and ulcers may form on the skin.
 - Increased oral secretions and gastrointestinal cramping, nausea, pain, or vomiting may occur.
 - Muscle spasm, respiratory, and cardiovascular collapse may follow.
- **Lionfish:**
 - In North Carolina this would typically occur in a residence/ business as lionfish are often kept as pets in saltwater aquariums.
 - Remove any obvious protruding spines and irrigate area with copious amounts of saline.
 - The venom is heat labile so immersion in hot water, 110 – 114°F for 30 to 90 minutes is the treatment of choice but do not delay transport if indicated.
- **Stingrays:**
 - Typical injury is swimmer stepping on ray and muscular tail drives 1 – 4 barbs into victim.
 - Venom released when barb is broken.
 - Typical symptoms are immediate pain which increases over 1 – 2 hours.
 - Bleeding may be profuse due to deep puncture wound.
 - Nausea, vomiting, diarrhea, muscle cramping, and increased urination and salivation may occur.
 - Seizures, hypotension, and respiratory or cardiovascular collapse may occur.
 - Irrigate wound with saline. Extract the spine or barb unless in the abdomen or thorax, Contact Medical Control for advice.
 - Immersion in hot water, if available, for 30 to 90 minutes but do not delay transport.
- Patients can suffer cardiovascular collapse from both the venom and/ or anaphylaxis even in seemingly minor envenomation.
- Sea creature stings and bites impart moderate to severe pain.
- Arrest the envenomation by inactivation of the venom as appropriate.
- Ensure good wound care, immobilization and pain control.



Overdose/ Toxic Ingestion

History


- Ingestion or suspected ingestion of a potentially toxic substance
- Substance ingested, route, quantity
- Time of ingestion
- Reason (suicidal, accidental, criminal)
- Available medications in home
- Past medical history, medications

Signs and Symptoms

- Mental status changes
- Hypotension / hypertension
- Decreased respiratory rate
- Tachycardia, dysrhythmias
- Seizures
- S.L.U.D.G.E.
- D.U.M.B.B.E.L.S

Differential


- Tricyclic antidepressants (TCAs)
- Acetaminophen (Tylenol)
- Aspirin
- Depressants
- Stimulants
- Anticholinergic
- Cardiac medications
- Solvents, Alcohols, Cleaning agents
- Insecticides (organophosphates)



Toxin-Environmental Protocol Section



Overdose/ Toxic Ingestion





Overdose/ Toxic Ingestion

The goal of Narcan is to restore respirations;
NOT to restore consciousness.

Pearls

- **Recommended Exam: Mental Status, Skin, HEENT, Heart, Lungs, Abdomen, Extremities, Neuro**
 - **Opioids and opiates may require higher doses of Naloxone to improve respiration, in certain circumstances up to 10 mg.**
 - **Time of Ingestion:**
 1. **Most important aspect is the TIME OF INGESTION, the substance(s), amount ingested, and any co-ingestants.**
 2. **Every effort should be made to elicit this information before leaving the scene.**
 - **Charcoal Administration:**

The American Academy of Clinical Toxicology DOES NOT recommend the routine use of charcoal in poisonings.

 1. Consider Charcoal within the FIRST HOUR after ingestion. If a potentially life threatening substance is ingested or extended release agent(s) are involved and \geq one hour from ingestion, Contact Medical Control or NC Poison Control Center for direction.
 2. If NG would be necessary to administer Charcoal, then DO NOT administer unless known to be adsorbed, airway secured by intubation, and ingestion is less than ONE HOUR confirmed and potentially lethal.
 3. Charcoal in general, should only be given to a patient who is alert and awake such that they can self-administer the medication.
 - Do not rely on patient history of ingestion, especially in suicide attempts. Make sure patient is still not carrying hiding other medications or has any weapons.
 - **Pediatric:**

Age specific blood pressure 0 – 28 days > 60 mmHg, 1 month - 1 year > 70 mmHg, 1 - 10 years > 70 + (2 x age)mmHg and > 10 years > 90 mmHg.

Pediatric IV Fluid maintenance rate:

 - 4 mL for the first 10 kg of weight +
 - 2 mL for the second 10 kg of weight +
 - 1 mL for every additional kg in weight after 20 kg
- | Example: 34 kg pediatric | |
|--------------------------|-----------------------|
| First 10 kg: | 4 mL/kg/hr = 40 mL/hr |
| Second 10 kg: | 2 mL/kg/hr = 20 mL/hr |
| Final 14 Kg: | 1 mL/kg/hr = 14 mL/hr |
| Total: 74 mL/hr rate | |
- **Bring bottles, contents, emesis to ED.**
 - **S.L.U.D.G.E:** Salivation, Lacrimation, Urination, Defecation, GI distress, Emesis.
 - **D.U.M.B.B.E.L.S:** Diarrhea, Urination, Miosis, Bradycardia, Bronchorrhea, Emesis, Lacrimation, Salivation.
 - **Tricyclic:** 4 major areas of toxicity: seizures, dysrhythmias, hypotension, decreased mental status or coma; rapid progression from alert mental status to death.
 - **Acetaminophen:** initially normal or nausea/ vomiting. If not detected and treated, causes irreversible liver failure.
 - **Aspirin:** Early signs consist of abdominal pain and vomiting. Tachypnea and altered mental status may occur later. Renal dysfunction, liver failure, and or cerebral edema among other things can take place later.
 - **Depressants:** decreased HR, decreased BP, decreased temperature, decreased respirations, non-specific pupils.
 - **Stimulants:** increased HR, increased BP, increased temperature, dilated pupils, seizures.
 - **Anticholinergic:** increased HR, increased temperature, dilated pupils, mental status changes.
 - **Cardiac Medications:** dysrhythmias and mental status changes.
 - **Solvents:** nausea, coughing, vomiting, and mental status changes.
 - **Insecticides:** increased or decreased HR, increased secretions, nausea, vomiting, diarrhea, pinpoint pupils.
 - **Nerve Agent Antidote kits** contain 2 mg of Atropine and 600 mg of pralidoxime in an autoinjector for self administration or patient care. These kits may be available as part of the domestic preparedness for Weapons of Mass Destruction.
 - **EMR and EMT may administer naloxone by IN / IM route only and may administer from EMS supply. Agency medical director may require Contact of Medical Control prior to administration and may restrict locally.**
 - **When appropriate contact the North Carolina Poison Control Center for guidance, reference Policy 18.**
 - Consider restraints if necessary for patient's and/or personnel's protection per the Restraint Procedure.



WMD-Nerve Agent Protocol

History

- Exposure to chemical, biologic, radiologic, or nuclear hazard
- Potential exposure to unknown substance/hazard

Signs and Symptoms

- **S**alivation
- **L**acrimation
- **U**rination; increased, loss of control
- **D**efecation / Diarrhea
- **G**I Upset; Abdominal pain / cramping
- **E**mesis
- **M**uscle Twitching
- Seizure Activity
- Respiratory Arrest

Differential

- Nerve agent exposure (e.g., VX, Sarin, Soman, etc.)
- Organophosphate exposure (pesticide)
- Vesicant exposure (e.g., Mustard Gas, etc.)
- Respiratory Irritant Exposure (e.g., Hydrogen Sulfide, Ammonia, Chlorine, etc.)

Call for help/ additional resources
Stage until scene safe

Obtain history of exposure
Observe for specific toxidromes
Initiate triage and/or decontamination as indicated.

Contact
Carolinas Poison Control
1-800-222-1222
Or
Agency Specific Number

Symptom Severity

Asymptomatic

Monitor and Reassess
Every 15 minutes for symptoms
Initiate Treatment per Appropriate Arm

Minor Symptoms:
Respiratory Distress + SLUDGEM

IV or IO Access Protocol UP 6

Nerve Agent Kit IM
2 Doses Rapidly
if available

Major Symptoms:
Altered Mental Status, Seizures,
Respiratory Distress, Respiratory Arrest

IV or IO Access Protocol UP 6

Nerve Agent Kit IM
3 Doses Rapidly
if available

Atropine 2 mg IV / IO / IM
Pediatric: See Pearls
IV / IO / IM
Repeat every 3 to 5 minutes until symptoms resolve

Pralidoxime (2PAM)
600 mg IV / IO / IM
Pediatric: 15 – 25 mg / kg
IV / IO / IM
Over 30 minutes

Seizure Protocol UP 13

CDC/ ASPR
CHEMPACK Program

NC -57 EMS containers
-43 locations

Almost all citizens within 50 miles of CHEMPACK
See Page 2 and Pearls

Multiple Patients

YES

NO

Consider
Activation and deployment of CHEMPACK

CHEMPACK ACTIVATION:
(insert local number)

Healthcare Coalition Activation
(insert local number)

Notify Destination or
Contact Medical Control




WMD-Nerve Agent Protocol

Pearls

- **Recommended Exam: Mental Status, Skin, HEENT, Heart, Lungs, Gastrointestinal, Neuro**
- **Follow local HAZMAT protocols for decontamination and use of personal protective equipment.**
- **Adult/ Pediatric Atropine Dosing Guides:**
 - Confirmed attack: Begin with 1 Nerve Agent Kit for patients less than 7 years of age, 2 Nerve Agent Kits from 8 to 14 years of age, and 3 Nerve Agent Kits for patients 15 years of age and over.
 - If Triage/ MCI issues exhaust supply of Nerve Agent Kits, use pediatric atropines (if available).
 - Usual pediatric doses: 0.5 mg ≤ 40 pounds (18 kg), 1 mg dose if patient weighs between 40 to 90 pounds (18 to 40 kg), and 2 mg dose ≥ 90 pounds (≥ 40 kg).
- Each Nerve Agent Kit contains 600 mg of Pralidoxime (2-PAM) and 2 mg of Atropine.
- **Seizure Activity: Any benzodiazepine by any route is acceptable.**
- For patients with major symptoms, there is no limit for atropine dosing.
- Carefully evaluate patients to ensure they do not have exposure to other agent(s) (e.g., narcotics, vesicants, etc.)
- The main symptom that the atropine addresses is excessive secretions, so atropine should be given until secretions improve/ dry.
- EMS personnel, public safety officers and EMR/ EMT may carry, self-administer, or administer atropine/ pralidoxime to others by protocol. Agency medical director may require Contact of Medical Control prior to administration.
- **CHEMPACK Program:**
 - For multiple patients, call for **CHEMPACK** deployment per local emergency management and healthcare coalition plans.
 - 1 EMS CHEMPACK supports 454 patients.**
 - Medication in CHEMPACK may be used regardless of expiration date.

EMS Type CHEMPACK Container 454 Person Treatment Capacity			
Product	Cases	Units per case	Total Units
Mark 1 Auto-injector	5	240	1,200
-OR-			
ATNAA Auto-injector	6	200	1,200
-OR-			
Atropen 2mg Auto-injector	9	135	1,224
Pralidoxime 300mg Auto-injector	5	240	1,200
-AND-			
Diazepam 10mg Auto-injector	2	300	600
Seizalam (Midazolam) 5mg/ml vial 10ml	1	100	100
Atropen 0.5mg Auto-injector	1	225	225
Atropen 1mg Auto-injector	1	225	225
Atropine Sulfate 0.4mg/ml vial 20ml	1	100	100
Pralidoxime 1gm inj. 20ml	1	275	275
Sterile Water 20ml vials	1	150	150





Suspected Viral Hemorrhagic Fever Ebola

EMS Dispatch Center

- Use Emerging Infectious Disease (EID) Surveillance Tool with the following chief complaints:
Typical Flu-Like Symptoms
and/or
Unexpected Bleeding
(not trauma or isolated nose bleed related)
- Use EID Card (or equivalent) with the following protocols (or equivalent)
EMD 6 Breathing Problem
EMD 10 Chest Pain
EMD 18 Headache
EMD 21 Hemorrhage (medical)
EMD 26 Sick Person
- Ask the following:
In the past 21 days have you been to Africa or been exposed to someone who has?
If YES:
Do you have a fever?

Evolving Protocol:

Protocol subject to change at any time dependent on changing outbreak locations.

Monitor for protocol updates.

Viral Hemorrhagic Fevers:

Ebola is one of many.

DO NOT DISPATCH FIRST RESPONDERS

Dispatch EMS Unit only
Discretely notify EMS Supervisor or command staff

NO

EMS

Do not rely solely on EMD personnel to identify a potential viral hemorrhagic fever patient – constrained by time and caller information

Obtain a travel history / exposure history and assess for clinical signs and symptoms

EMS Immediate Concern

- Traveler from area with known VHF (Ebola) with or without symptoms
- Traveler from a Country, with active Ebola outbreak, within past 21 days

AND

Fever, Headache	Joint and Muscle aches	Weakness, Fatigue
Vomiting and/or Diarrhea	Abdominal Pain	Anorexia
Bleeding		

NO

Exit to Appropriate Protocol(s)

YES

EMS
Personal Protective Equipment

Refer to page 2
Place surgical mask on patient
Use Non-rebreather mask if Oxygen Needed
Donning and Doffing Guidelines

NO Routine
Aerosol Generating Procedures

Avoid aerosol generating procedures unless medically necessary
NIPPV / Nebulizer therapy / Intubation / BIAD / Suctioning

No Routine
IV or IO Lines

Avoid routine IV or IO access unless medically necessary
If IV / IO necessary:
Stop vehicle to lessen exposure risk

EMS Personnel / Equipment /
Transport Unit Requires
Decontamination

Refer to Page 3



**Notify Destination as soon and as discretely as possible
DO NOT ENTER facility with patient until instructed
Follow entry directions from hospital staff**



Special Circumstances Section

Suspected Viral Hemorrhagic Fever Ebola

PARTICULAR ATTENTION MUST BE PAID TO PROTECTING MUCOUS MEMBRANES OF THE EYES, NOSE, and MOUTH FROM SPLASHES OF INFECTIOUS MATERIAL OR SELF INOCULATION FROM SOILED PPE / GLOVES.

THERE SHOULD BE NO EXPOSED SKIN

DONNING PPE: **BEFORE** you enter the patient area.

Recommended PPE

PAPR: A PAPR with a full face shield, helmet, or headpiece. Any reusable helmet or headpiece must be covered with a single-use (disposable) hood that extends to the shoulders and fully covers the neck and is compatible with the selected PAPR.

N95 Respirator: Single-use (disposable) N95 respirator in combination with single-use (disposable) surgical hood extending to shoulders and single-use (disposable) full face shield. If N95 respirators are used instead of PAPRs, careful observation is required to ensure healthcare workers are not inadvertently touching their faces under the face shield during patient care.

Single-use (disposable) fluid-resistant or impermeable gown that extends to at least mid-calf or coverall without integrated hood. Coveralls with or without integrated socks are acceptable.

Single-use (disposable) nitrile examination gloves with extended cuffs. Two pairs of gloves should be worn. At a minimum, outer gloves should have extended cuffs.

Single-use (disposable), fluid-resistant or impermeable boot covers that extend to at least mid-calf or single-use (disposable) shoe covers. Boot and shoe covers should allow for ease of movement and not present a slip hazard to the worker.

Single-use (disposable) fluid-resistant or impermeable shoe covers are acceptable only if they will be used in combination with a coverall with integrated socks.

Single-use (disposable), fluid-resistant or impermeable apron that covers the torso to the level of the mid-calf should be used if Ebola patients have vomiting or diarrhea. An apron provides additional protection against exposure of the front of the body to body fluids or excrement. If a PAPR will be worn, consider selecting an apron that ties behind the neck to facilitate easier removal during the doffing procedure

DOFFING PPE: OUTSIDE OF PPE IS CONTAMINATED! DO NOT TOUCH

1) PPE must be carefully removed without contaminating one's eyes, mucous membranes, or clothing with potentially infectious materials.

Use great care while doffing your PPE so as not to contaminate yourself (e.g. Do not remove your N-95 facemask or eye protection BEFORE you remove your gown). There should be a dedicated monitor to observe donning and doffing of PPE. It is very easy for personnel to contaminate themselves when doffing. A dedicated monitor should observe doffing to insure it is done correctly. Follow CDC guidance on doffing.

2) PPE must be double bagged and placed into a regulated medical waste container and disposed of in an appropriate location.

3) Appropriate PPE must be worn while decontaminating / disinfecting EMS equipment or unit.

3) Re-useable PPE should be cleaned and disinfected according to the manufacturer's reprocessing instructions.

Hand Hygiene should be performed by washing with soap and water with hand friction for a minimum of 20 seconds.

Alcohol-based hand rubs may be used if soap and water are not available.

EVEN IF AN ALCOHOL-BASED HAND RUB IS USED, WASH HANDS WITH SOAP AND WATER AS SOON AS

FEASIBLE.

THE USE OF GLOVES IS NOT A SUBSTITUTE FOR HAND WASHING WITH SOAP & WATER


For any provider exposure or contamination contact occupational health.

If the patient is being transported via stretcher then a disposable sheet can be placed over them.

Pearls

- **Transmission to another individual is the greatest after a patient develops fever. Once there is fever, the viral load in the bodily fluids appears to be very high and thus a heightened level of PPE is required.**
- **Patient contact precautions are the most important consideration.**
- **Incubation period 2-21 days**
- **Ebola must be taken seriously; however using your training, protocols, procedures and proper Personal Protective Equipment (PPE), patients can be cared for safely.**
- When an infection does occur in humans, the virus can be spread in several ways to others. The virus is spread through direct contact (through broken skin or mucous membranes) with a sick person's blood or body fluids (urine, saliva, feces, vomit, and semen) objects (such as needles) that have been contaminated with infected body fluids.
- Limit the use of needles and other sharps as much as possible. All needles and sharps should be handled with extreme care and disposed in puncture-proof, sealed containers. Safety devices must be employed immediately after use.
- **Ebola Information: For a complete review of Ebola go to:**
<http://www.cdc.gov/vhf/ebola/index.html>
<https://www.cdc.gov/vhf/ebola/clinicians/emergency-services/ems-systems.html>

Suspected Viral Hemorrhagic Fever Ebola



- 1) EMS personnel performing decontamination / disinfection should wear recommended PPE
When performing Decontamination EMS Personnel MUST wear appropriate PPE, which includes:
 - Gloves (Double glove)
 - Fluid resistant (impervious) Tyvek Like Full length (Coveralls)
 - Eye protection (Goggles)
 - N-95 face mask
 - Fluid resistant (impervious)-Head covers
 - Fluid resistant (impervious)-Shoe / Boot covers
- 2) Face protection (N-95 facemask with goggles) should be worn since tasks such as liquid waste disposal can generate splashes.
- 3) Patient-care surfaces (including stretchers, railings, medical equipment control panels, and adjacent flooring, walls and work surfaces) are likely to become contaminated and should be decontaminated and disinfected after transport.
- 4) A blood spill or spill of other body fluid or substance (e.g., feces or vomit) should be managed through removal of bulk spill matter, cleaning the site, and then disinfecting the site. For large spills, a chemical disinfectant with sufficient potency is needed to overcome the tendency of proteins in blood and other body substances to neutralize the disinfectant's active ingredient. An EPA-registered hospital disinfectant with label claims for viruses that share some technical similarities to Ebola (such as, norovirus, rotavirus, adenovirus, poliovirus) and instructions for cleaning and decontaminating surfaces or objects soiled with blood or body fluids should be used according to those instructions.
(Alternatively, a 1:10 dilution of household bleach (final working concentration of 500 parts per million or 0.5% hypochlorite solution) that is prepared fresh daily (i.e., within 12 hours) can be used to treat the spill before covering with absorbent material and wiping up. After the bulk waste is wiped up, the surface should be disinfected as described in the section above).
- 5) Contaminated reusable patient care equipment should be placed in biohazard bags (double-bagged) and labeled for decontamination and disinfection.
- 6) Reusable equipment should be cleaned and disinfected according to manufacturer's instructions by appropriately trained personnel wearing correct PPE.
- 7) Avoid contamination of reusable porous surfaces that cannot be made single use. Use only a mattress and pillow with plastic or other covering that fluids cannot get through.
- 8) To reduce exposure, all potentially contaminated textiles (cloth products) should be discarded. This includes non-fluid-impermeable pillows or mattresses. They should be considered regulated medical waste and placed in biohazard red bags. They must be double-bagged prior to being placed into regulated medical waste containers.

Pearls

- **Ebola Information:** For a complete review of Ebola EMS Vehicle Disinfection go to:
<https://www.cdc.gov/vhf/ebola/clinicians/emergency-services/ems-systems.html>

Suspected Viral Hemorrhagic Fever Ebola

Decedent Known or suspected carrier of HVF / Ebola Requires Transportation

Only personnel trained in handling infected human remains, and wearing full PPE, should touch, or move any Ebola-infected remains.
Handling human remains should be kept to a minimum.

Donning / Doffing PPE

PPE should be in place BEFORE contact with the body

- 1) Prior to contact with body, postmortem care personnel must wear PPE consisting of: surgical scrub suit, surgical cap, impervious Tyvex-Coveralls, eye protection (e.g., face shield, goggles), facemask, shoe covers, and double surgical gloves.
- 2) Additional PPE (leg coverings,) might be required in certain situations (e.g., copious amounts of blood, vomit, feces, or other body fluids that can contaminate the environment).

PPE should be removed immediately after and discarded as regulated medical waste.

- 1) Use caution when removing PPE as to avoid contaminating the wearer.
- 2) Hand hygiene (washing your hands thoroughly with soap and water or an alcohol based hand rub) should be performed immediately following the removal of PPE. If hands are visibly soiled, use soap and water.

Preparation of Body Prior to Transport

- 1) At the site of death, the body should be wrapped in a plastic shroud. Wrapping of the body should be done in a way that prevents contamination of the outside of the shroud.
- 2) Change your gown or gloves if they become heavily contaminated with blood or body fluids.
- 3) Leave any intravenous lines or endotracheal tubes that may be present in place.
- 4) Avoid washing or cleaning the body.
- 5) After wrapping, the body should be immediately placed in a leak-proof plastic bag not less than 150 μ m thick and zippered closed. The bagged body should then be placed in another leak-proof plastic bag not less than 150 μ m thick and zippered closed before being transported to the morgue.

Surface Decontamination

- 1) Prior to transport to the morgue, perform surface decontamination of the corpse-containing body bags by removing visible soil on outer bag surfaces with EPA-registered disinfectants which can kill a wide range of viruses.
- 2) Follow the product's label instructions. Once the visible soil has been removed, reapply the disinfectant to the entire bag surface and allow to air dry.
- 3) Following the removal of the body, the patient room should be cleaned and disinfected.
- 4) Reusable equipment should be cleaned and disinfected according to standard procedures.

Transportation of VHV / Ebola Remains

PPE is required for individuals driving or riding in a vehicle carrying human remains. DO NOT handle the remains of a suspected / confirmed case of Ebola. The remains must be safely contained in a body bag where the outer surface of the body bag has been disinfected prior to the transport.

Pearls

- **Ebola Information:** For a complete review of Handling Remains of Ebola Infected Patients go to: <http://www.cdc.gov/vhf/ebola/hcp/guidance-safe-handling-human-remains-ebola-patients-us-hospitals-mortuaries.html>

SC 1

Any local EMS System changes to this document must follow the NC OEMS Protocol Change Policy and be approved by OEMS

High Consequence Pathogens (Respiratory Diseases, SARS, MERS-CoV, COVID-19)

EMD Dispatch Center Screening

1. All calls requiring response from EMS system:

**Ask: Do you have FEVER AND/OR RESPIRATORY SYMPTOMS?
(cough, breathing difficulty, or other respiratory symptoms?)**

EMD Systems:

- PDS – Card 36 Pandemic Flu
- APCO – COVID-19 Pandemic Vital Points Card
- PowerPhone – Pandemic Influenza Card

Evolving Protocol:

Protocol subject to change at any time dependent on changing outbreak locations.

Monitor for protocol updates.

EMD Screen Positive

Notify

All Responding Agencies:

- Positive screening (agency specific code)
- First Responder Response:
Follow local system guidance

EMD Screen Negative

First Responders and EMS Screening

Do not rely solely on EMD personnel to identify a potential exposure patient:

- EMD may be constrained by time and caller information.
- First arriving provider (FR or EMS):**
If call nature allows, send 1 provider only into scene to complete a quick screen. Stand at a distance of ≥ 6 feet and perform screening question. Patients with Fever and/or Cough (or other respiratory symptoms are at risk of Influenza and/or COVID-19).
Chills, muscle aches, sore throat, or sudden loss of taste or smell.
If patient screens positive:
Place facemask or covering over patient's mouth and nose and provider dons appropriate PPE based on clinical situation.
- First Responders should stage and limit number of providers entering scene only necessary for care to limit potential exposures and use of PPE.
- Request additional resources as needed. See Page 4.

Negative
FR or EMS Screening

Exit to
Appropriate
Protocol(s)

PPE Supply Chain Disruptions:

- Prioritize respirators (N95 or equivalent) to aerosol-generating procedures until supply chain restored.
- Prioritize gowns to aerosol-generating procedures.
- It is reasonable for providers to wear a facemask during their duty-shift and change only when soiled or damaged. Adjust use based on supply chain.

Positive
FR or EMS Screening

EMS PPE

EMS
General Treatment
Considerations

Exit to
Appropriate Protocol(s)

Patient:

- Use non-rebreather mask if oxygen needed
- If unable to tolerate mask, have patient cover mouth and nose when coughing

Providers utilize:

- Follow PPE precautions listed below:**
- Exam gloves and eye protection
- Facemask minimum
- Aerosol generating procedure:**
- Respirator (N95, P100, PAPR, or equiv.)
- Goggles, gown (disposable gown, coveralls, or equivalent)
- Create negative pressure in care compartment (See Pearls)

Personnel in ambulance cab utilize:

- Facemask for driver and passenger

Aerosol generating procedures:

NIPPV / Nebulizer therapy / Intubation / BIAD / Suctioning / CPR

Use all PPE devices and strategies listed above

- Notify receiving facility of infection control requirements prior to arrival.**

High Consequence Pathogens

(Respiratory Diseases, SARS, MERS-CoV, COVID-19)

Pearls

- **First Responders: Because community spread is now present, every patient contact should be considered to have potential for infection with COVID-19. Limit number of FR when caring for patients to limit exposures and PPE use.**
- **Place facemask on any patient complaining of respiratory problems with or without a fever.**
- **Dispatch Screening:**
 - If caller interrogation results in positive screen first responders are assigned based on local agency direction.
 - This screening process will result in many False Positive screens in order to be very sensitive.
- **First Responder and EMS Screening:**
 - Limit distance initially to ≥ 6 feet and conduct a quick screening using the EMD specific question. If this results in a positive screen, immediately place a facemask on the source patient and all providers don appropriate PPE and limit provider number to that which necessary for patient care.
- **Close Contact and Duration Definition:**
 - Healthcare provider exposure is defined as being within 6 feet for ≥ 15 minutes with a patient with suspected illness.
 - Unprotected (no or incorrect PPE) with direct contact with body fluids, including respiratory generated body fluids.
- **Transport:**
 - Occupants in cab of vehicle all should wear facemasks. Riders should be discouraged in order to limit PPE use.
 - Limit number of providers in vehicle required to provide patient care in order to limit exposures.
 - Ensure use of correct PPE for crew and passengers when aerosol-producing procedures utilized.
- **Recommend facemask and gloves with every patient contact. It is reasonable to wear eye protection on every patient contact.**
- **Reasonable to wear simple/surgical mask during entire duty-shift when not able to maintain social distance of ≥ 6 feet among fellow providers when not engaged in patient care.**
- **Negative Pressure in care compartment:**
 - Door or window available to separate driver's and care compartment space:**
 - Close door/window between driver's and care compartment and operate rear exhaust fan on full.
 - No door or window available to separate driver's and care compartment space:**
 - Open outside air vent in driver's compartment and set rear exhaust fan to full.
 - Set vehicle ventilation system to non-recirculating to bring in maximum outside air.
 - Use recirculating HEPA ventilation system if equipped.
- **Airborne precautions:**
 - Standard PPE with fit-tested N95 mask (or PAPR respirator) and utilization of a gown or coveralls, change of gloves after every patient contact, and strict hand washing precautions. This level is utilized with Aspergillus, SARS/MERS/COVID-19, Tuberculosis, Measles (rubeola) Chickenpox (varicella-zoster), Smallpox, Influenza, disseminated herpes zoster, or Adenovirus/Rhinovirus.
- **Contact precautions:**
 - Standard PPE with utilization of a gown or coveralls, change of gloves after every patient contact, and strict hand washing precautions.
 - This level is utilized with GI complaints, blood or body fluids, C diff, scabies, wound and skin infections, MRSA.
 - Clostridium difficile (C diff) is not inactivated by alcohol-based cleaners and washing with soap and water is indicated.
- **Droplet precautions:**
 - Standard PPE plus a standard surgical mask for providers who accompany patients in the treatment compartment and a surgical mask or NRB O2 mask for the patient.
 - This level is utilized when Influenza, Meningitis, Mumps, Streptococcal pharyngitis, Pertussis, Adenovirus, Rhinovirus, and undiagnosed rashes.
- **All-hazards precautions:**
 - Standard PPE plus airborne precautions plus contact precautions.
 - This level is utilized during the initial phases of an outbreak when the etiology of the infection is unknown or when the causative agent is found to be highly contagious (e.g. SARS, MERS-CoV, COVID-19).
- **COVID-19 (Novel Coronavirus): For most current criteria to guide evaluations of patients under investigation:**
 - <https://www.cdc.gov/coronavirus/2019-ncov/symptoms-testing/symptoms.html>

High Consequence Pathogens

(Respiratory Diseases, SARS, MERS-CoV, COVID-19)

Decontamination Recommendations

EMS Personnel Requires Decontamination

Driver:

- Should wear full PPE as described when caring for patient.
- Remove all PPE, except respiratory (N95, PAPR, or equivalent) and perform hand hygiene prior to entering cab to prevent contamination of driver's compartment. **Cab occupants only need to wear facemasks if respirator not already used.**

Wash hands:

- Thoroughly after transferring patient care and/or cleaning ambulance

Maintain records:

- All prehospital providers exposed to patient at the scene and during ambulance transport (self-monitoring for symptoms for 14 days is recommended, even if wearing appropriate PPE).
This does not mean the providers can no longer work.
- List all prehospital provider names (students, observers, supervisors, first response etc.) in the Patient Care Report.

EMS Equipment / Transport Unit Requires Decontamination

Safely clean vehicles used for transport:

- Follow standard operating procedures for the containment and disposal of regulated medical waste.
- Follow standard operating procedures for containing and reprocessing used linen.

Wear appropriate PPE when:

- Removing soiled linen from the vehicle. Avoid shaking the linen.
- Clean and disinfect the vehicle in accordance with agency standard operating procedures.
- Personnel performing the cleaning should wear a disposable gown and gloves (a respirator should not be needed) during the clean-up process; the PPE should be discarded after use.
- All surfaces that may have come in contact with the patient or materials contaminated during patient care (e.g., stretcher, rails, control panels, floors, walls, work surfaces) should be thoroughly cleaned and disinfected using an **EPA-registered disinfectant** appropriate for SARS, MERS-CoV, or coronavirus in healthcare settings in accordance with manufacturer's recommendations. **Keep doors open to patient care compartment while cleaning to allow air exchanges.**

EMS Provider Exposure Risk and Monitoring Recommendations

Close Contact				Close Contact			
Less than 6 feet for ≥ 15 minutes				Less than 6 feet for ≥ 15 minutes			
Source patient NOT WEARING A MASK				Source patient WEARING A MASK			
PPE Utilized	Exposure Risk	Monitoring	Work Restrictions	PPE Utilized	Exposure Risk	Monitoring	Work Restrictions
NONE	HIGH	Self-monitor Supervision	If symptomatic: Fever and Respiratory symptoms (cough, difficulty breathing or other respiratory symptoms) THEN Exclude from work: • At least 72 hours after fever resolution with no use of fever reducing medications. AND • At least 10 days since symptom onset.	NONE	MEDIUM	Self-monitor Supervision	If symptomatic: Fever and Respiratory symptoms (cough, difficulty breathing or other respiratory symptoms) THEN Exclude from work: • At least 72 hours after fever resolution with no use of fever reducing medications. AND • At least 10 days since symptom onset.
No facemask N95 or PAPR	HIGH			No facemask N95 or PAPR	MEDIUM		
No Eye Protection	MEDIUM			No Eye Protection	LOW		
No Gown/ Coveralls or Gloves	LOW			No Gown/ Coveralls or Gloves	LOW		
All recommended PPE Except facemask instead of N95 or PAPR	LOW			All recommended PPE Except facemask instead of N95 or PAPR	LOW		

Placing a simple/surgical mask on the patient within 15 minutes of contact decreases exposure risk.

Return to Work Practice and Work Restrictions (if excluded from work OR exposure to suspected or known COVID-19 patient):

- Prior to duty shift, measure temperature and assess for illness symptoms either by provider, infection control officer, or occupational or public health.
- Self-monitoring with oversight by agency's infection control officer, occupation or public health department per agency policy.
- Wear mask at all times and restrict care of immunocompromised patients (Cancer, Transplant, Steroid use) until all symptoms have resolved or 14 days after onset of illness, whichever is longest.
- Social distance: Employee should maintain 6 feet of separation as work duties permit in the workspace.
- Remove from work if employee becomes symptomatic.

- <https://www.cdc.gov/coronavirus/2019-ncov/hcp/guidance-for-ems.html>
- <https://www.cdc.gov/coronavirus/2019-ncov/healthcare-facilities/hcp-return-work.html>
- <https://www.epa.gov/pesticide-registration/list-n-disinfectants-use-against-sars-cov-2-covid-19>

Special Circumstances Section

High Consequence Pathogens (Respiratory Diseases, SARS, MERS-CoV, COVID-19)

First Responder Guidance

COVID-19 Declared Pandemic with both State and Federal Emergencies Declared

- Many systems are heavily dependent on First Responder agencies to supplement critical prehospital medical care services.
- Community spread is now evident both in NC and in the US.
- Every patient, regardless of medical or injury complaint, is at risk of COVID-19 and all should undergo routine screening questions.
- While EMD is a first step, all providers must screen every patient contact and don appropriate PPE based on clinical situation and COVID-19 screening.
- The citizens we serve continue to have a variety of illness and injury unrelated to COVID-19.

Limiting PPE use:

First Responders should consider staging with all incidents and sending 1 provider (or more dependent on situation) into the scene to assess for fever and respiratory complaints.

Request staged resources as needed only to provide necessary medical care.

Where patients do not require immediate intervention, first responders may stay in contact with patient, but remain beyond 6 feet until EMS providers arrive to begin assessment and further care.

Consider calling patient on mobile phones to maintain contact and provide reassurance and explain current situation.

PPE Crisis or Alternative Strategies

N95 Respirators

- Use only for aerosol generating procedures (Nebulizer, NIPPV, Suctioning, BVM, BIAD, Intubation).
- Use facemasks in all other scenarios.
- Use respirators (N95 or equivalent) beyond the manufacturing expiration date when not soiled, ripped, torn, or otherwise damaged. Securing straps should also be in good repair and operational:
Visually inspect straps, nose bridge/foam, and mask in general.
Perform seal check: <https://www.youtube.com/watch?v=pGXiUyAoEd8>
- Models tested by CDC and are believed to function properly beyond expiration date:
3M: 1860, 1860s, 1870, 8210, 9010, 8000 Medline/Alpha Protech NON27501
Gerson 1730 Moldex: 1512, 2201
- Minimize providers caring for patient to the extent possible to conserve.
- Use Self-Contained Breathing Apparatus (SCBA) if needed.
- Re-use respiratory (N95 or equivalent) masks and place in paper bag between use. Do not touch inside of mask. Wash hands thoroughly before removing mask.
- When to discard a respirator (N95 or equivalent):
After using during an aerosol producing procedure.
Contamination with blood, body fluids or secretions, following close contact with known COVID-19 patient.


Gowns:

- Use only for aerosol generating procedures (Nebulizer, NIPPV, Suctioning, BVM, BIAD, Intubation).
- Use only for close patient contact, lifting, moving, or transferring where provider contacts patients body.
- May use removable and washable coveralls.

<https://www.cdc.gov/coronavirus/2019-ncov/hcp/respirators-strategy/index.html>



Hospice or Palliative Care Patient (Optional)



Request CP or MIH Medic To Scene
If available

Exit to
Appropriate Adult or
Pediatric Protocol(s)

Contact Hospice Service
and
Ensure Hospice Response
to scene
Local Hospice Agency(s)
Contact Number

Exit to
Appropriate Adult or
Pediatric Protocol(s)

Involve Hospice Service in
transport decision and/or make
aware of transport and destination
IV or IO Access Protocol UP 6
Parenteral Access
Existing Catheters Procedure PAS 7
If available
Port-A-Cath Access Procedure PAS 13
If available

P Administer Medications from **eKit**
Follow instructions in **eKit**
or
Equivalent Medications
from EMS Supply
See Pearls

P Administer Medications from **eKit**
Follow instructions in **eKit**
or
Equivalent Medications
from EMS Supply
See Pearls

Exit to
Appropriate Adult or Pediatric Protocol(s)
Consider
Treatment in place with Treat/ No transport
In consultation with Hospice Nurse or Provider

Special Circumstances Section



Hospice or Palliative Care Patient (Optional)

Acute Pain / Air Hunger:

Severity	Medication		
	Morphine (IV/IM/SQ)	Dilaudid (IV/IM/SQ)	Fentanyl (IV/IM/SQ)
Mild	2 mg	0.5 mg	25 mcg
Moderate	4 mg	1 mg	50 mcg
Severe	8 mg	2 mg	100 mcg
Titration	2 mg q 15 minutes IV	0.5 mg q 15 minutes IV	25 mcg q 15 minutes IV

Due to pain associated with IM injection, IM administration should only be used if alternative medications or routes of administration are not available. PICC lines may be accessed for use by EMS with sterile techniques. May access port-a-cath if appropriate equipment is available and provider is trained.

If using IM or SQ injections, delay repeat dosing by 30 minutes to prevent dose stacking.

Consider using moderate / severe dose in opiate tolerant patients:

Opiate tolerant patients have typical daily dose of narcotic is equivalent to ≥ 60 mg of oral Morphine per day (60 OME (Oral Morphine Equivalents)).

Examples of opiate dosages equivalent to 60 mg of oral Morphine:

- | | |
|--------------------------------|--------------------------|
| 40 mg/day of Oxycodone | 60 mg/day Hydrocodone |
| 25 mcg/hr Fentanyl Transdermal | 15 mg/day of Methadone |
| 200 mg/day of Tapentadol | 16 mg/day of Oxymorphone |
| Suboxone | |

Consider total use of multiple types of opiates. If in doubt about the patient's level of opiate tolerance, or amount of total daily opiate use, treat with a lower initial dose of opiate.

Anxiety / Agitation:

Severity	Medication			
	Ativan (IV/IM/SQ)	Versed (IV/IM/SQ)	Valium (IV/IM/SQ)	Haldol (IV/IM/SQ)
Mild / Moderate	0.5 mg	1 mg	2 mg	2 mg
Severe	1 mg	2 mg	5 mg	4 mg

May repeat dose in 15 minutes for IV administration, or 30 minutes for IM or SQ injections.

Nausea / Vomiting:

Zofran IV / IM	Phenergan IV / IM	Haldol IV / IM / SQ	Ativan IV / IM / SQ
4 mg	25 mg	2 mg	0.5 mg

Pearls

- **MOST form Section A and DNR forms are equivalent – if valid, Do Not Resuscitate.**
- **MOST form and DNR forms may be revoked by Health Care Power of Attorney or other appropriate surrogate decision-makers.**
- Palliative care is specialized care for patients with a chronic and/ or terminal illness which focuses on managing symptoms exacerbation and the stress of illness.
- Hospice care is specialized care (similar to palliative care) for patients within the last 6 months of life.
- Hospice patient may not have a DNR or MOST form completed and still be enrolled in Hospice care.
- **Emergency Kits (eKit):**
May be given to patient by Hospice to use at home for acute symptom exacerbation. Each eKit is individualized and will be different for each patient, but typically addresses pain, nausea/ vomiting, anxiety, and/ or secretions. (EMS is able to administer if within provider's scope of practice.)
- **Interaction on-scene with Hospice personnel:**
Hospice nurses are valuable resources in helping patients/ families make care/ transport decisions. EMS should discuss care/ transport decision with Hospice nurse. After medication administration, if no transport occurs, care may be transferred to Hospice nurse.



Mass Vaccination/Immunization Medication Distribution

History

- Follow local public health department criteria for specific immunization or medication administered.
- Patient receiving medication or vaccination must be without evidence of active infection.
- AEMT and Paramedic providers may participate
- EMT may participate when DHHS/NCMB allows special provision during local or state emergency.

Situation

- Local implementation of this protocol must be done as a component of the EMS system's local public health department community immunization or medication distribution program.
- May initiate protocol when a community has limited public health department resources or when local or state health emergency is declared.

Review immunization/vaccination or medication guide provided by the local public health department:

- | | |
|----------|---|
| B | <ul style="list-style-type: none"> • Patient selection criteria per local public health department (may vary) • Vaccine/immunization or medication indications • Vaccine/immunization or medication contraindications • Vaccine/immunization or medication distribution procedure |
| B | <ul style="list-style-type: none"> • EMT may provide vaccinations when DHHS/NCMB allows special provision during local or state emergency. |

Confirm patient eligibility for the vaccination or medication including:

- Age
- Medical history
- Contraindications
- Allergies

Eligibility confirmed?

YES

NO

Administer vaccination or medication:

- | | |
|----------|--|
| B | <ul style="list-style-type: none"> • Dose dependent on local public health department • Route dependent on local public health department (PO, IN, IM, IV, SQ) |
|----------|--|

Administer Over-the-Counter medication and/or vaccination (if applicable):

- | | |
|----------|---|
| B | <ul style="list-style-type: none"> • Undergo specific "just-in-time" training • Dose dependent on local public health department • Route dependent on local public health department (PO, IN, IM). SQ when specified by NCOEMS. • Complete required local public health department documentation • Provide post immunization or medication written instructions and monitoring |
|----------|---|

Do not administer:

- Refer to local public health department providers/officials for further care and instructions.

Allergic Reaction or Complications



- Exit to age appropriate Protocol(s)
- Notify appropriate local public health department provider/official

Pearls

- **Purpose:**
Provide protocol driven process for EMS providers to assist with public health immunization or medication distribution initiatives.
- **Documentation of the immunization or medication:**
Complete using local public health department approved record system.
Creation of an EMS patient care report is not required and is not required to submit to NCOEMS.
Must create a log of all patient contacts associated with the immunization or medication distribution program maintained by the EMS system.
If local public health department is maintaining a log of all patients, EMS may use the public health log and keep copies in the EMS system.
- **Injection site:**
Most common injection site for subcutaneous is tissue of an upper arm; follow procedure USP-4 otherwise.
Injection volume is limited to 1 - 2 mL per site unless specific guidance is given per local public health department.
Most common sites for intramuscular injections are upper arm, buttocks, and thighs, follow procedure USP-4.
Injection volume is limited to 1 mL in the upper arm, unless specific guidance is given per local public health department; follow procedure USP-4 otherwise.
Injection volume is limited to 2 mL (1 mL in pediatrics) in buttocks and thighs, unless specific guidance is given per local public health department; follow procedure USP-4 otherwise.



SARS CoV2 Monoclonal Antibody Administration

History

- FDA has issued an Emergency Use Authorization permitting the administration of REGEN-COV (casirivimab and imdevimab) for the treatment of mild to moderate COVID-19.
- Monoclonal antibodies are used to neutralize and prevent progression of the SARS CoV2 virus.

Situation

- Local implementation of this protocol must be done as a component of the EMS system's local public health department community immunization or medication distribution program.
- May initiate protocol when a community has limited public health department resources or when local or state health emergency is declared.

Review monoclonal antibody eligibility criteria:

- https://www.ems.gov/pdf/EMS_Template_Protocol_for_COVID-19_Monoclonal_Antibody_Administration_August_2021.pdf
- See page 2.
- Determine medication route for either intravenous or subcutaneous administration

Allergic Reaction or Complications



- Exit to age appropriate Protocol(s)
- Notify appropriate local public health department provider/official

Confirm patient eligibility for monoclonal antibody including:

- Age
- Medical history
- Contraindications
- Allergies

Eligibility confirmed?

NO

YES

Do not administer:

- Refer to local public health department providers/officials for further care and instructions.

Intravenous Administration:

Mix

Casirivimab 600 mg and Imdevimab 600 mg
In 100 mL NS

Infuse IV piggyback in a NS line at KVO
over 21 minutes (310 mL/hr)

Infuse through NS primed micron filter 0.20 or 0.22 size

Subcutaneous Administration:

Draw up

Casirivimab 600 mg (2.5 mL) in 2 separate syringes
and
Imdevimab 600 mg (2.5 mL) in 2 separate syringes
Administer 4 SQ injections in separate muscle areas

Avoid the waistline and 5 cm periumbilically

Injections: Subcutaneous and Intramuscular
Procedure USP - 4

Pearls

- **Purpose:**
Provide protocol driven process for EMS providers to assist with public health medication distribution initiatives.
- **Documentation of the medication:**
Creation of an EMS patient care report is required and is required to submit to NCOEMS.
Must create a log of all patient contacts associated with the medication distribution program maintained by the EMS system.
If local public health department is maintaining a log of all patients, EMS may use the public health log and keep copies in the EMS system.
- **Injection site:**
Most common injection site for subcutaneous is tissue of an upper arm; follow procedure USP-4 otherwise.
Injection volume is limited to 1 - 2 mL per site unless specific guidance is given per local public health department.
Most common sites for intramuscular injections are upper arm, buttocks, and thighs, follow procedure USP-4.
Injection volume is limited to 1 mL in the upper arm, unless specific guidance is given per local public health department; follow procedure USP-4 otherwise.
Injection volume is limited to 2 mL (1 mL in pediatrics) in buttocks and thighs, unless specific guidance is given per local public health department; follow procedure USP-4 otherwise.



SARS CoV2 Monoclonal Antibody Administration

Eligibility criteria:

- Age > 12 and weight \geq 40 kg.
- Not requiring hospitalization
- Not requiring oxygen therapy
- High risk for disease progression
 - Age \geq 65
 - Obesity
 - Pregnancy
 - Chronic kidney disease
 - Dementia
 - Diabetes
 - Immunocompromised or immunosuppressive treatments
 - Cardiovascular disease (MI, CVA, CHF, hypertension, hyperlipidemia, diabetes)
 - Chronic lung disease (COPD, asthma, interstitial lung disease)
 - Cancer
 - Sickle cell disease
 - Liver disease
 - Neurodevelopmental disorders, metabolic syndromes, or congenital abnormalities
 - Medical technology dependent, tracheostomy, gastrostomy, or NIPPV/ventilator
- Post-Exposure Prophylaxis (PEP)
 - Not fully vaccinated and immunocompromised or taking immunosuppressive medications
 - Only 1 of 2 doses and/or less than 2 weeks from 2d dose in 2 dose series or less than 2 weeks from 1st does in vaccine only requiring 1 dose.*
 - OR:** Individuals at high risk of exposure to a SARS-CoV-2 infected individual
(Nursing home or prison resident)
- High risk of death
 - Age \geq 80
 - Male sex
 - Black and South Asian descent



Scene Rehabilitation: General

Injury / Illness / Complaint should be treated using appropriate treatment protocol beyond need for oral or IV hydration.



- Initial Process**
1. Personnel logged into General Rehabilitation Section
 2. VS Assessed / Recorded (If HR > 110 then obtain Temp)
Carbon Monoxide monitoring if indicated
 3. Personnel assessed for signs / symptoms
 4. Remove PPE, Body Armor, Haz-Mat Suits, Turnout Gear, Other equipment as indicated

Significant Injury
Cardiac Complaint: Signs / Symptoms
Respiratory Complaint: Serious Signs / Symptoms
Respiratory Rate < 8 or > 40
Systolic Blood Pressure ≤ 80

Exit to Scene Rehabilitation Responder Protocol



NO

Heat or Cold stress

HEAT STRESS

Active Cooling Measures
Forearm immersion, cool shirts, cool mist fans etc.
Rest 10 – 20 Minutes

COLD STRESS

Active Warming Measures
Dry responder, place in warm area
Hot packs to axilla and / or groin
Rest 10 – 20 minutes

NO

Rehydration Techniques
12 – 32 oz Oral Fluid over 20 minutes
Oral Rehydration may occur along with Active Cooling Measures
Firefighters should consume 8 ounces of fluid between SCBA change-out

Rehydration Techniques
12 – 32 oz Oral Fluid over 20 minutes
Oral Rehydration may occur along with Active Warming Measures
Firefighters should consume 8 ounces of fluid between SCBA change-out

Reassess responder after 20 Minutes in General Rehabilitation Section
Reassess VS

Responder Cannot Wear Protective Gear
Extend Rehabilitation Time Until VS Improve

HR ≥ 110

Temp ≥ 100.6

NO

NO

Temp ≥ 100.6

HR ≥ 110

NO

NO

Extend Rehabilitation Time Until VS Improve

Discharge Responder from General Rehabilitation Section
Reports for Reassignment

VITAL SIGN CAVEATS

Blood Pressure:

Prone to inaccuracy on scenes. Must be interpreted in context.

Firefighters have elevated blood pressure due to physical exertion and is not typically pathologic.

Firefighters with Systolic BP ≥ 160 or Diastolic BP ≥ 100 may need extended rehabilitation. However this does not necessarily prevent them from returning to duty.

Temperature:

Firefighters may have increased temperature during rehabilitation.

Special Operations Section

Scene Rehabilitation: General

Pearls

- **This protocol is optional and given only as an example. Agencies may and are encouraged to develop their own.**
- **Rehabilitation officer has full authority in deciding when responders may return to duty and may adjust rest / rehabilitation time frames depending on existing conditions.**
- **Rehabilitation goals:**
 - Relief from climatic conditions.**
 - Rest, recovery, and hydration prior to incident, during, and following incident.**
 - Active and / or passive cooling or warming as needed for incident type and climate conditions.**
- **May be utilized with adult responders on fire, law enforcement, rescue, EMS and training scenes.**
- **Responders taking anti-histamines, blood pressure medication, diuretics or stimulants are at increased risk for cold and heat stress.**
- **General indications for rehabilitation:**
 - 20-minute rehabilitation following use of a second 30-minute SCBA, 45-minute SCBA or single 60-minute SCBA cylinder.
 - 20-minute rehabilitation following 40 minutes of intense work without SCBA.
- **General work-rest cycles:**
 - 10-minute self-rehabilitation following use of one 30-minute SCBA cylinder or performing 20 minutes of intense work without SCBA.
- **Serious signs / symptoms:**
 - Chest pain, dizziness, dyspnea, weakness, nausea, or headache.
 - Symptoms of heat stress (cramps) or cold stress.
 - Changes in gait, speech, or behavior.
 - Altered Mental Status.
 - Abnormal Vital Signs per agency SOP or Policy / Procedure.
- **Rehabilitation Section:**
 - Integral function within the Incident Management System.
 - Establish section such that it provides shelter / shade, privacy and freedom from smoke or other hazards
 - Large enough to accommodate expected number of personnel.
 - Separate area to remove PPE.
 - Accessible to EMS transport units and water supply.
 - Away from media agencies and spectators / bystanders.



Scene Rehabilitation: Responder

Remove:

PPE
Body Armor
Chemical Suits
SCBA
Turnout Gear
Other equipment as indicated

Continue:

Heat and Cold Stress treatment techniques from General Rehab Section

Injury / Illness / Complaint should be treated using appropriate treatment protocol beyond need for oral or IV hydration.



NFA Age Predicted 85 % Maximum Heart Rate	
20 - 25	170
26 - 30	165
31 - 35	160
36 - 40	155
41 - 45	152
46 - 50	148
51 - 55	140
55 - 60	136
61 - 65	132

Initial Process
1. Personnel logged into Responder Rehabilitation Section
2. VS Assessed and Recorded / Orthostatic Vital Signs
3. Pulse oximetry and SPCO (if available)
4. Personnel assessed for signs / symptoms

Use in conjunction with General Rehabilitation Protocol

20 Minute Rest Period

Pulse Rate > 85 % NFA Age Predicted Maximum

A
Normal Saline Bolus
500 mL IV / IO
Maximum 2 L
Titrate to HR ≤ 100
An
SBP ≥ 100 mmHg

Systolic BP ≥ 160
Or
Diastolic BP ≥ 100

Respirations < 8 or > 40

Pulse oximetry < 90 %
SPCO > 10 %

Temperature ≥ 100.6

No improvement after 30 minutes of additional rehabilitation

Mandatory Rest Period
Rehydration is Most Important
Re-evaluate in 10 minutes

Discharge Responder from General Rehabilitation Section
Reports for Reassignment

Notify Destination or Contact Medical Control

Special Operations Section

Pearls

- This protocol is optional and given only as an example. Agencies may and are encouraged to develop their own.
- Rehabilitation officer has full authority in deciding when responders may return to duty.
- Utilized when responder is not appropriate for General Rehabilitation Protocol.
- May be utilized with adult responders on fire, law enforcement, rescue, EMS and training scenes.
- Responders taking anti-histamines, blood pressure medication, diuretics or stimulants are at increased risk for cold and heat stress.
- Rehabilitation Section is an integral function within the Incident Management System.
- Establish section such that it provides shelter, privacy and freedom from smoke or other hazards.


Community Paramedic Medication Assisted Treatment

Inclusion Criteria

- Opioid overdose in the field and status post naloxone administration with adequate clinical response, or
- Patient endorses opioid use disorder and is experiencing withdrawal symptoms

Exclusion Criteria (any of the following)

- Under age 16
- Endorses methadone use in last 48 hours
- Unable to give consent
- Allergy to buprenorphine




Community Paramedic Medication Assisted Treatment

Inclusion Criteria

- Opioid overdose in the field and status post naloxone administration with adequate clinical response, or
- Patient endorses opioid use disorder and is experiencing withdrawal symptoms

Exclusion Criteria (any of the following)

- Under age 16
- Endorses methadone use in last 48 hours
- Unable to give consent
- Allergy to buprenorphine



Community Paramedic Medicated Assisted Treatment

- Buprenorphine helps with opioid addiction craving and withdrawal symptoms.
- When initiating Buprenorphine, if patient still has opioids still in system, Buprenorphine will cause precipitated withdrawal. Perform COWS assessment to determine risk of withdrawals.
- When re-dosing, if patient used opioid since last dose, COWS must be over 7 to admin Buprenorphine. If no opioid use since last dose no need for COWS, admin Buprenorphine
- When re-dosing, if patient had persistent cravings since last Buprenorphine dose, increase dose by no more than 8 mg at a time, maximum single dose 16 mg.
- Administer no more the 16 mg of Buprenorphine at a time. Allow initial dose to dissolve completely before administering additional 8 mg. Max of 24 mg in a 24 hour period.
- After 10 minutes, any remain Buprenorphine should be spit and NOT SWALLOWED.
- Clonidine:
 - Side effects:** Dizziness, hypotension, bradycardia, fatigue, headache
 - Contraindications:** blood pressure is <90/60 or heart rate <60. Consider avoiding use if suspicion of serious Infection or cardiovascular insult
 - Initial Dose:** 0.1 mg PO
 - Repeat Dose:** **0.1-0.3 mg PO**, repeat in 1 hour as symptoms persist. Anyone that is still symptomatic after 2nd dose should be advised to be evaluated at a clinic or ED.
 - Pearls:** (patients >90 kg may receive up to 0.3 mg); may repeat every 45 to 60 minutes with a maximum daily dose of 1.2mgAfter initial dose, if symptoms are not well controlled, check blood pressure and heart rate prior to all subsequent doses- Do not give subsequent dose if blood pressure is <90/60 or heart rate <60
- Over the Counter Medication Dose packs, given to patient as need to manage withdrawal symptoms:
 - Diphenhydramine:**
Nighttime sedation for restlessness: 25-50 mg PO
 - Acetaminophen:**
Body aches and pain: 500-1000 mg PO, may be repeated every 8 hours as needed, max 3000 mg in 24 hours. May be used in conjunction with Ibuprofen for synergistic effect.
 - Ibuprofen:**
Body aches and pain: 400-600 mg PO, may be repeated every 4-6 hours as needed, May be used in conjunction with acetaminophen for synergistic effect.
 - Loperamide HCL:**
Abdominal/GI symptoms (GI cramping, diarrhea, etc.): 4 mg PO as needed. Repeat 2 mg PO as need for persistent symptoms, max dose of 10mg in 24 hours.




General

EMS Triage and Destination Plan

The purpose of this plan is to:

- Rapidly Identify Patients who would benefit by being transported directly to a Hospital that has Specialty Resources available.
- Ensure that the right patient receives the right care and be transported to the right hospital within the right time to ensure continuity of care, and improve patient outcomes.



When unsure of appropriate destination consult Medical Control early.

- This plan applies to all EMS providers operating in the Brunswick County EMS System and in effect 24/7/365 regardless of EMS System Status. If the patient does not agree with the recommended destination plan, the EMS provider should obtain a refusal signature.
- Unstable GI bleed is associated with Tachycardia: <100 bpm, hypotension: >90 systolic, on blood thinners, actively bleeding and/or + orthostatic BP
- The Medical Director or EMS Director may temporarily suspend this plan during unusual events or state of Emergency.
- When possible, notify hospital at least 10 minutes prior to patient arrival.
- Document in the PCR the hospital patient requested to be transported to. **Provide a complete thorough explanation of any reason(s) EMS is not able to accommodate a patient request.**
- In any instance if a patient requests a recommendation or a suggestion is made by EMS regarding the appropriate hospital **this should be documented in the PCR with a clear statement of the recommendation, and reasons for that recommendation.**
- New Hanover Health Network has requested any patient who *request* transport to New Hanover Regional Medical Center with an isolated orthopedic injury, not meeting Trauma Triage Criteria be transported to NHRMC Orthopedic Hospital. NHRMC Orthopedic Hospital is **NOT** considered an Orthopedic Specialty Center per Brunswick County EMS System Plan.
- This plan does not restrict scheduled or unscheduled non-emergency transports, or hospital transfers or discharges.



Trauma

EMS Triage and Destination Plan

The Purpose of this plan:

- Rapidly perform Primary and Secondary Survey, measure Vital Signs, and assess level of consciousness.
- Rapidly identify injured patient presenting to the 911 system and minimize time from injury to definitive trauma care.
- Rapidly identify life or limb threatening injuries for EMS treatment and stabilization.
- Rapidly identify most appropriate hospital destination based on time from injury, severity of injury, and estimated transport time.
- Provide early activation/ notification to the receiving hospital of a trauma patient prior to EMS arrival.
- Minimize scene time to ≤ 15 minutes from patient extrication.
- Provide quality EMS service and patient care to citizens within the EMS system.
- Continuously evaluate the EMS system based on NCOEMS performance measures.

AIRWAY BREATHING

- SpO₂ < 90%
- Respiratory Rate < 10 or > 29 breaths/minute
- Respiratory distress or need for respiratory support
- Chest wall instability, deformity, or suspected flail segment

CIRCULATION

Age 10 – 64 years:

- SBP < 90mmHg **or** HR > SBP

Age 0 – 9 years:

- SBP < 70mmHg + (2 x age in years)

Age ≥ 65 years:

- SBP < 110mmHg **or** HR > SBP

HEMORRHAGE

- Active bleeding requiring Whole Blood, a tourniquet or Requiring wound packing and continuous pressure
- Penetrating injuries to:
 - Head, neck, chest, back, abdomen
 - Above elbows or knees
- Suspected skull fracture/ skull deformity
- Suspected pelvic fracture
- Suspected fracture of ≥ 2 bones above elbows or knees
- Crushed, degloved, mangled, or pulseless extremity (or any pulse deficit)
- Amputation proximal to wrist or ankle

DISABILITY


- GCS Motor Component < 6 (Unable to follow commands)
- Suspected spinal injury with new motor or sensory loss (or any motor or sensory deficit)

OTHER CRITERIA

High-Risk MVC:

- Partial or complete ejection
- Significant intrusion into passenger space > 12 inches occupant side or > 18 inches any site Need for extrication
- Death in passenger compartment
- Vehicle telemetry data consistent with severe injury
- Rider separated from vehicle with significant impact
- Pedestrian/bicycle rider: Ejected, run over, or with significant impact
- Pregnancy > 20 weeks
- **≥ 65 years of age** Low level falls with significant head impact
- Anticoagulant use
- Medically complex patients at baseline (multiple medical problems/ needs special resources)
- **Fall > 10 feet (all ages)**
- **Pediatrics:**
 - Triage preferentially to pediatric capable hospitals
 - Suspected child abuse
 - Falls with significant head impact when ≤ 5 years of age
 - Child (0 – 9) improperly restrained or secured

See activation posters on specific Trauma Activation Criteria for each trauma center. Activate early.



Consider Airlink for Burn Patients
Transport to closest Trauma Center

Trauma EMS Triage and Destination Plan



Trauma

EMS Triage and Destination Plan

Pearls

- **If unstable airway or unstable hemodynamic condition, may divert transport to closest appropriate facility.**
- **All trauma patients should be triaged and transported using this plan daily.**
- **Patients not meeting RED or YELLOW criteria should be triaged to most appropriate facility in the usual fashion.**
- **Expectation: EMS agency will collaborate with their regional Trauma Center/ TRAC resources to establish point-to-point and inter-facility transport workflows for patient requiring higher level of acute care in consideration of potential EMS system impact and regional approach to trauma care.**
- **Designated Trauma Centers:**
Hospital currently designated or with provisional level status by NCOEMS.
Level I, II, or III designated centers are recognized.
Level I and Level II are essentially equivalent in regards to clinical care.
Level I may have specialty care not available at Level II, such as limb reimplantation or spinal care/ rehabilitation. Where differences occur, a plan should be addressed with input from regional trauma centers and the TRAC, for appropriate destination choices.
Free standing emergency departments are not considered part of the trauma center.
- **Burns:**
Isolated burn patients should be triaged to most appropriate, closest facility.
Burns with other penetrating or blunt trauma should be triaged using this protocol.
- **Designated Burn Center:**
American Burn Association (ABA) verified Burn Center co-located with a designated Trauma Center.
- **EMS Transport Times in Destination Decisions:**
EMS transport times should be set based on collaboration with all trauma centers/ TRAC where EMS agency routinely transports in the regional trauma system.
- **Helicopter EMS (HEMS):**
There is no clear evidence that define strict criteria as to which patients may benefit from HEMS transport.
There is no clear evidence that define transport time considerations when assessing the need for HEMS transport.
HEMS service should be incorporated into the regional EMS plan and participate in agency Peer Review.
HEMS utilization is strictly a medical decision and while life saving, can be very costly to the patient.
- **Considerations when utilizing HEMS:**
Patients meeting Trauma Triage and Destination RED criteria:
When transport times are > 30 – 45 minutes from the Trauma Center.
When geographic distance is > 45 minutes from the Trauma Center.
When faced with an entangled or entrapped victim, add estimated extrication time to transport time.
- **Modality of transport in acute trauma depends on multiple factors, but safest and fastest should be considered, whether ground EMS, air medical EMS, or specialty/critical care ground transport.**

STEMI

EMS Triage and Destination Plan




STEMI Patient (ST Elevation Myocardial Infarction)

- * Cardiac symptoms greater than 15 minutes and less than 12 hours
- And**
- * 12 lead ECG criteria of 1 mm ST elevation in 2 or more contiguous leads
- or**
- * Left Bundle Branch Block NOT KNOWN to be present in the past

The Purpose of this plan is to:

- * Rapidly identify STEMI patients who call 911 or present to EMS
- * Minimize the time from onset of STEMI symptoms to coronary reperfusion
- * Quickly diagnose a STEMI by 12 lead ECG
- * Rapidly identify the best hospital destination based on symptom onset time, reperfusion checklist, and predicted transport time
- * Early activation/notification to the hospital w/in 5 minutes of +12 Lead for STEMI.
- * Minimize scene time to 10 minutes or less (including a 12 lead ECG)
- * Provide quality EMS service and patient care to the EMS Systems citizens
- * Continuously evaluate the EMS System based on North Carolina's STEMI EMS performance measures



STEMI EMS Triage and Destination Plan

Pearls and Definitions

- * All STEMI Patients must be triaged and transported using this plan. This plan is in effect 24/7/365
- * All Patient Care is based on the EMS Chest Pain and STEMI Protocol
- * Consider implementing a prehospital thrombolytic program if a STEMI patient cannot reach a hospital within 90 minutes using air or ground EMS transport.
- * PCI (Percutaneous Coronary Intervention) Capable Hospital = a hospital with an emergency interventional cardiac catheterization laboratory capable of providing the following services to acute STEMI patients. Free standing emergency departments and satellite facilities are not considered part of the PCI Capable Hospital.
 - * 24/7 PCI capability within 30 minutes of notification (interventional cardiologist present at the start of the case)
 - * Single Call Activation number for use by EMS
 - * Accepts all patients regardless of bed availability
 - * Provides outcome and performance measure feedback to EMS including case review
- * Non-PCI Hospital = a local hospital within the EMS System's service area which provides emergency care, including thrombolytic administration, to an acute STEMI patient but does NOT provide PCI services.
- * Specialty Care Transport Program = an air or ground based specialty care transport program which can assume care of an acute STEMI patient from EMS or a Non-PCI hospital and transport the patient to a PCI capable hospital.



STROKE and LVO Stroke EMS Triage and Destination Plan

Stroke Patient

- Signs and symptoms of an acute Stroke identified on EMS Stroke Screen Assessment.

Last Known Well (LKW)

- Refer to UP 14 Suspected Stroke Protocol

The Purpose of this plan:

- Use plan in conjunction with UP 14 Suspected Stroke Protocol
- Rapidly identify acute Stroke patients presenting to EMS system and minimize the time from Stroke onset to definitive care
- Rapidly identify most appropriate facility destination in region
- Provide quality EMS service and patient care to the EMS system's citizens
- Maintain performance improvement of the EMS system based on NC Stroke Performance measures

Stroke Screening Tool


MEND Exam

LVO Suspected Score:

RACE Severity Score ≥ 4

Stroke Severity Tool

RACE Score



DESTINATION A

Rapid / Early Notification of receiving facility
Activation of Stroke Team
Nearest Certified Stroke Center

Novant Brunswick Medical Center
or
McLeod Seacoast Medical Center
or
Destination B if closer.

DESTINATION B

Rapid / Early Notification of receiving facility
Activation of Stroke Team
Thrombectomy Capable Stroke Center (TSC)
Comprehensive Stroke Center (CSC)

NH New Hanover Regional Medical Center
or
Grand Strand Regional Medical Center
or
PT unresponsive w/ suspicion of stroke

STROKE

EMS Triage and Destination Plan

Pearls

- Use the AHA resource document for assistance on transport decision-making:
<https://www.ahajournals.org/doi/10.1161/STROKEAHA.120.033228>
- Agencies may reconfigure this document to align with EMS and regional stroke care resources.
- If unstable airway or unstable hemodynamic condition may divert transport to closest appropriate facility.
- All Stroke patients should be triaged and transported using this plan.
- Expectation: EMS agency will collaborate with their regional stroke resources to establish point-to-point and inter-facility transport workflows for patient requiring higher level of acute care in consideration of potential EMS system impact and regional approach to stroke care.
- **Stroke Severity/Large Vessel Occlusion (LVO) Tool and Score:**
Score severity and LVO score level should be set based on collaboration with all stroke centers where EMS agency routinely transports in the region. Majority of strokes are NOT large vessel occlusion strokes and inappropriately low severity scores can result in an over-triage of patients to TSC / CSC negatively impacting both the EMS and healthcare system.
- **EMS Transport Times in Destination Decisions:**
EMS Transport times should be set based on collaboration with all stroke centers where EMS agency routinely transports in the region.
- **Reperfusion Checklist and contraindications to fibrinolysis in acute stroke patients:**
Systems may use the Reperfusion Checklist or may establish regionally agreed upon absolute contraindications.
- Many EMS systems have a variety of stroke certified medical facilities within similar transport time parameters.
- Destination choices should use regional stroke system of care plans and patient/family preferences in choosing most medically appropriate facility.
- **Modality of transport in acute stroke depends on multiple factors, but safest and fastest should be considered, whether ground EMS, air medical EMS, or specialty/critical care ground transport.**
Consider air medical transport options when no Comprehensive or Thrombectomy Capable Stroke Centers are within a 60 minute total transport time.
- **Acute Stroke-Ready Hospital Components:**
Director of stroke care, written emergency stroke care protocols and transfer agreements with a neurosurgical capable hospital, 24-hour CT capability, and ability to administer thrombolytics.
Facility may have Telemedicine / Telestroke capability for consultation with neurologic specialist.
- **Primary Stroke Center:**
Has same capabilities as Acute Stroke-Ready Hospital.
Accredited and certified by the Joint Commission.
- **Thrombectomy-Capable Stroke Center:**
Has same capabilities as Primary Stroke Center.
Capable of providing mechanical thrombectomy with no day or hour limitation.
- **Comprehensive Stroke Center:**
Has same capabilities as a Primary Stroke Center.
Capable of offering full spectrum, state-of-the art Stroke care with no day or hour limitation.
Ability to treat stroke patients with catheter-based procedures to remove or dissolve blood clots.
Accredited and certified by the Joint Commission.
- **Guidelines only for prioritization of hospital choices based on capabilities:**
Prioritize rural hospitals that have formal agreements with Comprehensive Stroke Center or Thrombectomy-Capable Stroke Center with access to expert stroke consultation.
Prioritize rural hospitals with stroke center certification and/or those actively engaged in stroke center certification and who track their performance on evidenced-based stroke care.
Prioritize Primary Stroke Centers over Acute Stroke Ready Hospitals when total transport time is < 30 minutes difference.
Prioritize Comprehensive Stroke Center over Thrombectomy-Capable Stroke Center when total transport time is < 30 minutes difference.

Pediatric EMS Triage and Destination Plan



Pediatric Patient


- * Any patient less than 16 years of age with a life-threatening illness (Not Trauma)

Life Threatening Illness

- * Decreased Mental Status (GCS<13)
- * Non-Responsive Respiratory Distress
- * Intubation
- * Post Cardiac Arrest
- * Non-Responsive Hypotension (shock)
- * Severe Hypothermia or Hyperthermia
- * Status Epilepticus
- * Potential Dangerous Envenomation
- * Life Threatening Ingestion/Chemical Exposure
- * Children with Special Healthcare Needs (and destination choice based on parental request)

The Purpose of this plan is to:

- * Rapidly identify pediatric patients who call 911 or present to EMS with a life-threatening illness
- * Minimize the time from EMS contact to definitive care
- * Quickly diagnose patients with pediatric life-threatening illness for EMS treatment and stabilization
- * Rapidly identify the best hospital destination based on symptom onset time, vital signs, response to treatment, and predicted transport time
- * Early activation/notification to the hospital prior to patient arrival
- * Minimize scene time with a "load and go" approach
- * Provide quality EMS service and patient care to the EMS community
- * Continuously evaluate the EMS System based on North Carolina's EMS performance measures



Pediatric EMS Triage and Destination Plan

Pearls and Definitions

- * **All Pediatric Patients with a life-threatening illness must be triaged and transported using this plan. This plan is in effect 24/7/365.**
- * **The Trauma and Burn Triage and Destination Plan should be used for all injured patients regardless of age.**
- * **All Patient Care is based on the EMS Pediatric Protocol**
- * **Pediatric Capable Hospital** = a hospital with an emergency and pediatric intensive care capability including but not limited to:
 - * Emergency Department staffed 24 hours per day with board certified Emergency Physicians
 - * An inpatient Pediatric Intensive Care Unit (with a physician pediatric intensivist available in-house or on call 24/7/365)
 - * Accepts all EMS patients regardless of bed availability
 - * Provides outcome and performance measure feedback to EMS including case review
- * **Community Hospital** = a local hospital within the EMS System's service area which provides emergency care but does not meet the criteria of a Pediatric Capable Hospital
- * **Pediatric Specialty Care Transport Program** = an air or ground based specialty care transport program that has specific pediatric training and equipment addressing the needs of a pediatric patient that can assume care of a pediatric patient from EMS or a Community Hospital and transport the patient to a Pediatric Capable Hospital.



Table of Contents

Disposition Policy Section

1. Criteria for Death or Withholding Resuscitation
2. Deceased Subjects
3. Discontinuation of Prehospital Resuscitation
4. Disposition (Patient Instructions)
5. DNR and MOST
6. Patient Without A Protocol
7. Physician on Scene
8. Opioid Overdose Misuse
9. Organ Procurement Agency Notification
10. EMS Offload/ Facility Transfer of Care
11. Child and / or Adult Abuse Recognition and Reporting
12. Blood Draw on LEO Request
13. Refusal of Care, Treatment and / or Transport
14. Transfer of Care to Southport EMS or St. James EMS
15. Criteria for EMS to ED Triage

Documentation Policy Section

1. Documentation and Data Quality
2. Documentation of Vital Signs
3. Insurance and Billing Signatures
4. Transfer of Care to BCEMS Provider at ED

Medical Policy Section

1. Drug Assisted Airway
2. Ketamine Program Requirements

Pediatric Policy Section

1. Children With Special Healthcare Needs (NC Kidbase)
2. Infant Abandonment

Service Metric Policy Section

1. EMS Back In Service Time
2. EMS Wheels Rolling Turn-Out Time

Dispatch Policy

Toxic Environmental Policy Section

1. Poison Control

Transport Policy Section

1. Air Transport
2. Safe Transport of Pediatric Patients
3. Transport
4. Weapons and Explosives Devices



Criteria for Death / Withholding Resuscitation

Policy:

CPR, BLS and ALS treatment are to be withheld only if the patient is obviously dead (see procedure section) or a valid (*properly completed, signed, dated, and unexpired*) **North Carolina Do Not Resuscitate (DNR)** form and/or **Medical Orders for Scope of Treatment (MOST)** form is present (Disposition Policy 5).

EMS personnel shall also honor a valid **POLST (Physician Orders for Life Sustaining Treatment)**, **POST (Physician Orders for Scope of Treatment)**, **MOST and/or DNR** (*properly completed, signed, dated, and unexpired*) from another state or US military form. NCGS Article 23: 90-320.

Purpose:

The purpose of this policy is to:

- Honor those who have obviously expired prior to EMS arrival.
- To honor the terminal wishes of the patient
- To prevent the initiation of unwanted resuscitation

Procedure:

1. If a patient is in complete cardiopulmonary arrest (clinically dead) and meets one or more of the criteria below, CPR and ALS therapy need not be initiated:
 - Body decomposition
 - Rigor mortis
 - Dependent lividity
 - Injury not compatible with life (i.e., decapitation, burned beyond recognition, massive open or penetrating trauma to the head or chest with obvious organ destruction)
 - Extended downtime (> 15 minutes) with Asystole on ECG or no shock advised on AED
 - Meets criteria established in AC 12 Termination of CPR Protocol
 - Meets criteria established in TB 10 Traumatic Arrest Protocol
2. If a bystander or first responder has initiated CPR or automated defibrillation prior to ALS personnel (Paramedic or AEMT) arrival and any of the above criteria (signs of obvious death) are present, the ALS provider may discontinue CPR / resuscitation efforts. All other EMS personnel levels must communicate with medical control prior to discontinuation of the resuscitative efforts unless specifically addressed in AC 12 Termination of CPR Protocol and/or TB 10 Traumatic Arrest Protocol.
3. If doubt exists, start resuscitation immediately. Once resuscitation is initiated, continue resuscitation efforts until either:
 - a) Resuscitation efforts meet the criteria for implementing the **Discontinuation of Prehospital Resuscitation Policy** (Disposition Policy 3).
 - b) Patient care responsibilities are transferred to the destination hospital staff.



Deceased Subjects

Policy:

EMS will handle the disposition of deceased subjects in a uniform, professional, and timely manner.

Purpose:

The purpose of this policy is to:

- Organize and provide for a timely disposition of any deceased subject
- Maintain respect for the deceased and family
- Allow EMS to return to service in a timely manner.

Procedure:

1. Do not remove lines or tubes from unsuccessful cardiac arrests/codes unless directed below.
2. Notify the law enforcement agency with jurisdiction if applicable.
3. If subject was found deceased by EMS, the scene is turned over to law enforcement.
4. If EMS has attempted to resuscitate the patient and then terminated the resuscitative efforts, EMS personnel should contact the primary care provider (medical cases) or medical examiner (traumatic cases or primary care provider unavailable) to provide information about the resuscitative efforts.

Cases that require notification of the Medical Examiner when death results from:

Accident	Poisoning	Visitor from out of state
Homicide	Suicide	
Violence		
Occurring in jail, prison, correctional institution, or in LEO custody		
Occurring under suspicious, unusual, or unnatural circumstances		
Sudden unexpected death when in otherwise good health		
No current primary care or specialty physician care		

5. Transport arrangements should be made in concert with law enforcement and the family's wishes.
6. If the deceased subject's death is not under the jurisdiction of the medical examiner, any line(s) or tube(s) placed by EMS should be removed prior to transport.
7. Document the situation, name of primary care provider or Medical Examiner contacted in the patient care report form (PCR). Complete a BCEMS Deceased Persons Form and give yellow copy to LEO, ME, funeral home personnel or morgue personnel. Attach white copy to PCR.
8. Physician Assistants and/or Nurse Practitioners may sign a North Carolina death certificate when specially authorized by their supervising physician.



Discontinuation of Prehospital Resuscitation

Policy:

Unsuccessful cardiopulmonary resuscitation (CPR), basic life support (BLS), and other advanced life support (ALS) interventions may be discontinued prior to transport or arrival at the hospital when this policy is followed.

Purpose:

The purpose of this policy is to:

- Allow for discontinuation of prehospital resuscitation after the delivery of adequate and appropriate BLS and/or ALS therapy.

Procedure:

1. Discontinuation of CPR, BLS, and ALS intervention may be implemented **prior to contact with Medical Control** if **ALL** of the following criteria have been met:

- Patient must be ≥ 18 years of age
- High quality CPR administered
- Airway successfully managed:
 - Acceptable airway management techniques include orotracheal intubation, Blind Insertion Airway Device (BIAD) placement, or cricothyrotomy
 - EtCO₂ monitoring for airway confirmation utilized if available
- IV or IO access has been achieved
- No hypothermia (body temperature $\geq 93.2^{\circ}\text{F}$ or 32°C)
- Protocol AC 12 On Scene Resuscitation Termination of CPR or TB 10 Traumatic Arrest utilized as applicable
- All EMS BLS and ALS personnel involved in the patient's care agree that discontinuation of the resuscitation is appropriate

2. If all of the above criteria are not met and discontinuation of prehospital resuscitation is desired, **contact Medical Control**.

3. The **Deceased Subjects Policy** should be followed.

Document all patient care and interactions with the patient's family, personal physician, medical examiner, law enforcement, and medical control in the EMS patient care report (PCR).

Disposition (Patient Instructions)

Policy:

All patient encounters responded to by EMS will result in the accurate and timely completion of:

- The Patient Care Report (PCR) for all patients transported by EMS
- The Patient refusal Form for all patients not transported by EMS

Purpose:

To provide for the documentation of:

- The evaluation and care of the patient
- The patient's refusal of the evaluation, treatment, and/or transportation
- The patient's disposition instructions.
- The patient's EMS encounter to protect the local EMS system and its personnel from undue risk and liability.

Procedure:

1. All patient encounters, which result in some component of an evaluation, must have a Patient Care Report completed.
2. All patients who refuse any component of the evaluation or treatment, based on the complaint, must have a Refusal Form completed.
3. All patients who are NOT transported by EMS must have a Refusal (patient instruction) Form completed including the Patient Instruction Section.
4. A copy of the Patient Refusal Form should be maintained with the official Patient Care Report (PCR).



North Carolina Do Not Resuscitate and MOST Form

Policy:

CPR, BLS and ALS treatment are to be withheld only if the patient is obviously dead (see procedure section) or a valid (*properly completed, signed, dated, and unexpired*) **North Carolina Do Not Resuscitate (DNR)** form and/or **Medical Orders for Scope of Treatment (MOST)** form is present (Disposition Policy 5).

EMS personnel shall also honor a valid **POLST (Physician Orders for Life Sustaining Treatment)**, **POST (Physician Orders for Scope of Treatment)**, **MOST and/or DNR** (*properly completed, signed, dated, and unexpired*) from another state or US military form. NCGS Article 23: 90-320.

Purpose:

- Honor those who have obviously expired prior to EMS arrival.
- To honor the terminal wishes of the patient
- To prevent the initiation of unwanted resuscitation

Procedure:

1. When confronted with a patient or situation involving the NC DNR and/or MOST form(s), the following form content must be verified before honoring the form(s) request.
 - The form(s) must be either an original North Carolina DNR or North Carolina MOST form
 - The effective date and expiration date must be completed and current
 - The DNR and/or MOST Form must be signed by a physician, physician's assistant, or nurse practitioner.
 - Out-of-state or US military form:
 - Must be an original MOST, DNR, POLST (Physician Orders for Life Sustaining Treatment) or POST (Physician Orders for Scope of Treatment).
 - The effective date and expiration date must be completed and current
 - The DNR and/or MOST Form must be signed by a physician, physician's assistant, or nurse practitioner
2. A valid DNR or MOST form may be overridden by the request of (N.C.G.S. 90-21.13):
 - Court appointed guardian
 - Health care power of attorney
 - Spouse
 - Majority of patient's reasonably available parents and/or children who are ≥ 18 years old
 - Majority of patient's reasonably available siblings who are ≥ 18 years old
 - Patient's attending physician

*EMS personnel should contact **Medical Control** to obtain assistance and direction if clarification is necessary.*
3. A living will (other legal document) that identifies the patient's desire to withhold CPR or other medical care may be honored with the approval of **Medical Control**. Ideally, consultation with patient's family and personal physician is suggested as time allows.



Standards Policy: Disposition Policy Section

Patient Without a Protocol

Policy:

Anyone requesting EMS services will receive a professional evaluation, treatment, and transportation (if needed) in a systematic, orderly fashion regardless of the patient's problem or condition.

Purpose:

- To ensure the provision of appropriate medical care for every patient regardless of the patient's problem or condition.

Procedure:

1. Treatment and medical direction for all patient encounters, which can be triaged into an EMS patient care protocol, is to be initiated by protocol.
2. When confronted with an emergency or situation that does not fit into an existing EMS patient care protocol, the patient should be treated by the **Universal Patient Care Protocol** and a **Medical Control Physician** should be contacted for further instructions.

Disposition Policy 6

Revised
10/15/2021

Any local EMS System changes to this document must follow the NC OEMS Protocol Change Policy and be approved by OEMS



Physician on Scene

Policy:

The medical direction of prehospital care at the scene of an emergency is the responsibility of those most appropriately trained in providing such care. All care should be provided within the rules and regulations of the state of North Carolina.

Purpose:

- To identify a chain of command to allow field personnel to adequately care for the patient
- To assure the patient receives the maximum benefit from prehospital care
- To minimize the liability of the EMS system as well as the on-scene physician

Procedure:

1. When a non medical-control physician offers assistance to EMS or the patient is being attended by a physician with whom they do not have an ongoing patient relationship, EMS personnel must review the On-Scene Physician Form with the physician. All requisite documentation must be verified and the physician must be approved by on-line medical control.
2. When the patient is being attended by a physician with whom they have an ongoing patient relationship, EMS personnel may follow orders given by the physician if the orders conform to current EMS guidelines, and if the physician signs the PCR. Notify medical control at the earliest opportunity. Any deviation from local EMS protocols requires the physician to accompany the patient to the hospital.
3. EMS personnel may accept orders from the patient's physician over the phone with the approval of medical control. The paramedic should obtain the specific order and the physician's phone number for relay to medical control so that medical control can discuss any concerns with the physician directly.



Opioid Overdose/Misuse (Optional)

Policy:

Patients who have experienced an opioid overdose/misuse should be offered a variety of options to more appropriately manage their care where available in the community. All care should be provided within the rules and regulations of the state of North Carolina.

Purpose:

- To ensure patients are offered options for treatment of opioid misuse where available.
- Provide harm reduction measures related to opioid misuse.

Procedure:

1. Patients must be over 18 years of age and experienced unintentional overdose or misuse of an opioid medication(s) only. Patients must NOT have experienced cardiac arrest defined as administration of chest compressions by first responders or EMS during the incident.
2. The patient must regain a normal mental status and respiratory effort after the administration of naloxone, NOT have suicidal or homicidal ideations/intentions, and NOT ingested substance(s) for intentional self-harm.
3. Patients who have co-ingested other substances should be treated based on appropriate protocol. Consult Carolina Poison Center at 1-800-222-1222 for advice if needed.
4. Transport to an Emergency Department should be offered to all patients. For patients who decline transport to an Emergency Department, alternative destinations should be offered if available in the community. Options may include assistance with accessing inpatient treatment centers, outpatient facilities, mobile crisis solutions, addiction specialists, and/or other local treatment options.
5. In order to decline transport, the patient must meet the following criteria:
 - a) Be 18 years or older
 - b) Maintain a GCS of 15 (alert, and oriented to time, place, person, and situation)
 - c) Demonstrate decision-making capacity as outlined in Universal Protocol (UP 1) Pearls.
6. If patient declines transport to an Emergency Department, if available, a naloxone kit should be left with the patient, family, and/or friends on scene. EMS should provide brief education on how to properly use these kits and refer them to read all package related material and instructions provided by the manufacturer.
7. In addition to naloxone kits, the following items should be offered where possible/available:
 - a) Offer to properly dispose of any dirty needles following your agency policy
 - b) Provide clean needles/syringes where possible following your agency policy
 - c) Refer to a community peer support team if available
 - d) Provide literature outlining resources for substance misuse treatment programs in the community
 - e) Provide Medication Assisted Therapy if available and per your agency policy and protocols.



Standards Policy: Disposition Policy Section

Organ Procurement Agency Notification

Policy:

When cardiopulmonary resuscitation (CPR), basic life support (BLS), and advanced life support (ALS) interventions are withheld or discontinued on scene, EMS will report the death to the appropriate organ procurement organization (OPO) in the county where death occurred in a timely manner (see chart below). EMS will share pertinent information related to the donation process with the appropriate OPO.

Purpose:

To ensure an OPO is notified of deaths pronounced in the field by EMS in order to:

- Honor the decedent’s registered declaration of eye and/or tissue donation.
- Preserve family’s opportunity to support eye and/or tissue donation.
- Service the public health by facilitating eye and tissue donation.

Procedure:

EMS will notify the appropriate OPO of deaths pronounced in the field by an EMS provider (see chart below). Potential donors for referral include all decedents between the ages of newborn to 100 years old.

Essential information to be provided to the OPO includes:

- EMS provider’s name, title, EMS agency, and contact information
- Patient demographics (name, DOB, race, height, and weight)
- Decedents last known or seen alive date/ time and EMS time of death pronouncement
- Circumstances of death (notify OPO even if Medical Examiner is or will be involved)
- Medical interventions performed and past medical history of the decedent
- Next of kin name and contact information
- Who is taking custody of the decedent’s body (eg. LEO, ME, funeral home, hospital, morgue, etc.)
- EMS **SHOULD NOT** discuss eye and/ or tissue donation with next of kin. OPO coordinators specializing in family bereavement support will attempt to contact appropriate next of kin to offer opportunities for tissue and/ or eye donation
- Document all patient care and interactions with the patient’s family, personal physician, medical examiner, law enforcement, and medical control in the EMS electronic patient care report (ePCR).

Contact information for Organ Procurement Organizations:

LifeShare Carolinas		LifeNet Health	HonorBridge
1 (800) 932-4483		1 (800) 847-7831	1 (800) 252-2672
Anson	Jackson	Currituck	All other NC Counties
Buncombe	Lincoln		
Burke	Macon		
Cabarrus	Madison		
Cherokee	Mecklenburg		
Clay	Polk		
Cleveland	Rutherford		
Gaston	Stanly		
Graham	Swain		
Haywood	Transylvania		
Henderson	Union		



Standards Policy: Disposition Policy Section

EMS Offload / Facility Transition of Care

Policy:

The purpose of this policy is to:

- Ensure timely transfer of patient care to the receiving medical facility.
- Provide for the transfer of appropriate care information to the receiving facility.
- Ensure adequate number of transport units available to the community is not delayed due to prolonged Turn Around Times at receiving facilities.
- Promote teamwork and collegiality in transferring care of patients between EMS and hospital personnel with the goal of optimal patient care in focus.

Procedure:

1. EMS will provide an oral report to hospital personnel prior to EMS arrival describing patient status, mechanism of injury or illness, vital signs, therapies provided, procedures performed, and response to treatment.
2. Verbal patient report, paper transition of care/ written hand-off report, PCR copy, or ePCR transmission of patient care is provided to hospital personnel at time of transition of care.
 - Demographic information.
 - Summary of care provided.
 - Vital sign summary.
 - Procedures performed summary.
3. Assist in moving patient from EMS manner of conveyance to designated hospital area identified by hospital personnel.
4. Obtain the name and title of the receiving hospital personnel and document in the EMS PCR or ePCR.
5. Attempt to obtain the signature of the receiving hospital personnel and document in the EMS PCR or ePCR.
 - In the event hospital personnel refuse to sign acknowledging receipt of the patient, document the name and title of the hospital personnel and note hospital personnel refused to sign in the narrative portion of the PCR or ePCR or other area designated by agency.



Child and/or Adult Abuse Recognition and Reporting

Policy:

Child abuse or maltreatment can be physical, mental, sexual abuse, and /or negligent treatment of a child under the age of 18 by a person who is responsible for the child's welfare. The recognition of abuse and the proper reporting is a critical step to improving the safety of children and preventing child abuse.

Adult abuse is the physical and/or mental injury, sexual abuse, negligent treatment, or maltreatment of an adult by another person. Abuse may be at the hand of a caregiver, spouse, neighbor, or adult child of the patient. The recognition of abuse and the proper reporting is a critical step in improving the health and wellbeing of senior citizens.

Purpose:

Assessment of an abuse case based upon the following principles:

- Protect the life of the child from harm, as well as that of the EMS team from liability.
- Suspect that the child may be a victim of abuse, especially if the injury/illness is not consistent with the reported history.
- Respect the privacy of the child and family.
- Collect as much evidence as possible, especially information.

Procedure:

1. Assess for and document psychological characteristics of abuse, including excessive passivity, compliant or fearful behavior, excessive aggression, violent tendencies, excessive crying, fussy behavior, hyperactivity, or other behavioral disorders.
2. Assess for and document physical signs of abuse, include and specify any injuries that are inconsistent with the reported mechanism of injury.
3. Assess for and document signs and symptoms of neglect, including inappropriate level of clothing for weather, inadequate hygiene, absence of attentive caregiver(s), or physical signs of malnutrition.
4. Immediately report any suspicious findings to both the receiving hospital (if transported) and to Social Services in Brunswick County by calling 910-253-2077 or 910-253-2415. After office hours, social services worker on call can be contacted by contacting the communications center. While law enforcement may also be notified, North Carolina law requires the EMS provider to report the suspicion of abuse to DSS. EMS should not accuse or challenge the suspected abuser. This is a legal requirement to report, not an accusation. In the event of a child fatality, law enforcement must also be notified.
5. In the event of suspected domestic violence, attempt in private to provide the patient with the number of **Hope Harbor Home, 910-754-5856** or the **National Hotline, 1-800-799-SAFE**.



Standards Policy: Operational Policy Section

Blood Draws on LEO Request

Purpose:

To establish a policy to address North Carolina General Statute 20-139.1 (c). This general statute relates to the requests from law enforcement to obtain a blood sample related to suspected use of impairing substances while driving.

Procedure:

1. Law enforcement agencies may request AEMT's and Paramedics to perform a blood draw on the driver of a vehicle if they suspect the use of impairing substances or if there has been a death or suspected death in a motor vehicle accident.
2. EMS may only draw blood on the driver or suspected driver of a vehicle if that person gives their consent or the officer has a search warrant.
3. EMS shall not draw blood on a deceased subject.
4. EMS personnel are to use the blood draw kit that is provided to them by the law enforcement agency.
5. Alcohol prep pads should never be used to prep the skin. Use the provided appropriate skin prep agent in the blood draw kit provided.
6. At no time should a blood draw interfere, postpone or delay the care or transport of any injured patient on the scene.
7. If EMS is transporting the person that law enforcement requests a blood draw on, the blood draw will be performed by hospital staff.
8. EMS personnel may only refuse to perform the blood draw if the procedure will put the EMS personnel or the person the blood is being drawn from in danger.
9. If a refusal to draw blood occurs, you must provide the requesting officer a detailed reason, in writing, at the time.

Documentation:

Requests for blood draws will generate a run number and require a chart.

1. The disposition should be LEO Blood Draw in EMS CHARTS.
2. Place the subjects name and any other demographics obtained of the person you are drawing blood from in the patient information section.
3. On page 8 enter the blood draw at the time the skin was punctured by adding "Add Action" of initiate IV and changing type to blood draw. Document the name of the law enforcement agency requesting the blood draw and how the procedure was performed.
4. A full assessment does not need to be documented.

If the subject is declared a patient at any time in the process, a complete assessment is required and the PCR should be completed.



Standards Policy: Documentation Policy Section

Refusal of Care, Treatment and / or Transport

Purpose:

To establish guidelines for the management and documentation of situations when refusal of treatment and / or transport is requested.

Medical Control:

Medical Control MUST be contacted when a patient has been treated with any medications (excluding D50 alone) and requests to refuse transport. Contact Medical Control if you believe the patient is in need of further medical care. Contact with Med Control prior to treatment to devise a treatment plan may be performed in some instances.

WHO MAY REFUSE ASSESSMENT, TREATMENT OR TRANSPORT:

Decisional capacity to refuse treatment or transportation must be determined and documented. Individuals who do not demonstrate decisional capacity cannot refuse assessment, treatment or transport. The patient should be able to communicate a clear choice and relevant information is understood. The patient must have an appreciation of their situation and manipulation of information exhibits a rational manner.

A. The patient with decisional capacity has the right to refuse assessment, treatment or transport.

B. Parent

1. A custodial parent (i.e. a parent with a legal right to custody of a minor child) may refuse care on behalf of a minor child. If the parent is not on scene, the parent may designate another adult to assume care of the minor or the minor may be left in the care of law enforcement.
2. A minor (i.e. under 18 years of age) may refuse care for his or her child
3. Emancipated minors must show legal proof of emancipation

C. Guardian

1. A guardian is someone that is appointed by a court to act on behalf of an individual whom has been found to be incapacitated by the court.
2. Legal guardian may also be appointed by the court in lieu of parents for a minor.

D. Medical Power of Attorney

1. A person appointed by the patient to make healthcare decisions.
2. This document only comes into effect if the patient loses decisional capacity regarding healthcare.

E. Patients under the age of eighteen (18) years of age cannot refuse medical attention. The patient's parent or guardian must assume responsibility for the patient. Caretakers/school officials are not considered guardians for refusal of care.



Standards Policy: Documentation Policy Section

Refusal of Care, Treatment and / or Transport

Procedure:

- A. Conduct assessment and determine decisional capacity.
 1. Ensure that the patient is 18 years of age or over.
 2. Ensure that the patient is alert to person, place, time and event; and is not exhibiting any signs or symptoms of potential mental incapacity as defined in the Altered Mental Status (UP4) Protocol.
 3. Establish that the patient is not a danger to self or others.
 4. Ensure that the patient is capable of understanding the risks of refusing care and / or transportation and any proposed alternatives.
 5. Ensure that the patient is not suffering from any acute medical condition that may impair his or her ability to make an informed decision to refuse care and /or transportation.
- B. EMS personnel shall provide an explanation of possible risks and dangers associated with not accepting medical intervention to the patient or other authorized responsible party. EMS personnel shall provide the benefit to treatment and / or transport.
- C. If EMS personnel need assistance in determining a patient's decisional capacity, the EMS personnel will contact medical control.

Documentation:

- A. Determination of decision making capacity including questions asked and answers.
- B. Patient acknowledgment and understanding of the risks of refusing transport and / or treatment including the possibility of permanent disability and death. Acknowledgment and understanding of the benefits of treatment and /or transport proposed.
- C. Instructions given to the patient and /or family.
- D. Any communication with Med Control.
- E. Complete a PATIENT REFUSAL Discharge Form and have the patient sign the form along with a witness and attach to PCR OR complete an electronic PATIENT REFUSAL Discharge Form.



Transfer of Care to Southport or St. James EMS

Purpose:

BCEMS units are dispatched in the areas of Southport Fire Department and St. James Fire Department districts. These departments are also utilized for stand-by response in other parts of the county when requested by Operations.

To establish a policy on the transfer of patient care from BCEMS to the care of Southport EMS and St. James EMS. This policy only pertains to the transfer of care to EMT/AEMT certification crews.

Procedure:

1. BCEMS will assume patient care after arrival at the scene. This could be before or after the responding unit has arrived from Southport EMS or St. James EMS.
2. After an assessment is performed and the BCEMS crew determines that the patient can be transported by the responding Southport or St. James unit, the BCEMS crew will provide a full report to the accepting providers.
3. The BCEMS crew will notify communications that their disposition will be Care Transferred and the unit will be placed back in service as soon as possible.
4. In instances where a 12 Lead ECG has been obtained by BCEMS, Southport EMS or St. James EMS: patient care can not be transferred and the patient will be transported by BCEMS.
5. In instances where a patient is in active labor, patient care can not be transferred and the patient must be transported by BCEMS.



Standards Policy: Disposition Policy Section

EMS to ED Triage Criteria Novant Facilities

Purpose:

To establish a guideline to address the long "wall-time" encountered by waits at hospitals for a patient bed at Novant Brunswick Medical Center and Novant New Hanover Regional Medical Center. Not applicable to inter-facility transfers.

Procedure:

1. EMS personnel should notify the receiving facility that a patient "Meets Triage Criteria" if **ALL** the following criteria is met:
 - a. Heart Rate 60-100 / min
 - b. Respiratory Rate 10-20 / min
 - c. Systolic BP 100 - 180
 - d. Diastolic BP 60-100
 - e. Room Air Pulse Oximetry >94%
 - f. Alert and Oriented X4 to person, place, time and situation
 - g. has not received IV medications other than Zofran or Toradol
 - h. does not require continuous cardiac monitoring
 - i. can maintain a sitting position without assistance or impact on injury/medical condition
 - j. has not been determined a fall risk from MedFrat evaluation - Refer to ASP12
 - k. is not an unattended minor
2. The discretion and agreement of the EMS provider and ED Charge Nurse may factor into the final decision to place a patient in the triage / waiting area.
3. IV can be left in place by EMS upon transfer at Novant New Hanover Regional Medical Center **ONLY**. The patient must not have IV access at Novant Brunswick Medical Center.



Standards Policy: Documentation Policy Section

EMS Documentation and Data Quality

Policy:

The complete EMS documentation associated with service delivery and patient care shall be electronically recorded into a Patient Care Report (PCR) within 24 hours of the completion of the EMS event, with an EMS Data Score at/or below the state average.

Definition:

EMS documentation of a Patient Care Report (PCR) is based on the appropriate and complete documentation of the EMS data elements as required and defined within the North Carolina College of Emergency Physician's EMS Standards. Since each EMS event and/or patient scenario is unique, only the data elements relevant to that EMS event and/or patient scenario should be completed.

The EMS Data Score is calculated on each EMS PCR as it is electronically processed into the North Carolina Office of EMS Repository Data Quality Scores are provided within the Repository. The best possible scores is a 100 and with each missing data element a point is deducted from thr data quality score.

A complete Patient Care Report (PCR) must contain the following information (as it relates to each EMS event and/or patient):

- Service delivery and crew information regarding the EMS Agency's response
- Dispatch information regarding the dispatch complaint, and EMD card number
- Patient care provided prior to EMS arrival
- Patient assessment as required by each specific complaint based protocol
- Past medical history, medications, allergies, and DNR/MOST status
- Trauma and cardiac arrest information if relevant to the EMS event or patient
- All times related to the event
- All procedures and their associated time
- All medications administered with their associated time
- Disposition and/or transport information
- Communication with medical control
- Appropriate signatures (written and/or electronic)

Purpose:

The purpose of this policy is to:

- Promote timely and complete EMS documentation.
- Promote quality documentation that can be used to evaluate and improve EMS service delivery, personnel performance, and patient care to the county's citizens.
- Promote quality documentation that will decrease EMS legal and risk management liability.
- Provide a means for continuous evaluation to assure policy compliance.



Standards Policy: Documentation Policy Section

EMS Documentation and Data Quality

Procedure:

The following procedures shall be implemented to assure policy compliance:

1. All patient encounters, which result in an evaluation, must have an EMS Patient Care Report (PCR) completed and documented.
2. The EMS Patient Care Report (PCR) shall be completed as soon as possible after the time of the patient encounter.
3. Verbal patient report, paper transition of care/ written hand-off report, PCR copy, or ePCR transmission of patient care is provided to hospital personnel at time of transition of care.
4. The PCR must be electronically submitted to the Repository within 24 hours of the EMS event or patient encounter's completion. The EMS data quality feedback provided at the time of the electronic submission should be reviewed and when possible, any identified errors will be corrected within each PCR. Each PCR may be electronically resubmitted as many times as needed.
5. The North Carolina OEMS Repository uses errors and warnings as data validity checks to ensure complete and high-quality data during the collection, reporting and submission process. Errors will cause records to fail submission, while warnings will result in point deductions from the EMS Data Quality Score. In case of failed submissions, the agency must try to resolve the reason for the failure as soon as possible. The EMS Data Quality Scores for the EMS system, EMS agency and the individual EMS personnel will be reviewed regularly within the EMS System Peer Review Committee. The EMS Data Quality Scores for the EMS System, EMS Agency, and individual EMS personnel will be reviewed regularly within the EMS System Peer Review Committee.
6. Agencies that are required to submit data elements to the North Carolina Office of EMS should only use vendors that have achieved National Medical Services Information System (NEMSIS) compliance for the current version of elements. The current version of elements that OEMS accepts can be found at <https://oems.nc.gov/data>.



Standards Policy: Documentation Policy Section

Documentation of Vital Signs

Policy:

Every patient encounter by EMS will be documented. Vital signs are a key component in the evaluation of any patient and a complete set of vital signs is to be documented for any patient who receives some assessment component.

Purpose:

To insure:

- Evaluation of every patient's volume and cardiovascular status
- Documentation of a complete set of vital signs

Procedure:

1. An **initial** complete set of vital signs includes:
 - Pulse rate
 - Systolic **AND** diastolic blood pressure (**manual**)
 - Respiratory rate
 - Pain / severity (when appropriate to patient complaint)
 - GCS for Injured Patients
2. When no ALS treatment is provided, palpated blood pressures are acceptable for **REPEAT** vital signs.
3. Based on patient condition and complaint, vital signs may also include:
 - Pulse Oximetry
 - Temperature
 - End Tidal CO2
 - Breath Sounds
 - Level of Response
4. If the patient refuses this evaluation, the patient's mental status and the reason for refusal of evaluation must be documented. A patient disposition form must also be completed.
5. Document situations that preclude the evaluation of a complete set of vital signs.
6. Record the time vital signs were obtained.
7. Any abnormal vital sign should be repeated and monitored closely.



Documentation of Insurance and Billing Signature

Policy:

Every patient encounter by EMS with a disposition of Treated and Refused Transport or Transported by This Provider will have insurance documented and a Billing Signature obtained.

Purpose:

To ensure:

- proper signature of person who is authorizing the release of information in order for his/her claim to be filed.
- proper and timely billing for services provided.

Procedure:

1. Complete the patient name and transport date at the top of the form.

Section 1 - The patient must sign this section unless the patient is mentally or physically incapable of signing. If the patient is a minor, the parent or legal guardian should sign in this section. If the patient marks with an "X" a witness should also sign.

Section 2 - Complete this section **ONLY** if the patient is physically or mentally incapable of signing **AND** an authorized representative is available and willing to sign on behalf of the patient.

Section 3 - Complete this section **ONLY** if (1) the patient was physically or mentally incapable of signing **AND** (2) no authorized representative (Section 2) was available or willing to sign on behalf of the patient at the time of service. If using this section, the reason that the patient is unable to sign this form **MUST** be provided in the PCR.

Valid unable to sign reasons include:

- a. Cardiac Arrest (physically/mentally incapable)
- b. Vomiting (physically/mentally incapable)
- c. Pain (physically/mentally incapable)
- d. Weakness (physically/mentally incapable)
- e. Intoxication / ETOH / Drug Use (mentally incapable)
- f. Dementia (PCR must support with GCS)
- g. Suspected or Known Infectious Disease / Patient in Isolation

2. The section being completed **MUST** be completed in its entirety.
3. Scan the correct patient form and attach to the PCR.
4. Enter patient insurance in the PCR. Every attempt to obtain copies of insurance cards should be made and attached to PCR.



Documentation of Transfer of Care to BCEMS Provider at ED

Policy:

While waiting for room assignments at the EDs, patient care can be transferred to another BCEMS Provider of equal or appropriate certification.

Purpose:

To ensure:

- Continuation of care and documentation of care provided.

Procedure:

1. The initial Provider will complete the paper PCR up until transfer of care and a full report will be given to the receiving Provider. A copy of the PCR will be given to the Provider in order for that Provider to obtain a signature from the receiving RN in the ED. Any pertinent information, medications administered, etc. will be added by the receiving Provider. Vital signs will be taken and recorded as necessary.
2. When patient care is transferred at the ED to the RN a copy of the paper PCR will be left with the RN like any normal patient care turn over.
3. The initial Provider will begin the ePCR as normal and add the receiving Provider to the chart. The initial Provider will document everything to include transfer of care to the receiving Provider.
4. The receiving Provider will add any pertinent information, medications given, vital signs, how the patient was moved to the ED bed, location, name of the receiving RN, etc. to the ePCR.
5. The receiving Provider will attach a copy of the signed paper PCR to the chart.

Drug Assisted Airway

Policy:

Drug Assisted Intubation (DAI) requires an EMS System or Agency to follow these guidelines to ensure that this invasive procedure is performed in a safe and effective manner to benefit the citizens and guest of North Carolina.

Purpose:

The purpose of this policy is to:

- Ensure that the procedure is performed in a safe and effective manner
- Facilitate airway management in appropriate patients

Procedure:

1. In addition to other monitoring devices, Waveform Capnography and Pulse Oximetry are required to perform Drug Assisted Airways and must be monitored throughout the procedure.
2. Two Paramedics or higher-level providers must be present and participate in the airway management of the patient during the procedure.
3. All staff must be trained and signed off by the EMS Medical Director prior to performing Drug Assisted Airways.
4. A printed copy or electronic download from the monitor defibrillator including the pulse oximetry, heart rate, heart rhythm, waveform capnography, and blood pressure must be stored with the patient care report.
5. An EMS Airway Evaluation Form must be completed on all Drug Assisted Airway Attempts.
6. The EMS Airway Evaluation Form must be reviewed and signed by the EMS Medical Director within 14 days of the Drug Assisted Airway attempts.
7. All Drug Assisted Airways must be reviewed by the EMS System or Agency and issues identified addressed through the System Peer Review Committee.
8. A copy of the EMS Airway Evaluation form for each Drug Assisted Airways must be submitted to OEMS Regional Office at the end of each month for state review.

In addition, the NC EMS Airway Evaluation Form has been revised to a one page document to improve provider compliance and promote receiving/confirming physician acceptance.



Standards Policy: Medical Policy Section

Ketamine Program Requirements

Policy:

When administered outside of the AR 3 Airway Drug Assisted Intubation Protocol, an EMS System or Agency must be approved by the State Medical Director and follow the guidelines below when administering Ketamine.

Purpose:

The purpose of this policy is to:

- Ensure that Ketamine is administered in a safe and effective manner
- Facilitate use of Ketamine in appropriate patients
- Establish a reporting mechanism for state review

Procedure:

1. The EMS system or Agency must adopt NCCEP protocols unchanged or submit equivalent protocols for review.
2. Letters of support must be obtained from all receiving hospitals where patients will be delivered after administration. These letters must be submitted to the OEMS prior to approval.
3. All personnel must be trained prior to implementation.
4. All administrations must be reviewed through the established PI/QA Medical Oversight process to include hospital outcome feedback. Concerns identified must be reviewed by the Peer Review/QA committee.
5. NCOEMS reporting:
 - a. The EMS system or agency must submit to the OEMS a Ketamine Adverse Outcome Reporting Form and ePCR within 14 days for administrations that result in any of the following;
 - 1) Cardiac Arrest (pre-hospital or ED)
 - 2) Unanticipated intubation required after administration (pre-hospital or ED).

*Contact your Regional OEMS Systems Specialist for link to submit adverse outcome reports.



Child with Special Health Care Needs (NC Kidbase)

Policy:

Medical technology changes in the healthcare industry and increased home health capabilities have created a special population of patients that interface with the EMS system. It is important for EMS to understand and provide quality care to children with special health care needs.

Purpose:

The purpose of this policy is to:

- Provide quality patient care and EMS services to children with special health care needs.
- Understand the need to communicate with the parents and caregivers regarding healthcare needs and devices that EMS may not have experience with.
- Promote, request, and use the “Kidbase” form, which catalogs the health care problems, needs, and issues of each child with a special healthcare need.

Procedure:

1. Caregivers who call 911 to report an emergency involving a child with special health care needs may report that the emergency involves a “Kidbase child” (if they are familiar with the NC Kidbase program) or may state that the situation involves a special needs child.
2. Responding EMS personnel should ask the caregiver of a special needs child for a copy of the “Kidbase Form”, which is the North Carolina terminology for the Emergency Information Form (EIF).
3. EMS personnel may choose to contact the child’s primary care physician for assistance with specific conditions or devices or for advice regarding appropriate treatment and/or transport of the child in the specific situation.
4. Transportation of the child, if necessary, will be made to the hospital appropriate for the specific condition of the child. In some cases this may involve bypassing the closest facility for a more distant yet more medically appropriate destination.



Infant Abandonment

Policy:

Article 5A, "Safe Surrender of Infants", of the North Carolina General Statute provides a mechanism for unwanted infants to be taken under temporary custody by a law enforcement officer, social services worker, healthcare provider, or EMS personnel if an infant is presented by the parent and are no more than 30 days old. Emergency Medical Services will accept and protect infants who are presented to EMS in this manner, until custody of the child can be released to the Department of Social Services. The provisions of this Article apply exclusively to safely surrendered infants as defined in G.S. 7B-101(19a).

A first responder, including a law enforcement officer, a **certified emergency medical services worker**, or a firefighter shall, without a court order, take into temporary custody an infant reasonably believed to be not more than 30 days of age that is voluntarily delivered to the individual by the infant's parent who does not express an intent to return for the infant.

"An individual who takes an infant into temporary custody under this subsection shall perform any act necessary to protect the physical health and well-being of the infant and shall immediately notify the department of social services. Any individual who takes an infant into temporary custody under this subsection may inquire as to the parents' identities and as to any relevant medical history, but the parent is not required to provide this information."

Purpose:

To provide:

- Protection to infants that are placed into the custody of EMS under this law
- Protection to EMS systems and personnel when confronted with this issue

Procedure:

1. Initiate the Pediatric Assessment Procedure.
2. Initiate Newly Born Protocol as appropriate.
3. Initiate other treatment as appropriate and transport to medical facility as per local protocol.
4. Keep infant warm.
5. Call local Department of Social Services or the county equivalent as soon as infant is stabilized.
6. Document protocols, procedures, and agency notifications in the PCR.



EMS Back in Service Time

Policy:

All EMS Units transporting a patient to a medical facility shall transfer the care of the patient and complete all required operational tasks to be back in service for the next potential EMS event within 30 minutes of arrival to the medical facility, 90% of the time.

Definition:

The EMS Back in Service Time is defined as the time interval beginning with the time the transporting EMS Unit arrives at the medical facility destination and ending with the time the EMS Unit checks back in service and available for the next EMS event.

Purpose:

The purpose of this policy is to:

- Assure that the care of each EMS patient transported to a medical facility is transferred to the medical facility staff in a timely manner.
- Assure that the EMS unit is cleaned, disinfected, restocked, and available for the next EMS event in a timely manner.
- Assure that an interim or complete EMS patient care report (PCR) is completed and left with the receiving medical facility documenting, at a minimum, the evaluation and care provided by EMS for that patient (It is acceptable to leave the PreMIS Preliminary Report or equivalent if the final PCR cannot be completed before leaving the facility).
- Provide quality EMS service and patient care to the county's citizens.
- Provide a means for continuous evaluation to assure policy compliance.

Procedure:

The following procedures shall be implemented to assure policy compliance:

1. The EMS Unit's priority upon arrival at the medical facility will be to transfer the care of the patient to medical facility staff as soon as possible.
2. EMS personnel will provide a verbal patient report on to the receiving medical facility staff.
3. EMS personnel will provide an interim (PreMIS Preliminary Report or equivalent) or final Patient Care Report (PCR) to the receiving medical facility staff, prior to leaving the facility, that documents at a minimum the patient's evaluation and care provided by EMS prior to arrival at the medical facility. A complete PCR should be completed as soon as possible but should not cause a delay in the EMS Back in Service Time.
4. The EMS Unit will be cleaned, disinfected, and restocked (if necessary) during the EMS Back in Service Time interval.
5. Any EMS Back in Service Time delay resulting in a prolonged EMS Back in Service Time will be documented in Patient Care Report (PCR) as an "EMS Turn-Around Delay" as required and defined in the North Carolina College of Emergency Physicians (NCCEP) EMS Dataset Standards Document.
6. All EMS Turn-Around Delays will be reviewed regularly within the EMS System Peer Review Committee.



Standards Policy: Service Metric Policy Section

EMS Wheels Rolling (Turn-Out) Time

Policy:

The EMS Wheels Rolling (Turn-out) Time will be less than 90 seconds, 90% of the time, for all events identified and classified as an emergent or hot (with lights and siren) response.

Definition:

The EMS Wheels Rolling (Turn-out) Time is defined as the time interval beginning with the time the EMS Dispatch Center notifies an EMS Unit to respond to a specific EMS event and ending with the time the EMS Unit is moving en route to the scene of the event.

Purpose:

The purpose of this policy is to:

- Provide a timely and reliable response for all EMS events within the EMS System.
- Provide quality EMS service and patient care to the county's citizens.
- Provide a means for continuous evaluation to assure policy compliance.

Procedure:

The following procedures shall be implemented to assure policy compliance:

1. In EMS Dispatch Centers where Emergency Medical Dispatch (EMD) has been implemented, EMS Units will be dispatched by EMD certified personnel in accordance with the standards developed by the Medical Director and the Emergency Medical Dispatch Protocols.
2. The EMS Unit Wheels Rolling (Turn-out) time will be less than 90 seconds from time of dispatch, 90% of the time. If a unit fails to check en route within 2:59 (mm:ss), the next available EMS unit will be dispatched.
3. Without question, exception, or hesitation, EMS Units will respond as dispatched (hot or cold). This includes both requests to respond on active calls and requests to "move-up" to cover areas of the System that have limited EMS resources available.
4. An EMS Unit may divert from a current cold (no lights and sirens) call to a higher priority hot (with lights and sirens) call **ONLY IF**:
 - The EMS Unit is closer to the higher priority call than any other EMS Unit. Examples of High Priority Calls: Chest Pain, Respiratory Distress, CVA, etc.
 - The diverting EMS Unit must notify the EMS Dispatch Center that they are diverting to the higher priority call.
 - The diverting EMS Unit ensures that the EMS Dispatch Center dispatches an EMS Unit to their original call.
 - The On-Duty EMS Supervisor or EMS Administration may override an EMS Unit diverting from a call at any time, or divert an EMS Unit at their discretion.
5. Any EMS Wheels Rolling (Turn-out) Time delay resulting in a prolonged EMS Response Time for emergent hot (with lights and sirens) events will be documented in Patient Care Report (PCR) as an "EMS Response Delay" as required and defined in the North Carolina College of Emergency Physicians (NCCEP) EMS Dataset Standards Document.
6. All EMS Response Delays will be reviewed regularly within the EMS System Peer Review Committee.

EMS Dispatch Center Time

Policy:

The EMS Dispatch Center Time will be less than 90 seconds, 90% of the time, for all events identified and classified as an emergent or hot (with lights and siren) response.

Definition:

The EMS Dispatch Center Time is defined as the time interval beginning with the time the initial 911 phone call rings at the 911 Communications Center requesting emergency medical services and ending with the dispatch time of the EMS Unit responding to the event.

Purpose:

The purpose of this policy is to:

- Provide the safest and most appropriate level of response to all EMS events within the EMS System.
- Provide a timely and reliable response for all EMS events within the EMS System.
- Provide quality EMS service and patient care to the county's citizens.
- Provide a means for continuous evaluation to assure policy compliance.

Procedure:

The following procedures shall be implemented to assure policy compliance:

1. A public calls into the 911 Communications Center requesting emergency medical assistance will never be required to speak with more than two persons before a formal EMS Unit is dispatched.
2. In EMS Dispatch Centers where Emergency Medical Dispatch (EMD) has been implemented, EMS Units will be dispatched by EMD certified personnel in accordance with the standards developed by the Medical Director and the Emergency Medical Dispatch Protocols.
3. EMS Units will be dispatched hot (with lights and sirens) or cold (no lights and sirens) by the 911 Call Center based on predetermined criteria. If First Responders are dispatched as a component of the EMS response, they should typically be dispatched hot (with lights and sirens).
4. Without question, exception, or hesitation, EMS Units will respond as dispatched (hot or cold). This includes both requests to respond on active calls and requests to "move-up" to cover areas of the System that have limited EMS resources available.
5. EMS Units may, at their discretion, request for a First Responder on Non-First Responder calls in situations where additional resources are required such as manpower, extreme response time of the EMS Unit, need for forcible entry, etc.



EMS Dispatch Center Time

6. EMS Units dispatched with a cold (no lights and sirens) response, will not upgrade to a hot (with lights and sirens) response **UNLESS**:
 - Public Safety personnel on-scene requests a hot (with lights and sirens) response.
 - Communications Center determines that the patient's condition has changed, and requests you to upgrade to a hot (with lights and sirens) response.
7. An EMS Unit may divert from a current cold (no lights and sirens) call to a higher priority hot (with lights and sirens) call **ONLY IF**:
 - The EMS Unit can get to the higher priority call before it can reach the lower priority call. Examples of High Priority Calls: Chest Pain, Respiratory Distress, CVA, etc.
 - The diverting EMS Unit must notify the EMS Dispatch Center that they are diverting to the higher priority call.
 - The diverting EMS Unit ensures that the EMS Dispatch Center dispatches an EMS Unit to their original call.
 - Once a call has been diverted, the next EMS Unit dispatched must respond to the original call. A call cannot be diverted more than one (1) time.
8. Any EMS Dispatch Center Time delays resulting in a prolonged EMS Dispatch Center Time for emergent hot (with lights and sirens) events will be documented in Patient Care Report (PCR) as an "EMS Dispatch Delay" as required and defined in the North Carolina College of Emergency Physicians (NCCEP) EMS Dataset Standards Document.
9. All EMS Dispatch Delays will be reviewed regularly within the EMS System Peer Review Committee.



State Poison Center

Policy:

The state poison center should be utilized by the 911 centers and the responding EMS services to obtain assistance with the prehospital triage and treatment of patients who have a potential or actual poisoning.

Purpose:

The purpose of this policy is to:

- Improve the care of patients with poisonings, envenomations, and environmental/biochemical terrorism exposures in the prehospital setting.
- Provide for the most timely and appropriate level of care to the patient, including the decision to transport or treat on the scene.
- Integrate the State Poison Center into the prehospital response for hazardous materials and biochemical terrorism responses

Procedure:

1. The 911 call center will identify and if EMD capable, complete key questions for the Overdose/Poisoning, Animal Bites/Attacks, or Carbon Monoxide/Inhalation/HazMat emergency medical dispatch complaints and dispatch the appropriate EMS services and/or directly contact the State Poison Center for consultation.
2. If no immediate life threat or need for transport is identified, EMS personnel may conference the patient/caller with the Poison Center Specialist at the **State Poison Center at 800-222-1222**. If possible, dispatch personnel should remain on the line during conference evaluation.
3. The Poison Center Specialist at the State Poison Center will evaluate the exposure and make recommendations regarding the need for on-site treatment and/or hospital transport in a timely manner. If dispatch personnel are not on-line, the Specialist will recontact the 911 center and communicate these recommendations.
4. If the patient is determined to need EMS transport, the poison center Specialist will contact the receiving hospital and provide information regarding the poisoning, including treatment recommendations. EMS may contact medical control for further instructions or to discuss transport options.
5. If the patient is determined not to require EMS transport, personnel will give the phone number of the patient/caller to the Poison Center Specialist. The Specialist will initiate a minimum of one follow-up call to the patient/caller to determine the status of patient.
6. Minimal information that should be obtained from the patient for the state poison center includes:
 - Name and age of patient
 - Time of exposure
 - Signs and symptoms
 - Substance(s) involved
 - Any treatment given
7. Minimal information which should be provided to the state poison center for mass poisonings, including biochemical terrorism and HazMat, includes:
 - Substance(s) involved
 - Signs and symptoms
 - Time of exposure
 - Any treatment given



Air Transport

Policy:

For the purposes of this policy, air transport refers to rotary wing aircraft or helicopter (HEMS). HEMS should be considered whenever time-dependent conditions in patient care can be improved by decreasing transport time or by giving advanced care not commonly available from ground EMS services (i.e. blood products, advanced procedures, or advanced monitoring).

Purpose:

The purpose of this policy is to:

- Improve patient care in the prehospital setting by decreasing out of hospital time in time-dependent conditions.
- Allow for expedient transport in time-dependent conditions or mass casualty settings.
- Provide life-saving treatment such as blood products or advanced monitoring.
- Provide more timely access to interventional care in acute Stroke and ST-elevation myocardial infarction (STEMI) patients.
- Time-dependent conditions:
 - ST Elevation Myocardial Infarction (STEMI)
 - Stroke and Large Vessel Occlusion Stroke
 - Moderate to Severe traumatic conditions

Procedure:

There is no clear evidence that define strict criteria as to which patients may benefit or time consideration benefit when assessing the need for HEMS.

Patient transportation via ground EMS should not be delayed to wait for HEMS transportation. If the patient is packaged and ready for transport and HEMS is not on scene, or within a reasonable distance, transportation should be initiated by ground EMS.

Air transport should be considered if any of the following criteria apply:

- High priority patient with > 30 - 45 minute transport times.
- High priority patient with geographic hospital transport distance > 45 miles.
- Entrapped patients with > 20 minute estimated extrication time.
- Multiple casualty incident with red/ yellow tag patients.
- Multi-trauma or medical patient requiring life-saving treatment not available in prehospital environment (i.e., blood transfusion, invasive procedure, operative intervention).
- Time dependent medical conditions such as acute ST-elevation myocardial infarctions (STEMI) or acute Stroke that could benefit from the resources at a specialty center as per the EMS System's Stroke and STEMI Plans.

If a potential need for HEMS is anticipated, but not confirmed, HEMS can be placed on standby (*this significantly decreases flight time without the need for auto-launch*).

If scene conditions or patient situation improves after activation of HEMS, and later determined not to be necessary, ALS personnel or administrative personnel may cancel the request for HEMS.

Minimal Information which should be provided to HEMS include:

- Number and Age of patient(s)
- Weight of patient(s)
- Mechanism of injury or nature of illness.
- Potential hazards or HazMat involvement.



Standards Policy: Transport Policy Section

Safe Transport of Pediatric Patients

Policy:

Without special considerations children are at risk of injury when transported by EMS. EMS must provide appropriate stabilization and protection to pediatric patients during EMS transport.

Purpose:

To provide:

- Provide a safe method of transporting pediatric patients within an ambulance.
- Protect the EMS system and personnel from potential harm and liability associated with the transportation of pediatric patients.

Procedure:

1. Drive cautiously at safe speeds observing traffic laws.
2. Tightly secure all monitoring devices and other equipment.
3. Insure that all pediatric patients less than 40 lbs are restrained with an approved child restraint device secured appropriately to the stretcher or captains chair.
4. Insure that all EMS personnel use the available restraint systems during the transport.
5. Transport adults and children who are not patients, properly restrained, in an alternate passenger vehicle, whenever possible.
6. Do not allow parents, caregivers, or other passengers to be unrestrained during transport.
7. NEVER attempt to hold or allow the parents or caregivers to hold the patient during transport.



Transport

Policy:

All individuals served by the EMS system will be evaluated, treated, and furnished transportation (if indicated) in the most timely and appropriate manner for each individual situation.

Purpose:

To provide:

- Rapid emergency EMS transport when needed.
- Appropriate medical stabilization and treatment at the scene when necessary
- Protection of patients, EMS personnel, and citizens from undue risk when possible.

Procedure:

1. All trauma patients with significant mechanism or history for multiple system trauma will be transported as soon as possible. The scene time should be 10 minutes or less.
2. All acute Stroke and acute ST-Elevation Myocardial Infarction patients will be transported as soon as possible. The scene time should be 10 minutes or less for acute Stroke patients and 10 minutes or less (with 12 Lead ECG) for STEMI patients.
3. Other Medical patients will be transported in the most efficient manner possible considering the medical condition. Advanced life support therapy should be provided at the scene if it would positively impact patient care. Justification for scene times greater than 20 minutes should be documented.
4. No patients will be transported in initial response non-transport vehicles.
5. In unusual circumstances, transport in other vehicles may be appropriate when directed by EMS administration.



Standards Policy: Transport Policy Section

Optional: Weapons and Explosive Devices

Policy:

Pursuant to 10A NCAC 13P .0216; Weapons, whether lethal, less lethal or non-lethal, and explosives (concealed or visible) shall not be worn or carried aboard an ambulance or EMS non-transporting vehicle within the State of North Carolina when the vehicle is operating in any patient treatment or transport capacity or is available for such function.

Weapons authorized for use by EMS personnel attached to a law enforcement tactical team in accordance with the weapons policy as set forth in Rule .0201 of this Section may be secured in a locked, dedicated compartment or gun safe mounted within the ambulance or non-transporting vehicle for use when dispatched in support of the law enforcement tactical team, but are not to be worn or carried open or concealed by any EMS personnel in the performance of normal EMS duties under any circumstances. Rule .0201 requires every EMS System to have “a weapons plan for any weapon as set forth in Rule 10A NCAC 13P .0216 of this Section;” This NCCEP policy does not supersede local EMS system policy.

Purpose:

To ensure the safety of EMS personnel, patients, and the public at large.

Definitions:

Conducted electrical weapons and chemical irritants such as mace, pepper (oleoresin capsicum) spray, and tear gas shall be considered weapons for the purpose of this Rule.

- Weapon is defined as any device with a tube, including mechanical attachments, from which a projectile is delivered by force of an explosive and may be lethal or less lethal including conducted electrical device. Weapons may be either concealed or visible
- Chemicals considered weapons:
Mace or Pepper spray (OC - oleoresin capsicum, CS – orthochlorbenzalmalonitrile
CN – alphachloroacetaphenone)
- Projectile is typically represented by bullets, shells, or slugs and may be metallic or non-metallic in composition

Procedure:

- During scene size-up and during your secondary patient assessment all patients should be screened for weapons or explosive devices, preferably before entering an ambulance.
- Screen all patients regardless if they have been screened by Law Enforcement.
- Individuals riding in the ambulance (friends, family) should be asked about concealed weapons.
- In the event a weapon or explosive device is found in an ambulance during transportation, the weapon or explosive device will be safely secured or turned over to Law Enforcement.
- Ask all patients, including family/friends, transported if they have a concealed weapon.
- **“Do you have any firearms or weapons on your person”**
- Likely scenarios EMS may encounter with individual found carrying a weapon:
- Individual has capacity, or does not have capacity, but is cooperative
- **Individual does not have capacity and is not cooperative:**
- **In this situation EMS personnel should retreat immediately to safety and notify Law Enforcement to secure the scene**
- It is recommended that EMS systems have a lock-box of sufficient size to accommodate a typical hand-gun on each EMS vehicle or available to EMS crews to safely and securely stow and transport a weapon.
- This Rule shall not apply to duly appointed law enforcement officers.



Standards Policy: Transport Policy Section

Alert – Code Blood

Policy:

Code Blood should be utilized whenever patient care can be improved with the administration of blood products.

Purpose:

The purpose of this policy is to:

1. Improve patient care in the prehospital setting
2. Provide life-saving treatment such as blood transfusion
3. Rapid transport of whole blood to the crew requesting

Procedure:

Patient transport will not be delayed to wait for whole blood. If the patient is packaged and ready for transport and the QRV is not on scene or within a reasonable distance, the transportation will be initiated.

Low Titer O+ Whole Blood should be considered if any of the following criteria apply:

- BP < 70 (Adult)
- BP < 90 with heart rate > 110 (Adult)
- Witnessed traumatic arrest < 10 minutes prior to provider arrival and continuous CPR throughout downtime
- Age > 65: SBP < 100 AND heart rate > 100
- Age 0-28 days: SBP < 60
- Age ≥ 1 month: SBP < 70
- Age 1-9: SBP < 70 + (2 x age)
- Age ≥ 10: SBP < 90 (pediatrics)
- All ages: Shock Index Criteria > 1 (shock index = heart rate/systolic blood pressure)

If any of the above criteria are met, EMS unit will contact the on-duty supervisor via radio and advise of a **Code Blood**.

CODE BLOOD PATIENTS WILL BE TRANSPORTED TO THE CLOSEST TRAUMA CENTER.



Standards Procedure (Skill) Airway Section Airway: BIAD-i-Gel

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Clinical Indications for Blind Insertion Airway Device (BIAD) Use:

- Inability to adequately ventilate a patient with a Bag Valve Mask or longer EMS transport distances require a more advanced airway.
- Inability to secure an endotracheal tube in a patient who does not have a gag reflex.
- Appropriate intubation is impossible due to patient access or difficult airway anatomy.
- Do not leave in place for ≥ 4 hours.
- **This airway does not prevent aspiration of stomach contents.**

Clinical Contraindications:

- Deforming Facial Trauma
- Pulmonary Fibrosis
- Morbid Obesity

Procedure:

1. Pre-Oxygenate the patient with 100% Oxygen
2. Select the appropriate tube size for the patient.
3. Remove the device from the protective cradle and carefully check for any signs of damage.
4. Place water-soluble jelly in the middle of the protective cradle.
5. Lubricate the back of the i-Gel on the non-inflatable cuff and ensure no lubricant is in the cuff.
6. Lubricate each side and the tip of the non-inflatable cuff.
7. Grasp along the integral bite block and face the cuff outlet toward the patient's chin.
8. Insert the i-Gel into the mouth in the direction of the hard palate.
9. Glide the device down and back along the hard palate with continuous, gentle pressure, until a definitive resistance is felt.
10. Connect the i-Gel to an BVM and assess for breath sounds and air entry.
11. Confirm tube placement using esophageal bulb device.
12. Monitor oxygen saturation with pulse oximetry and heart rhythm with ECG.
13. **EtCO₂ monitoring is mandatory following placement of a BIAD once available on scene.**
14. Re-verify i-Gel placement after every move and upon arrival in the ED.
15. Document the procedure, time, and result (success) on/with the patient care report (PCR).
16. Complete an Airway Evaluation Form.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System. Assessment should include direct observation once per certification cycle.



Airway: Cricothyrotomy-Surgical

Clinical Indications:**P PARAMEDIC P**

- Failed Airway Protocol
- Management of an airway when standard airway procedures cannot be performed or have failed in a patient \geq 12 years old.

Procedure:

1. Have suction and supplies available and ready.
2. Locate the cricothyroid membrane utilizing anatomical landmarks.
3. Prep the area with an antiseptic swab (Betadine).
4. Attach a 5-cc syringe to an 18G - 1 & 1/2-inch needle.
5. Insert the needle (with syringe attached) perpendicularly through the cricothyroid membrane with the needle directed posteriorly.
6. During needle insertion, gentle aspiration should be applied to the syringe. Rapid aspiration of air into the syringe indicates successful entry into the trachea. Do not advance the needle any further. Attach forceps and remove syringe.
7. With the needle remaining in place, make a 1-inch vertical incision through the skin and subcutaneous tissue above and below the needle using a scalpel. Using blunt dissection technique, expose the cricothyroid membrane. This is a bloody procedure. The needle should act as a guide to the cricothyroid membrane.
8. With the needle still in place, make a horizontal stabbing incision approx. 1/2 inch through the membrane on each side of the needle. Remove the needle.
9. Using (skin hook, tracheal hook, or gloved finger) to maintain surgical opening, insert the cuffed tube into the trachea. (Cric tube from the kit or a #6 endotracheal tube is usually sufficient).
10. Inflate the cuff with 5-10cc of air and ventilate the patient while manually stabilizing the tube.
11. All of the standard assessment techniques for insuring tube placement should be performed (auscultation, chest rise & fall, end-tidal CO₂ detector, etc.) Esophageal bulb devices are not accurate with this procedure.
12. Secure the tube.
13. If Available apply end tidal carbon dioxide monitor (Capnography) and record readings on scene, en route to the hospital, and at the hospital.
14. Document ETT size, time, result (success), and placement location by the centimeter marks either on/with the patient care report (PCR). Document all devices used to confirm initial tube placement and after each movement of the patient.
15. Consider placing an NG or OG tube to clear stomach contents after the airway is secured.
16. **Monitor the airway continuously through Capnography and Pulse Oximetry.**
17. **An Airway Evaluation Form must be completed with all intubations.**

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Airway: Intubation Oral Tracheal

A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Inability to adequately ventilate a patient with a Bag Valve Mask or longer EMS transport distances require a more advanced airway.
- An unconscious patient without a gag reflex who is apneic or is demonstrating inadequate respiratory effort.
- A component of Drug Assisted Intubation

Procedure:

1. Prepare, position and oxygenate the patient with 100% Oxygen.
2. Select proper ET tube (and stylette, if used), have suction ready.
3. Using laryngoscope, visualize vocal cords. (Use Sellick maneuver/BURP to assist you).
4. Limit each intubation attempt to 30 seconds with BVM between attempts.
5. Visualize tube passing through vocal cords.
6. **Confirm and document tube placement using an end-tidal CO₂ monitoring or esophageal bulb device.**
7. Inflate the cuff with 3-to10 cc of air; secure the tube to the patient's face.
8. Auscultate for bilaterally equal breath sounds and absence of sounds over the epigastrium. If you are unsure of placement, remove tube and ventilate patient with bag-valve mask.
9. Consider using a Blind Insertion Airway Device if intubation efforts are unsuccessful.
10. If Available apply end tidal carbon dioxide monitor (Capnography) and record readings on scene, en route to the hospital, and at the hospital.
11. Document ETT size, time, result (success), and placement location by the centimeter marks either at the patient's teeth or lips on/with the patient care report (PCR). Document all devices used to confirm initial tube placement. Also document positive or negative breath sounds before and after each movement of the patient.
12. Consider placing an NG or OG tube to clear stomach contents after the airway is secured with an ET tube.
13. **End-tidal (EtCO₂) monitoring is mandatory following placement of an endotracheal tube.**
14. **An Airway Evaluation Form must be completed with all intubations and attached to the ePCR.**

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System. Assessment should include direct observation at least once per certification cycle.



Standards Procedure (Skill) Airway Section

Airway: Intubation Nasotracheal

A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- A spontaneously breathing patient in need of intubation (inadequate respiratory effort, evidence of hypoxia or carbon dioxide retention, or need for airway protection).
- Rigidity or clenched teeth prohibiting other airway procedures.
- Patient must be 12 years of age or older.

Procedure:

1. Premedicate the patient with nasal spray.
2. Select the largest and least obstructed nostril and insert a lubricated nasal airway to help dilate the nasal passage.
3. Preoxygenate the patient. Lubricate the tube. The use of a BAAM device is recommended.
4. Remove the nasal airway and gently insert the tube keeping the bevel of the tube toward the septum.
5. Continue to pass the tube listening for air movement and looking for to and from vapor condensation in the tube. As the tube approaches the larynx, the air movement gets louder.
6. Gently and evenly advance the tube through the glottic opening on the inspiration. This facilitates passage of the tube and reduces the incidence of trauma to the vocal cords.
7. Upon entering the trachea, the tube may cause the patient to cough, buck, strain, or gag. Do not remove the tube! This is normal, but be prepared to control the cervical spine and the patient, and be alert for vomiting.
8. Auscultate for bilaterally equal breath sounds and absence of sounds of the epigastrium. Observe for symmetrical chest expansion. The 15mm adapter usually rests close to the nostril with proper positioning.
9. Inflate the cuff with 5-10 cc of air.
10. **Confirm tube placement using an end-tidal CO₂ monitoring or esophageal bulb device.**
11. Secure the tube.
12. Reassess airway and breath sounds after transfer to the stretcher and during transport. These tubes are easily dislodged and require close monitoring and frequent reassessment.
13. Document the procedure, time, and result (success) on/with the patient care report (PCR).
14. **End-tidal (EtCO₂) monitoring is mandatory following placement of an endotracheal tube.**
15. **An Airway Evaluation Form must be completed with all intubations and attached to the ePCR.**

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Standards Procedure (Skill) Airway Section

Airway: Video Laryngoscopy McGrath

Clinical Indications:

- Patient requires advanced airway.



A	AEMT	A
P	PARAMEDIC	P

Procedure:

1. Preoxygenate the patient and use in conjunction with procedure ASP - 6. Position the patient in the optimal position for direct laryngoscopy.
2. Select the appropriate ETT size and disposable laryngoscope blade for the patient. Ready suction.
3. Power on the McGrath by a single push of the Power button.
4. Insert the blade into the right side of the mouth, moving the device to the central position while sweeping the tongue to the left. Visualize the vocal cords and facilitate the intubation.
5. Advance the blade into the vallecula.
6. Visualize the epiglottis on the screen. Lift the anatomy forward and upwards to expose a direct and indirect view of the glottis.
7. When the device is in the optimal position the glottis should be viewed in the central upper section of the screen.
8. Advance the tube gently and atraumatically through the vocal cords. Tube placement can be performed either by looking directly in the mouth, indirectly on the screen or a combination of both.
9. If a direct pathway for the tube was not created by sweeping the tongue or aligning the airway axes, a bougie may need to be used.
10. Indirectly visualize the tube placement through the vocal cords.
11. The screen view can be used to confirm the correct insertion depth of the endotracheal tube.
12. Auscultate for breath sounds and sounds over the epigastrium and look for the chest to rise and fall.
13. Secure the ETT tube with tape or mechanical tube holder.
14. Confirm tube placement using end-tidal CO₂ monitor.
15. End-tidal (EtCO₂) monitoring is mandatory following placement of an endotracheal tube.
16. Complete the Airway Evaluation Form.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System. Assessment should include direct observation at least once per certification cycle.



North Carolina College of Emergency Physicians Standards Procedure (Skill)

Airway: Drug Assisted Airway

Clinical Indications:

P PARAMEDIC **P**

- Need for advanced airway control in a patient who has a gag reflex or trismus (jaw clenching)
- Failure to protect the airway. Unable to ventilate and / or oxygenate. Impending airway compromise
- A minimum of 2 EMT-Paramedic on scene able to participate in patient care.
- This protocol is only for use in patients longer than a Length-based Resuscitation Tape. Patient MUST be >15 years old OR >49 kg.

Clinical Contraindications:

- Refer to drug list for contraindications regarding use of Succinylcholine and Rocuronium.

Procedure:

1. Perform focused neurological exam
2. Evaluate for difficult airway (LEMON)-see appendix
3. Prepare equipment (intubation kit, BVM, suction, DAI medications, BIAD, Cricothyrotomy kit, waveform capnography, other airway adjuncts as available)
4. Pre-oxygenate patient with 100% oxygen via NRB mask or BVM. Apneic oxygenation: May continue high-flow oxygen via NC during entire procedure
5. Monitor oxygen saturation with pulse oximetry and heart rhythm with ECG
6. Ensure functioning IV / IO access. Two (2) IV sites are preferable
7. In-line c-spine stabilization by second caregiver (in setting of trauma)
8. Administer Etomidate or Ketamine by rapid IV push
9. Administer Succinylcholine or Rocuronium, await fasciculation and jaw relaxation
10. Perform external laryngeal manipulation to improve view during laryngoscopy with the right hand.
11. Intubate trachea or place BIAD if intubation unsuccessful or felt to be unsuccessful during procedure.
12. Verify ET placement through auscultation, Capnography, and Pulse Oximetry
13. May repeat Succinylcholine or Rocuronium if inadequate relaxation
14. Release cricoid pressure (if utilized) and secure tube
15. **Continuous Capnography and Pulse Oximetry is required for DAI. Pre-intubation, minimal during intubation, and post-intubation readings must be recorded in the PCR.**
16. Re-verify tube placement after every move and upon arrival in the ED
17. Document ETT or BIAD size, time, result (success), and placement location by the centimeter marks either at the patient's teeth or lips on/with the patient care report (PCR). Document all devices/methods used to confirm initial tube placement initially and with patient movement.
18. Insert a gastric tube to clear stomach contents after the airway is secured.
19. **Completion of the Airway Evaluation Form is required including a signature from the receiving physician at the Emergency Department confirming proper tube placement. Attach a copy to the ePCR, The original document MUST be submitted to the QA Manager.**

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System. Assessment should include direct observation at least once per certification cycle.



Airway: Drug Assisted Airway Intubation "TIMEOUT" Checklist

P **PARAMEDIC** **P**

Clinical Indications:

- Age >15 years old OR Weight >49 kg
- This checklist should be performed with any patient identified to need advanced airway control who has a gag reflex.

Procedure:

Report current vitals **HR:_____ RR:_____ B/P_____ SPO2:_____ BGL_____**

***Resuscitation and correction of hypoxia and /or hypotension (< 90 systolic) are paramount prior to use of RSI medications.**

Challenge

1. Nasal Cannula (not EtCo2)
2. Medication to be given
3. Suction
4. BVM
5. Blade and Backup
6. ETT x2 w/ stylette
7. Bougie
8. NG Tube
9. C-collar / Tube Stabilizer
10. Alternative Airway
11. McGrath
12. Capnography (ET)
13. Roles Clear
 - a. Intubation
 - b. Medications
 - c. Monitor
14. SpO2
15. State Current Vitals

Response

- On @10 lpm
Assigned & Reviewed
On & In Reach
Connected
Near Head
Both Tested
Placed Near Head
Prepared
Under Pt's Neck
I-Gel/ OPA/NPA
State Battery Time
Prepared
Verbal "Yes" From ALL
- State Current %
HR__RR__BP__SPO2__BGL__

*** Keep Oxygen Saturation >90%**

*** Preferably >94%**

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System. Assessment should include direct observation at least once per certification cycle.



Airway: Tracheostomy Tube Change

A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Presence of Tracheostomy site.
- Urgent or emergent indication to change the tube, such as obstruction that will not clear with suction, dislodgement, or inability to oxygenate/ventilate the patient without other obvious explanation.

Procedure:

1. Have all airway equipment prepared for standard airway management, including equipment of orotracheal intubation and failed airway.
2. Have airway device (endotracheal tube or tracheostomy tube) of the same size as the tracheostomy tube currently in place as well as 0.5 size smaller available (e.g., if the patient has a #6.0 Shilley, then have a 6.0 and a 5.5 tube).
3. Lubricate the replacement tube(s) and check the cuff.
4. Remove the tracheostomy tube from mechanical ventilation devices and use a bag-valve apparatus to pre-oxygenate the patient as much as possible.
5. Once all equipment is in place, remove devices securing the tracheostomy tube, including sutures and/or supporting bandages.
6. If applicable, deflate the cuff on the tube. If unable to aspirate air with a syringe, cut the balloon off to allow the cuff to lose pressure.
7. Remove the tracheostomy tube.
8. Insert the replacement tube. Confirm placement via standard measures except for esophageal detection (which is ineffective for surgical airways).
9. If there is any difficulty placing the tube, re-attempt procedure with the smaller tube.
10. If difficulty is still encountered, use standard airway procedures such as oral bag-valve mask or endotracheal intubation (as per protocol). **More difficulty with tube changing can be anticipated for tracheostomy sites that are immature – i.e., less than two weeks old. Great caution should be exercised in attempts to change immature tracheostomy sites.**
11. Document procedure, confirmation, patient response, and any complications in the PCR

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System. Assessment for this skill should include direct observation at least once per certification cycle.



Airway: Endotracheal Tube Introducer (Bougie)

Clinical Indications:

- Patients meet clinical indications for oral intubation
- Initial intubation attempt(s) unsuccessful
- Predicted difficult intubation

A	AEMT	A
P	PARAMEDIC	P

Contraindications:

- Three attempts at orotracheal intubation (utilize failed airway protocol)
- Age less than eight (8) or ETT size less than 6.5 mm

Procedure:

1. Prepare, position and oxygenate the patient with 100% oxygen;
2. Select proper ET tube without stylet, test cuff and prepare suction;
3. Lubricate the distal end and cuff of the endotracheal tube (ETT) and the distal 1/2 of the Endotracheal Tube Introducer (Bougie) (note: Failure to lubricate the Bougie and the ETT may result in being unable to pass the ETT);
4. Using laryngoscopic techniques, visualize the vocal cords if possible using Sellick's/BURP as needed;
5. Introduce the Bougie with curved tip anteriorly and visualize the tip passing the vocal cords or above the arytenoids if the cords cannot be visualized;
6. Once inserted, gently advance the Bougie until you meet resistance or "hold-up" (if you do not meet resistance you have a probable esophageal intubation and insertion should be re-attempted or the failed airway protocol implemented as indicated);
7. Withdraw the Bougie ONLY to a depth sufficient to allow loading of the ETT while maintaining proximal control of the Bougie;
8. Gently advance the Bougie and loaded ET tube until you have hold-up again, thereby assuring tracheal placement and minimizing the risk of accidental displacement of the Bougie;
9. While maintaining a firm grasp on the proximal Bougie, introduce the ET tube over the Bougie passing the tube to its appropriate depth;
10. If you are unable to advance the ETT into the trachea and the Bougie and ETT are adequately lubricated, withdraw the ETT slightly and rotate the ETT 90 degrees COUNTER clockwise to turn the bevel of the ETT posteriorly. If this technique fails to facilitate passing of the ETT you may attempt direct laryngoscopy while advancing the ETT (this will require an assistant to maintain the position of the Bougie and, if so desired, advance the ETT);
11. Once the ETT is correctly placed, hold the ET tube securely and remove the Bougie;
12. Confirm tracheal placement according to the intubation protocol, inflate the cuff with 3 to 10 cc of air, auscultate for equal breath sounds and reposition accordingly;
13. When final position is determined secure the ET tube, reassess breath sounds, apply end tidal CO₂ monitor, and record and monitor readings to assure continued tracheal intubation.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System. Assessment should include direct observation at least once per certification cycle.



Airway: Foreign Body Obstruction

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Sudden onset of respiratory distress often with coughing, wheezing, gagging, or stridor due to a foreign-body obstruction of the upper airway.

Procedure:

1. Assess the degree of foreign body obstruction
 - Do not interfere with a mild obstruction allowing the patient to clear their airway by coughing.
 - In severe foreign-body obstructions, the patient may not be able to make a sound. The victim may clutch his/her neck in the universal choking sign.
2. **For an infant**, deliver 5 back blows (slaps) followed by 5 chest thrusts repeatedly until the object is expelled or the victim becomes unresponsive.
3. **For a child**, perform a subdiaphragmatic abdominal thrust (Heimlich Maneuver) until the object is expelled or the victim becomes unresponsive.
4. **For adults**, a combination of maneuvers may be required.
 - First, subdiaphragmatic abdominal thrusts (Heimlich Maneuver) should be used in rapid sequence until the obstruction is relieved.
 - If abdominal thrusts are ineffective, chest thrusts should be used. Chest thrusts should be used primarily in morbidly obese patients and in the patients who are in the late stages of pregnancy
5. If the victim becomes unresponsive, begin CPR immediately but look in the mouth before administering any ventilations. If a foreign-body is visible, remove it.
6. **Do not perform blind finger sweeps in the mouth and posterior pharynx. This may push the object farther into the airway.**
7. In unresponsive patients, AEMT and Paramedic level professionals should visualize the posterior pharynx with a laryngoscope to potentially identify and remove the foreign-body using Magill forceps.
8. Document the methods used and result of these procedures in the patient care report (PCR).

Certification Requirements:

Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.

Airway: Post-Endotracheal Intubation Checklist

A	AEMT	A
P	EMT- P	P

Clinical Indication:

- Immediately after Endotracheal Intubation Procedure is performed.

Procedure:

1. Ensure cuff of endotracheal tube is inflated per Endotracheal Intubation procedure.
2. Monitor continuous (EtCO₂) waveform capnography to ensure that waveform is consistent with correct airway placement.
3. If no waveform capnography (ETCO₂) is observed on the monitor, **IMMEDIATELY** remove endotracheal tube from airway and provide Bag Valve Mask ventilations with appropriate oxygen source connected and flowing.
 - Escalate per protocol and as needed with airway adjuncts and BIAD. If BIAD is utilized as a rescue device, immediately confirm placement with ETCO₂ waveform capnography.
4. If cricothyrotomy procedure is being considered, follow protocol for procedure as indicated. After placement confirm that ETCO₂ waveform is present.
5. Provide the patient with Bag Valve ventilations and ensure appropriate oxygen source is flowing.
6. Listen to the lungs for equal breath sounds in the upper and lower lung fields, bilaterally; also, check for the absence of sounds over the epigastrium / center of the upper abdomen.
7. Continue appropriate ventilation by bag ventilation or confirm clinically appropriate ventilator settings on the ventilator. Re-confirm Bag Valve ventilations or that ventilator is connected to appropriate oxygen source that is flowing.
8. Secure the airway device with a commercially available product. If ETT, document placement location by the centimeter mark either at the patient's teeth or lips on the patient care report (PCR).
9. If using a mechanical ventilator follow RSP 6 Procedure: Ventilator Operation
10. Document all devices used to confirm initial tube placement. Also, document presence or absence of breath sounds before and after each movement of the patient.
11. Re-check and document vital signs immediately after intubation and at least every 5 minutes.
12. Follow post-intubation protocol for analgesia, sedation and neuro-muscular blockade as indicated.
13. It is strongly recommended that an airway evaluation form be completed with all intubations.



Airway: Post Intubation Checklist

Procedure (continued):

14. Record ETCO₂ monitor readings on scene, en-route to hospital, and at the hospital after patient is moved to a hospital stretcher.
15. Verify tube placement frequently and always with each movement of patient.
16. Consider placing an NG or OG tube to clear stomach contents after airway device is secured with ET tube or Supraglottic airway with a gastric tube port.
17. Assess for worsening abdominal distention with every delivered breath, or concern of aspiration of gastric contents.

Certification Requirements:

Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS system.



Assessment: Adult

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Any patient requesting a medical evaluation that is too large to be measured with a Length-based **2017** Resuscitation Tape.

Procedure:

- Scene size-up, including universal precautions, scene safety, environmental hazards assessment, need for additional resources, by-stander safety, and patient/caregiver interaction
- Assess need for additional resources.
- Initial assessment includes a general impression as well as the status of a patient’s airway, breathing, and circulation.
- Assess mental status (e.g., AVPU) and disability (e.g., GCS).
- Control major hemorrhage and assess overall priority of patient.
- Perform a focused history and physical based on patient’s chief complaint.
- Assess need for critical interventions.
- Complete critical interventions and perform a complete secondary exam to include a baseline set of vital signs as directed by protocol.
- Maintain an on-going assessment throughout transport; to include patient response/possible complications of interventions, need for additional interventions, and assessment of evolving patient complaints/conditions.
- Document all findings and information associated with the assessment, performed procedures, and any administration of medications on the PCR.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Standards Procedure (Skill) Assessment / Screening Section Pain Assessment and Documentation

Clinical Indications:

- Any patient with pain.


Definitions:

- Pain is an unpleasant sensory and emotional experience associated with actual or potential tissue damage.
- Pain is subjective (whatever the patient says it is).

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Procedure:

- Initial and ongoing assessment of pain intensity and character is accomplished through the patient's self report.
- Pain should be assessed and documented in the PCR during initial assessment, before starting pain control treatment, and with each set of vitals.
- Pain should be assessed using the appropriate approved scale.
- Three pain scales are available: the 0 – 10, the Wong - Baker "faces", and the FLACC.
 - 0 – 10 Scale: the most familiar scale used by EMS for rating pain with patients. It is primarily for adults and is based on the patient being able to express their perception of the pain as related to numbers. Avoid coaching the patient; simply ask them to rate their pain on a scale from 0 to 10, where 0 is no pain at all and 10 is the worst pain ever.
 - Wong – Baker "FACES" scale: this scale is primarily for use with pediatrics but may also be used with geriatrics or any patient with a language barrier. The faces correspond to numeric values from 0-10. This scale can be documented with the numeric value.



From Hockenberry MJ, Wilson D, Winkelstein ML: Wong's Essentials of Pediatric Nursing, ed. 7, St. Louis, 2005, p. 1259. Used with permission. Copyright, Mosby.

- FLACC scale: this scale has been validated for measuring pain in children with mild to severe cognitive impairment and in pre-verbal children (including infants).

CATEGORIES	SCORING		
	0	1	2
FACE	No particular expression or smile	Occasional grimace or frown, withdrawn, disinterested.	Frequent to constant quivering chin, clenched jaw.
LEGS	Normal position or relaxed.	Uneasy, restless, tense.	Kicking, or legs drawn up.
ACTIVITY	Lying quietly, normal position moves easily.	Squirming, shifting back and forth, tense.	Arched, rigid or jerking.
CRY	No cry, (awake or asleep)	Moans or whimpers; occasional complaint	Crying steadily, screams or sobs, frequent complaints.
CONSOLABILITY	Content, relaxed.	Reassured by occasional touching hugging or being talked to, distractable.	Difficulty to console or comfort

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Assessment: Pediatric

Clinical Indications:

- Any child that can be measured with a Length-based Resuscitation Tape.

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Procedure:

- Scene size-up, including universal precautions, scene safety, environmental hazards assessment, need for additional resources, by-stander safety, and patient/caregiver interaction
- Assess patient using the pediatric triangle of ABCs:
 - Airway and appearance: speech/cry, muscle tone, inter-activeness, look/gaze, movement of extremities
 - Work of breathing: absent or abnormal airway sounds, use of accessory muscles, nasal flaring, body positioning
 - Circulation to skin: pallor, mottling, cyanosis
- Establish spinal immobilization if suspicion of spinal injury
- Establish responsiveness appropriate for age (AVPU, GCS, etc.)
- Color code using Broselow-Luten tape
- Assess disability (pulse, motor function, sensory function, papillary reaction)
- Perform a focused history and physical exam. Recall that pediatric patients easily experience hypothermia and thus should not be left uncovered any longer than necessary to perform an exam.
- Record vital signs (BP > 3 years of age, cap refill < 3 years of age)
- Include Immunizations, Allergies, Medications, Past Medical History, last meal, and events leading up to injury or illness where appropriate.
- Treat chief complaint as per protocol

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Blood Glucose Analysis

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Patients with suspected hypoglycemia (diabetic emergencies, change in mental status, bizarre behavior, etc.)

Procedure:

1. Gather and prepare equipment.
2. Blood samples for performing glucose analysis can be obtained through a finger-stick or when possible simultaneously with intravenous access.
3. Place correct amount of blood on reagent strip or site on glucometer per the manufacturer's instructions.
4. Time the analysis as instructed by the manufacturer.
5. Document the glucometer reading and treat the patient as indicated by the analysis and protocol.
6. Repeat glucose analysis as indicated for reassessment after treatment and as per protocol.
7. Perform Quality Assurance per manufacture recommendation.

Consideration:

- Brunswick County EMS uses the Assure Prism Multi blood glucose mointor. If the blood sugar result shows a reading of "Lo" the blood sugar is less than 20 mg/dL. If the blood sugar result shows a reading of "Hi" the blood sugar is greater than 600 mg/dL.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Capnography

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Capnography shall be used with the use of all invasive airway procedures including endotracheal, nasotracheal, cricothyrotomy, or Blind Insertion Airway Devices (BIAD).
- Capnography should also be used when possible with CPAP.

Procedure:

1. Attach capnography sensor to the BIAD, endotracheal tube, or oxygen delivery device.
2. Note CO₂ level and waveform changes. These will be documented on each respiratory failure, cardiac arrest, or respiratory distress patient.
3. The capnometer shall remain in place with the airway and be monitored throughout the prehospital care and transport.
4. Any loss of CO₂ detection or waveform indicates an airway problem and should be documented.
5. The capnogram should be monitored as procedures are performed to verify or correct the airway problem.
6. Document the procedure and results on/with the Patient Care Report (PCR) and the Airway Evaluation Form.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Pulse Oximetry

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Patients with suspected hypoxemia.

Procedure:

1. Apply probe to patient's finger or any other digit as recommended by the device manufacturer.
2. Allow machine to register saturation level.
3. Record time and initial saturation percent on room air if possible on/with the patient care report (PCR).
4. Verify pulse rate on machine with actual pulse of the patient.
5. Monitor critical patients continuously until arrival at the hospital. If recording a one-time reading, monitor patients for a few minutes as oxygen saturation can vary.
6. Document percent of oxygen saturation every time vital signs are recorded and in response to therapy to correct hypoxemia.
7. In general, normal saturation is 97-99%. Below 94%, suspect a respiratory compromise.
8. Use the pulse oximetry as an added tool for patient evaluation. Treat the patient, not the data provided by the device.
9. The pulse oximeter reading should never be used to withhold oxygen from a patient in respiratory distress or when it is the standard of care to apply oxygen despite good pulse oximetry readings, such as chest pain. Supplemental oxygen is not required if the oxyhemoglobin saturation is $\geq 94\%$, unless there are obvious signs of heart failure, dyspneic, or hypoxic to maintain to 94%.
10. Factors which may reduce the reliability of the pulse oximetry reading include but are not limited to:
 - Poor peripheral circulation (blood volume, hypotension, hypothermia)
 - Excessive pulse oximeter sensor motion
 - Fingernail polish (may be removed with acetone pad)
 - Carbon monoxide bound to hemoglobin
 - Irregular heart rhythms (atrial fibrillation, SVT, etc.)
 - Jaundice
 - Placement of BP cuff on same extremity as pulse ox probe.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Reperfusion Checklist

Clinical Indications:

Rapid evaluation of a patient with suspected acute stroke and/or acute myocardial infarction (STEMI) to:

- Determine eligibility and potential benefit from fibrinolysis..
- Rapid identification of patients who are not eligible for fibrinolysis and will require interventional therapy.

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Procedure:

1. Follow the appropriate protocol for the patient’s complaint to assess and identify an acute condition which could potentially benefit from fibrinolysis. If a positive finding is noted on one of the following assessments, proceed to step 2.
 - Perform a 12-lead ECG to identify an acute ST elevation myocardial infarction (STEMI).
 - Perform the Los Angeles Pre-hospital Stroke Screen to identify an acute stroke
2. Complete the Reperfusion Check Sheet to identify any potential contraindications to fibrinolysis. (See Appendix)
 - Systolic Blood Pressure greater than 180 mm Hg
 - Diastolic Blood Pressure greater than 110 mm Hg
 - Right vs. Left Arm Systolic Blood Pressure difference of greater than 15 mm Hg
 - History of structural Central Nervous System disease (age >= 18, history of aneurysm or AV-malformation, tumors, masses, hemorrhage, etc.)
 - Significant closed head or facial trauma within the previous 3 months
 - Recent (within 6 weeks) major trauma, surgery (including laser eye surgery), gastrointestinal bleeding, or severe genital-urinary bleeding
 - Bleeding or clotting problem or on blood thinners
 - CPR performed greater than 10 minutes
 - Currently Pregnant
 - Serious Systemic Disease such as advanced/terminal cancer or severe liver or kidney failure.
3. Identify if the patient is currently in heart failure or cardiogenic shock. For these patients, a percutaneous coronary intervention is more effective.
 - Presence of pulmonary edema (rales greater than halfway up lung fields)
 - Systemic hypoperfusion (cool and clammy)
4. If any contraindication is noted using the check list and an acute Stroke is suspected by exam or a STEMI is confirmed by ECG, activate the EMS Stroke Plan or EMS STEMI Plan for fibrinolytic ineligible patients. This may require the EMS Agency, an Air Medical Service, or a Specialty Care Transport Service to transport directly to a specialty center capable of interventional care within the therapeutic window of time.
5. Record all findings in the Patient Care Report (PCR).

Certification Requirements:

Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Stroke Screen: MEND Exam and RACE

Clinical Indications:

- Suspected Stroke Patient
- Rapid Arterial Occlusion Evaluation Scale (RACE)
-

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Procedure:

1. Assess and treat suspected stroke patients as per protocol.
2. Miami Emergency Neurological Deficit Exam (MEND), this is a rapid screening tool and can detect both anterior and posterior circulation strokes. This is a positive or negative test.
3. Rapid Arterial Occlusion Evaluation Scale (RACE), this tool is based on the items of the NIHSS with the highest predictive value for large vessel occlusion (LVO). This RACE scale has a score range of 0-9; ≥ 4 points is associated with detection of a LVO.
4. The acute stroke activation criteria and assessment form should be completed for all suspected stroke patients (see appendix).
5. Screen the patient for the following criteria:
 - Level of consciousness
 - Speech
 - Questions
 - Facial Palsy
 - Visual Fields
 - Horizontal Gaze
 - Arm motor function
 - Leg motor function
 - Sensory
 - Coordination
 - Aphasia (right side) - loss of ability to understand or express speech
 - Agnosia (left side) - inability to interpret sensations and hence to recognize things
6. All sections of the form must be completed.
7. The completed form should be attached to the ePCR. Leave a copy at the receiving facility.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Temperature Measurement cont.

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Normal body temperature is a range. the Table below shows that normal temperatures vary by site. Therefore readings from different sites, even if taken at the same time, should not be directly compared.

The Welch Allyn Sure Temp Plus Thermometer will accurately predict the temperature of the site where the probe is placed. Be sure that the correct Mode is selected on the thermometer. There is no need to add / subtract a degree. Report the temperature along with the site used.

NORMAL BODY TEMPERATURE RANGES

°F	0 - 2 years	3 - 10 years	11 - 65 years	> 65 years
Oral	—	95.9 - 99.5	97.6 - 99.6	96.4 - 98.5
Rectal	97.9 - 100.4	97.9 - 100.4	98.6 - 100.6	97.1 - 99.2
Axillary	94.5 - 99.1	96.6 - 98.0	95.3 - 98.4	96.0 - 97.4
Ear	97.5 - 100.4	97.0 - 100.0	96.6 - 99.7	96.4 - 99.5
Core	97.5 - 100.0	97.5 - 100.0	98.2 - 100.2	96.6 - 98.8

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Temperature Measurement

Clinical Indications:

- Monitoring body temperature in a patient with suspected infection, hypothermia, hyperthermia, or to assist in evaluating resuscitation efforts.

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P



Procedure:

- For adult patients that are conscious, cooperative, and in no respiratory distress, an oral temperature is preferred (steps 2 to 5 below). For infants or adults that do not meet the criteria above, an axillary temperature is preferred (steps 6 to 8 below).
- To obtain an oral temperature, ensure the patient has no significant oral trauma. Observe the Oral Mode indicator on the display (flashing oral icon). 

If this icon is not flashing, press the Mode Selection button  until the icon appears.
- Load a Welch Allyn disposable probe cover.
- Have the patient seal their mouth closed around thermometer.
- Place the probe tip deep into the patient's sublingual pocket as shown below. Do not place the probe tip to the patient to place in their mouth. Always hold the probe in place, maintaining tissue contact until the temperature has been captured. Remove the probe and eject the probe cover.

Proceed to step 9.
- To obtain a pediatric axillary temperature, press Mode Selection button  to select the Pediatric Axillary Mode. Observe the Pediatric Axillary Mode indicator on the display (flashing pediatric axillary icon)  To obtain an adult axillary temperature, press mode selection button to select Adult Axillary Mode. 
- Load a Welch Allyn probe disposable cover, adjust clothing to visualize the axilla. Avoid folds in the axilla and place the probe tip vertically as high as you can, as shown below.
- Place the arm snugly at the patient's side. Hold the arm in this position without movement of the arm or probe during the measurement cycle. Remove the probe and eject the probe cover.
- Record time, temperature, method (oral, rectal), and scale (C° or F°) in Patient Care Report (PCR).

sublingual pockets





Standards Procedure (Skill) Assessment / Screening Section

Orthostatic Blood Pressure Measurement

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Patient situations with suspected blood, fluid loss, or dehydration with no indication for spinal immobilization. Orthostatic vital signs are not routinely recommended.
- Patients \geq 8 years of age, or patients larger than the Handtevy tape
- Orthostatic Vital Signs are not sensitive nor specific for volume loss / dehydration and may induce syncope in some cases. Assessment of orthostatic vital signs are not routinely recommended. Local Medical Director should indicate and educate on situations where they may be helpful.

Procedure:

1. Gather and prepare standard sphygmomanometer and stethoscope.
2. With the patient supine, obtain pulse and blood pressure.
3. Have the patient sit upright.
4. After 30 seconds, obtain blood pressure and pulse.
5. If the systolic blood pressure falls more than 30 mmHg or the pulse rises more than 20 bpm, the patient is considered to be orthostatic.
6. If a patient experiences dizziness upon sitting or is obviously dehydrated based on history or physical exam, formal orthostatic examination should be omitted and fluid resuscitation initiated.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Verbal De-escalation

Agency Name:		SATISFACTORY	<input type="checkbox"/>
Provider Name:	Paramedic	UNSATISFACTORY	<input type="checkbox"/>
Instructor Name:	EMT AEMT Paramedic Physician		

Instructor:
 1. Evaluate providers skill performance using the check off list below.
 2. Circle performance indicator:

		EMR	
YES = Provider completed skill with no assistance from instructor.	B	EMT	B
NO = Provider unable to complete skill satisfactorily following instructor intervention.	A	AEMT	A
IL = Provider able to complete skill satisfactorily following Instructor Led (teaching) intervention.	P	Paramedic	P

Satisfactory performance indicated with ≥ 8 YES / IL completions. (Combination of both YES and IL)

YES	NO	IL	Verbalizes indications for Verbal de-escalation techniques: 1. Behavioral Health Crisis 2. Behavior Activity Rating Score ≥ 5
-----	----	----	--

YES	NO	IL	Verbalizes contraindications: None
-----	----	----	------------------------------------

YES	NO	IL	Demonstrates respect of patient's personal space <ul style="list-style-type: none"> Maintain about 6 feet of distance and do not position yourself between the patient and only exit Both you and patient should be able to exit the room without feeling blocked-in
-----	----	----	---

YES	NO	IL	Does not provoke patient during interaction <ul style="list-style-type: none"> Your body language must convey that you want to listen and that you do not want to inflict harm <ul style="list-style-type: none"> Your hands should be visible and open Do not face the patient head-on. Always stand at an angle Avoid prolonged staring or direct eye contact Make sure others are not provoking the patient (providers, family members, bystanders, providers, police officers)
-----	----	----	---

YES	NO	IL	Establishes rapport, initiates and maintains verbal contact <ul style="list-style-type: none"> One person should make and maintain verbal contact, introduce yourself and explain your role <ul style="list-style-type: none"> Multiple providers talking to the patient will create confusion and may escalate patient's behavior Emphasize you are there to keep the patient safe Ask the patient their name and how they want to be addressed
-----	----	----	--

YES	NO	IL	Use concise statements when talking <ul style="list-style-type: none"> Agitation creates problems in a patient's ability to process information Keep your conversation simple and short in nature allowing time for patient to process information Repeat your statements, requests, or commands to ensure understanding <ul style="list-style-type: none"> This is called a loop, you may need to repeat 2 – 12 times before patient understands
-----	----	----	---

YES	NO	IL	Identify wants, feelings, and stress causing the crisis <ul style="list-style-type: none"> "When you called 911, how did you think we could help you?" "I would like to know what caused you to become upset today so we can help you" Identifying a need can help to quickly de-escalate the situation
-----	----	----	---

YES	NO	IL	Listen closely to patient <ul style="list-style-type: none"> You should be able to repeat back what is said by the patient <ul style="list-style-type: none"> "Tell me if I have all this right" "Let me make sure I understand what you said"
-----	----	----	---

YES	NO	IL	Agree or agree to disagree <ul style="list-style-type: none"> If statements are truthful, then agree with the truth Agree in principle, maybe patient's statement is not true, but you can agree, that in general, the idea is true Agree with the odds, anyone may be upset by the same circumstances Do not agree with delusions, at that point you can agree to disagree
-----	----	----	--

YES	NO	IL	Set clear limits on acceptable behavior <ul style="list-style-type: none"> Set limits in a positive, matter-of-fact manner, and not in a threatening manner Inform the patient that harm to self or other providers will not be tolerated If the patient's behavior is frightening to providers, tell the patient so Remind the patient you are there to help, keep them safe, but the providers cannot be abused in the process
-----	----	----	---

YES	NO	IL	Offer choices to patient (if available) and remain positive in your interactions <ul style="list-style-type: none"> Offer choices that are realistic and that may provide comfort such as drinks, food, blankets, etc. If medication is needed, offer choice between PO and IM/IV, offer medication early in encounter
-----	----	----	---

YES	NO	IL	Debrief provider team following the incident (if restraints necessary, debrief patient as well) What went well? What could have gone better? What did we learn? Who needs to know?
-----	----	----	---

Instructor notes:



Verbal De-escalation

Clinical Information for Verbal De-escalation

Objective of Procedure:

- Verbal engagement with patient and establishing collaborative relationship with patient
- Preventing violent behavior
- Avoiding use of restraints
- Reducing patient anger and frustration
- Maintaining staff and patient safety
- Enabling patients to manage their emotions and regain personal control

Scope of Practice: EMR EMT AEMT Paramedic

Indications:

1. Behavioral Health Crisis
2. Behavior Activity Rating Score ≥ 5

Contraindications:

None

Clinical Presentation:

Patient experiencing a behavioral crisis defined as:

- Significantly deviates from society’s expectations and commonly held normal behavior
 - Behavior that is unusual for patient’s baseline
 - Bizarre
 - Threatening
 - Dangerous to self and/or others
 - Alarming to patient, family, or bystanders
 - Interferes with the patient's ability to perform basic life functions and activities of daily living
- Behavior Activity Rating Score ≥ 5

Potential Complications:

- Injury to patient, provider, or bystander
- Need to move to restraint procedure
- Exacerbation of agitated condition

Procedure references:

1. Palmer J. (2019). Joint Commission Issues De-escalation Guidebook for Healthcare Facilities and Workers. Patient Safety and Quality Healthcare (PSQH). <https://www.psqh.com/analysis/joint-commission-issues-de-escalation-guidebook-for-healthcare-facilities-and-workers/>
2. Richmond JS, Berlin JS, Fishkind AV, et al. (2012). Verbal De-escalation of the Agitated Patient: Consensus Statement of the American Association for Emergency Psychiatry Project BETA De-escalation Workgroup. West J Emerg Med 13(1):17-25. doi: 10.5811/westjem.2011.9.6864



Standards Procedure (Skill) Assessment / Screening Section

Fall Risk Screen (MedFrat) Novant Facilities

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- All patient situations in which the EMS provider believes that the patient is eligible for triage at Novant New Hanover Regional Medical Center and Novant Brunswick Medical Center.

Procedure:

If the answer is **YES** to **ANY** of the questions below, patient is a Fall Risk. If transporting to Novant New Hanover Regional Medical Center they will need a **Yellow** Fall Risk arm band. These can be found at New Hanover in the radio room and/or EMS entrance. Please notify care team of risk at both facilities.

Questions:

- Any history of falling in the last 3 months?
- Is the patient confused or disoriented?
- Is the patient sedated or intoxicated?
- Does the patient have an impaired gait?
- Does the patient use a mobility assistance device?
- Is the patient experiencing altered elimination (a change in the bodily process of expelling waste products from the body by emptying either the bowels or the bladder)?

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Standards Procedure (Skill) Assessment / Screening Section

Clinical Opiate Withdrawal Scale (COWS)

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- All patient situations in which the EMS provider believes that the patient is eligible for measuring symptoms for opiate withdrawals over a period of time.

Procedure:

For each item, write in the number that best describes the patient's signs or symptom. Rate on just the apparent relationship to opiate withdrawal.

Patient's Name _____	Date _____			
Enter scores at time zero, 30 mins after first dose, 2 hours after first dose, etc.				
	Times: _____			
Resting Pulse Rate: (record beats /min) <i>Measured after patient is sitting or lying for one minute.</i> 0 pulse rate 80 or below 1 pulse rate 81-100 2 pulse rate 101-120 4 pulse rate greater than 120				
Sweating: <i>over past ½ hour not accounted for by room temperature of patient activity.</i> 0 no report of chills or flushing 1 subjective report of chills or flushing 2 flushed or observable moistness on face 3 beads of sweat on brow or face 4 sweat streaming off face				
Restlessness: <i>Observation during assessment</i> 0 able to sit still 1 reports difficulty sitting still, but is able to do so 3 frequent shifting or extraneous movements of legs/arms 5 Unable to sit still for more than a few seconds				
Pupil Size 0 pupils pinned or normal size for room light 1 pupils possibly larger than normal for room light 2 pupils moderately dilated 5 pupils so dilated that only the rim of the iris is visible				
Bone or Joint Aches: <i>If PT was having pain previously only the additional component attributed to opiate withdrawal is scored.</i> 0 Not present 1 Mild diffuse discomfort 2 Patient reports severe diffuse aching of joint/ muscles 4 Patient is rubbing joints/ muscles and is unable to sit still because of discomfort				



Clinical Opiate Withdrawal Scale (COWS) cont.

<p>Runny nose or tearing: <i>Not accounted for by cold symptoms or allergies</i></p> <p>0 not present 1 nasal stuffiness or unusually moist eyes 2 nose running or tearing 4 nose constantly running or tears streaming down cheeks</p>				
<p>GI upset: <i>over last ½ hour</i></p> <p>0 no GI symptoms 1 stomach cramps 2 nausea or loose stool 3 vomiting or diarrhea 5 multiple episodes of diarrhea or vomiting</p>				
<p>Tremor: <i>observations of outstretched hands</i></p> <p>0 no tremor 1 tremor can be felt but not observed 2 slight tremor observed 4 gross tremor or muscle twitching</p>				
<p>Yawning: <i>observation during assessment</i></p> <p>0 no yawning 1 yawning once or twice during assessment 2 yawning three- or four-times during assessment 4 yawning several times/ minute</p>				
<p>Anxiety or Irritability</p> <p>0 none 1 patient reports increasing anxiety or irritability 2 patient obviously anxious 4 patient so irritable or anxious that participation in the assessment is difficult</p>				
<p>Gooseflesh skin</p> <p>0 skin is smooth 3 piloerections of skin can be felt or hairs standing up on arms 5 prominent piloerections</p>				
Total Scores				
Observer's Initials				

Score: 5-12 = mild; 13-24 = moderate; 25-36 = moderate withdrawal; more than 36 = severe withdrawal

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Cardiac: 12 Lead ECG


Clinical Indications:

- Suspected cardiac patient
- Suspected tricyclic overdose
- Electrical injuries
- Syncope

B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Procedure:

1. Assess patient and monitor cardiac status.
2. Administer oxygen as patient condition warrants.
3. If patient is unstable, definitive treatment is the priority. If patient is stable or stabilized after treatment, perform a 12 Lead ECG.
4. Prepare ECG monitor and connect patient cable with electrodes.
5. Enter the required patient information (patient name, etc.) into the 12 lead ECG device.
6. Expose chest and prep as necessary. Modesty of the patient should be respected.
7. Apply chest leads and extremity leads using the following landmarks:
 - RA -Right arm
 - LA -Left arm
 - RL -Right leg
 - LL -Left leg
 - V1 -4th intercostal space at right sternal border
 - V2 -4th intercostal space at left sternal border
 - V3 -Directly between V2 and V4
 - V4 -5th intercostal space at midclavicular line
 - V5 -Level with V4 at left anterior axillary line
 - V6 -Level with V5 at left midaxillary line
8. Instruct patient to remain still.
9. Press the appropriate button to acquire the 12 Lead ECG.
10. If the monitor detects signal noise (such as patient motion or a disconnected electrode), the 12 Lead acquisition will be interrupted until the noise is removed.
11. Monitor the patient while continuing with the treatment protocol.
12. The Lifepak 15 will automatically print a 12 Lead if changes are detected.
13. Download data as per guidelines and attach a copy of the 12 lead to the PCR.
14. Document the procedure, time, interpretation and results on/with the patient care report (PCR)
15. **Go to page 2 for a suspected Code STEMI**



Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Code STEMI Activation

Clinical Indications:

- Suspected ST-Elevation MI (STEMI), based on patient presentation and ECG


P PARAMEDIC P

Procedure:

1. Reading the 12 Lead:
 - a. Is there a **LBBB not known to be old?**
 - b. Absence or **presence of LVH?**
 - c. Absence or **presence of other STEMI imitators?**If any of those are present see below - STEMI imitators, when to suspect STEMI.
2. Activation of a Code STEMI should be performed within 5 minutes of an interpretation of STEMI by a Paramedic to the receiving facility. To activate state "**Code STEMI Activation**" on radio transmission with receiving facility. Transport to the closest STEMI hospital or preferred STEMI hospital due to patient with known cardiologist at that facility.
 - a. New Hanover Regional Medical Center activate via Pulsara.
 - b. Grand Strand Regional Medical Center activate via radio and advise you are transmitting the 12-Lead. Transmit to GSRMC.
 - c. McLeod Seacoast activate by calling 1-843-366-3634 and transmit the 12-Lead. (the **ONLY** time c-comm should be asked to activate CODE STEMI is in instances of radio failure)
3. Follow-up radio call-in should include:
 - a. Patient age and gender
 - b. Patient cardiologist or state they do not have one
 - c. Clinical presentation, history, symptoms
 - d. Vital signs
 - e. Treatments and procedures performed
 - f. Estimated Time of Arrival

STEMI Imitators, when to suspect STEMI:

1. **Benign Early Repolarization** - reciprocal changes present
2. **Bundle Branch Block** - left or right? look at V1 to identify
 - a. **LBBB** - Sgarbossa Criteria
 - Concordant ST elevation ≥ 1 mm (see image)
 - or
 - ST segment depression ≥ 1 mm in lead V1, V2, V3
 - b. **RBBB** - concordant ST elevation
3. **LVH** - concordant ST elevation (see image)
4. **Pericarditis** - reciprocal changes present



Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Cardiac: Cardioversion

P PARAMEDIC P

Clinical Indications:

- Unstable patient with a tachydysrhythmia (rapid atrial fibrillation, supraventricular tachycardia, ventricular tachycardia)
- Patient is not pulseless (the pulseless patient requires unsynchronized cardioversion, i.e., defibrillation)

Procedure:

1. Ensure the patient is attached properly to a monitor/defibrillator capable of synchronized cardioversion.
2. Have all equipment prepared for unsynchronized cardioversion/defibrillation if the patient fails synchronized cardioversion and the condition worsens.
3. Consider the use of pain or sedating medications.
4. Set energy selection to the appropriate setting.
5. Set monitor/defibrillator to synchronized cardioversion mode.
6. Make certain all personnel are clear of patient.
7. Press and hold the shock button to cardiovert. Stay clear of the patient until you are certain the energy has been delivered. NOTE: It may take the monitor/defibrillator several cardiac cycles to “synchronize”, so there may a delay between activating the cardioversion and the actual delivery of energy.
8. Note patient response and perform immediate unsynchronized cardioversion/defibrillation if the patient’s rhythm has deteriorated into pulseless ventricular tachycardia/ventricular fibrillation, following the procedure for Defibrillation-Manual.
9. If the patient’s condition is unchanged, repeat steps 2 to 8 above, using escalating energy settings.
10. Repeat until maximum setting or until efforts succeed. Consider discussion with medical control if cardioversion is unsuccessful after 2 attempts.
11. Note procedure, response, and time in the patient care report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System. Assessment should include direct observation at least once per certification cycle., or other mechanisms as deemed appropriate by the local EMS System.



Cardiac: External Pacing

P PARAMEDIC P

Clinical Indications:

- Patients with symptomatic bradycardia (less than 60 per minute) with signs and symptoms of inadequate cerebral or cardiac perfusion such as:
 - Chest Pain
 - Hypotension
 - Pulmonary Edema
 - Altered Mental Status, Confusion, etc.
 - Ventricular Ectopy
- Asystole, pacing must be done early to be effective.
- PEA, where the underlying rhythm is bradycardic and reversible causes have been treated.

Procedure:

1. Attach standard four-lead monitor.
2. Apply defibrillation/pacing pads to chest and back:
 - One pad to left mid chest next to sternum
 - One pad to mid left posterior chest next to spine.
3. Rotate selector switch to pacing option.
4. Adjust heart rate to 70 BPM for an adult and 100 BPM for a child.
5. Note pacer spikes on EKG screen.
6. Slowly increase output until capture of electrical rhythm on the monitor.
7. If unable to capture while at maximum current output, stop pacing immediately.
8. If capture observed on monitor, check for corresponding pulse and assess vital signs.
9. Consider the use of sedation or analgesia if patient is uncomfortable.
10. Document the dysrhythmia and the response to external pacing with ECG strips in the PCR.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System. Assessment should include direct observation at least once per certification cycle.



Standards Procedure (Skill) Cardiac Section

Cardiac: Cardiopulmonary Resuscitation (CPR)

Clinical Indications:

- Basic life support for the patient in cardiac arrest

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Procedure:

1. Assess the patient's level of responsiveness.
2. If no response, open the patient's airway with the head-tilt, chin-lift and look, listen, and feel for respiratory effort. If the patient may have sustained C-spine trauma, use the modified jaw thrust while maintaining immobilization of the C-spine. For infants, positioning the head in the sniffing position is the most effective method for opening the airway.
3. Check for pulse (carotid for adults and older children, brachial for infants) for at least 10 seconds. If no pulse, begin chest compressions based on chart below:

Age	Location	Depth	Rate
Infant	Over sternum, between nipples (inter-mammary line), 2-3 fingers	At least 1/3 AP diameter of chest About 1.5 inches 4 cm	Continuous compressions at least 100 – 120/minute
Child	Over sternum, just cephalad from xyphoid process, heel of one hand	At least 1/3 AP diameter of chest About 2 inches 5 cm	Continuous compressions at least 100 – 120/minute
Adult	Over sternum, just cephalad from xyphoid process, hands with interlocked fingers	At least 2 inches 5 cm	Continuous compressions at least 100 – 120/minute

4. If patient is an adult, go to step 5. If no respiratory effort in a pediatric patient, give two ventilations. If air moves successfully, go to step 5. If air movement fails, proceed to the Airway Obstruction Procedure.
5. Go to Cardiac Arrest Procedure. Begin ventilations in the adult as directed in the Cardiac Arrest Procedure
6. Provide 1 breath every 6 seconds with the BVM or BIAD. Use EtCO₂ to guide your ventilations as directed in the Cardiac Arrest Protocol.
7. Chest compressions should be provided in an uninterrupted manner. Only brief interruptions (< 5 seconds with a maximum of 10 seconds) are allowed for rhythm analysis, defibrillation, and performance of procedures
8. Document the time and procedure in the Patient Care Report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Cardiac: Defibrillation-Automated

Clinical Indications:

- Patients in cardiac arrest (pulseless, non-breathing).
- Age < 8 years, use Pediatric Pads if available.

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Contraindication:

- Pediatric patients who are so small that the pads cannot be placed without touching one another.

Procedure:

1. **If multiple rescuers available, one rescuer should provide uninterrupted chest compressions while the AED is being prepared for use.**
2. Apply defibrillator pads per manufacturer recommendations. Based on 2010 guidelines, place pads preferably in AP or AL position when implanted devices (pacemakers, AICDs) occupy preferred pad positions and attempt to avoid placing directly over device.
3. Remove any medication patches on the chest and wipe off any residue.
4. If necessary, connect defibrillator leads: white to the anterior chest pad and the red to the posterior pad.
5. Activate AED for analysis of rhythm.
6. **Stop CPR and clear the patient** for rhythm analysis. Keep interruption in CPR as brief as possible.
7. Defibrillate if appropriate by depressing the “shock” button. **Assertively state “CLEAR” and visualize that no one, including yourself, is in contact with the patient prior to defibrillation.** The sequence of defibrillation charges is preprogrammed for monophasic defibrillators. Biphasic defibrillators will determine the correct joules accordingly.
8. Begin CPR (chest compressions and ventilations) immediately after the delivery of the defibrillation.
9. After 2 minutes of CPR, analyze rhythm and defibrillate if indicated. Repeat this step every 2 minutes.
10. If “no shock advised” appears, perform CPR for two minutes and then reanalyze.
11. Transport and continue treatment as indicated.
12. **Keep interruption of CPR compressions as brief as possible. Adequate CPR is a key to successful resuscitation.**
13. **If pulse returns please use the Post Resuscitation Protocol**

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System. Assessment should include direct observation at least once per certification cycle.



Standards Procedure (Skill) Cardiac Section

Cardiac: Defibrillation-Manual

P PARAMEDIC P

Clinical Indications:

- Cardiac arrest with ventricular fibrillation or pulseless ventricular tachycardia

Procedure:

1. **Ensure that Chest Compressions are adequate and interrupted only when absolutely necessary.**
2. Clinically confirm the diagnosis of cardiac arrest and identify the need for defibrillation.
3. After application of an appropriate conductive agent if needed, apply defibrillation hands free pads (recommended to allow more continuous CPR) or paddles to the patient's chest in the proper position
 - Paddles: right of sternum at 2nd ICS and anterior axillary line at 5th ICS
 - Pads: anterior-posterior position

For patients with implanted pacers/defibrillators, paddles or pads can be in AP or AL positions. The presence of implanted pacers/defibrillators should not delay defibrillation. Attempt to avoid placing paddles or pads directly above device.

4. Set the appropriate energy level
5. Charge the defibrillator to the selected energy level. **Continue chest compressions while the defibrillator is charging.**
6. If using paddles, assure proper contact by applying 25 pounds of pressure on each paddle.
7. **Hold Compressions, assertively state, "CLEAR" and visualize that no one, including yourself, is in contact with the patient.**
8. Deliver the countershock by depressing the discharge button(s) when using paddles, or depress the **shock button** for hands free operation.
9. Immediately resume chest compressions and ventilations for 2 minutes. After 2 minutes of CPR, analyze rhythm and check for pulse only if appropriate for rhythm.
10. Repeat the procedure every two minutes as indicated by patient response and ECG rhythm.
11. Keep interruption of CPR compressions as brief as possible. Adequate CPR is a key to successful resuscitation.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System. Assessment should include direct observation at least once per certification cycle.



Cardiac: Defibrillation-Dual or Double

P PARAMEDIC P

Clinical Indications:


- Cardiac arrest with persistent ventricular fibrillation or pulseless ventricular tachycardia.
- Refractory ventricular fibrillation or pulseless ventricular tachycardia where ≥ 3 shocks delivered.

Procedure:

1. Ensure that Chest Compressions are adequate and interrupted only when absolutely necessary.

2. Clinically confirm the diagnosis of cardiac arrest and identify the need for defibrillation.
3. Prepare sites for second pad set attachment and apply defibrillation hands free pads:

- Pads: First defibrillator pads in anterior-posterior position
- Pads: Second defibrillator pads in anterior-lateral position:
- Ensure pads are not in contact with one another.



For patients with implanted pacers/defibrillators:
Avoid placing paddles or pads directly above device.

4. Set the appropriate energy level and assure controls for both defibrillator / monitors are accessible to provider performing defibrillation.
5. At next pulse / rhythm check, if refractory or persistent VF/VT continues:
Charge the defibrillator to the selected energy level.
Continue chest compressions while the defibrillator is charging.
6. Optional: Agencies may provide a single shock at this point with the second defibrillator / monitor to provide a change in energy vector delivered to the heart then move to step 7 if VF / VT persists.
- 7. When both monitor / defibrillators have reached selected energy setting:**
Hold Compressions, assertively state, "CLEAR" and visualize that no one, including yourself, is in contact with the patient.
2 options at this point:
Option 1 (double simultaneous): Provider depresses both defibrillator shock buttons simultaneously.
Option 2 (dual sequential): Provider depresses monitor 1 shock button and then immediately following, depresses monitor 2 shock button.
8. Immediately resume chest compressions and ventilations for 2 minutes. After 2 minutes of CPR, analyze rhythm and check for pulse only if appropriate for rhythm.
9. Repeat the procedure every two minutes as indicated by patient response and ECG rhythm.
10. Keep interruption of CPR compressions as brief as possible. Adequate CPR is a key to successful resuscitation.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System. Assessment should include direct observation at least once per certification cycle.



Cardiac: 15 Lead ECG



Clinical Indications:

- Suspected cardiac patient
- All patients with a normal 12 lead ECG
- All inferior STEMI's
- All patients with a strong suspicion of having posterior involvement

B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Procedure:

1. After performing a 12 Lead ECG
 - V4R**- move Lead V4 to the 5th intercostal space at the midclavicular line on the right side of the chest
 - V8** - move V5 wire to the 5th intercostal space mid-scapular line
 - V9** - move V6 wire between V8 and spine
2. Instruct patient to remain still.
3. Press the appropriate button to acquire the 12 Lead ECG.
4. If the monitor detects signal noise (such as patient motion or a disconnected electrode), the 12 Lead acquisition will be interrupted until the noise is removed.
5. Once acquired, re-label the 12-Lead with V4R, V8 and V9.
6. Monitor the patient while continuing with the treatment protocol.
7. Download data as per guidelines and attach a copy of the 12 lead to the PCR.
8. Document the procedure, time, and results on/with the patient care report (PCR)



Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Standards Procedure (Skill) Cardiac Section Cardiac: Mechanical CPR (LUCAS Device)

Clinical Indications:

The LUCAS device can be considered:

- During the management of cardiac arrest when there is an insufficient number of providers to continue high quality CPR.
- When the patient must be moved or transported and effective CPR can not be maintained or is impossible.
- If the use or set up of the device WILL NOT delay Manual Compressions or Defibrillations. In the event of a delay manual compressions will be continued without hesitation.
- In the event of mechanical failure Manual Compressions will be resumed immediately.


	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Do not use the LUCAS Device if:

- The patient is a pediatric patient and you are following PC1, PC4, and/or PC7.
- The patient has suffered a traumatic arrest.
- It is not possible to position the LUCAS safely or correctly around the patient's chest.
- The patient is too small; if the LUCAS alerts with 3 fast signals when lowering the suction cup.
- The patient is too large. If you cannot lock the upper part of the LUCAS to the back plate without compressing the patient's chest.

Procedure:

1. Manual high quality compressions will be administered according to current protocols.
2. Defibrillation / Pacing Pads must be positioned so that the pads and cables are not under the suction cup.
3. Open the LUCAS case and push the On/Off button on the control panel for one second and it will start the self test.
4. Remove the back plate from the case. Communicate with other team members as to when the back plate will be deployed, preferably on the next rhythm check.
5. Make sure the patients head is supported and carefully place the back plate under the patient , immediately below the armpits using one of these procedures:
 - raise the patient up by lifting the shoulders or pulling their arms a small distance to get the board under them OR
 - roll the patient , insert board and roll back
6. Resume CPR
7. Pull the RELEASE RINGS once to open the CLAW LOCKS. Then let go of the rings. Stop CPR briefly while attaching the upper part of the back plate. Listen for "click" sound. Pull up once to assure attachment.
8. Push down suction cup. Adjust position if needed. The lower edge of the suction cup should be immediately above the sternum.
9. Press the pause button to lock the start position. Push play (continuous) if advanced airway in place or push play (30:2) if no airway in place.
10. Attach stabilization strap and place patient arms in the Wrist Straps.
11. Pause for rhythm / pulse checks every two minutes, no longer than 5 seconds. LUCAS should continue through defibrillations.





Standards Procedure (Skill) Parenteral Access Section

Parenteral Access: Arterial Line Maintenance

Clinical Indications:

P **PARAMEDIC** **P**

- Transport of a patient with an existing arterial line.

Procedure:

1. Make certain arterial line is secured prior to transport, including intersection of arterial catheter and IV/Monitoring lines.
2. Use available equipment for monitoring of arterial pressures via arterial line.
3. Do not use the arterial line for administration of any fluids or medications.
4. If there is any question regarding dislodgement of the arterial line and bleeding results, remove the line and apply direct pressure over the site for at least five minutes before checking to ensure hemostasis.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Parenteral Access: Venous Blood Draw

Clinical Indications:

- Collection of a patient's blood for laboratory analysis

A	AEMT	A
P	PARAMEDIC	P

Procedure:

1. Utilize universal precautions as per OSHA.
2. Select vein and prep as usual.
3. Select appropriate blood-drawing devices.
4. Draw appropriate tubes of blood for lab testing.
5. Assure that the blood samples are labeled with the correct information (a minimum of the patient's name, along with the date and time the sample was collected).
6. Deliver the blood tubes to the appropriate individual at the hospital.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Standards Procedure (Skill) Parenteral Access Section

Parenteral Access: Central Line Maintenance

Clinical Indications:

P **PARAMEDIC** **P**

- Transport of a patient with a central venous pressure line already in place

Procedure:

1. Prior to transportation, ensure the line is secure.
2. Medications and IV fluids may be administered through a central venous pressure line. Such infusions must be held while the central venous pressure is transduced to obtain a central venous pressure, but may be restarted afterwards.
3. Do not manipulate the central venous catheter.
4. If the central venous catheter becomes dysfunctional, does not allow drug administration, or becomes dislodged, contact medical control.
5. Document the time of any pressure measurements, the pressure obtained, and any medication administration in the patient care report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Standards Procedure (Skill) Parenteral Access Section

Parenteral Access: Epidural Catheter Maintenance

Clinical Indications:

P **PARAMEDIC** **P**

- Presence of an epidural catheter in a patient requiring transport

Procedure:

1. Prior to transport, ensure catheter is secure and that transport personnel are familiar with medication(s) being delivered and devices used to control medication administration.
2. No adjustments in catheter position are to be attempted.
3. No adjustments in medication dosage or administration are to be attempted without direct approval from on-line medical control.
4. Report any complications immediately to on-line medical control.
5. Document the time and dose of any medication administration or rate adjustment in the patient care report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Standards Procedure (Skill) Parenteral Access Section

Parenteral Access: Ventricular Catheter Maintenance

P **PARAMEDIC** **P**

Clinical Indications:

- Transport of a patient with an intra-ventricular catheter in place

Procedure:

1. Prior to transport, ensure the catheter is secure.
2. Prior to transport, determine from the referring hospital/physician the desired patient position (e.g., supine, head of bed elevated 30 degrees, etc.).
3. Prior to transport, determine the height at which the drain is to be maintained, given the patient position desired from #2 above (if applicable).
4. Do not manipulate or move the drain.
5. If the patient or height of the drain is altered, immediately correct based on the pre-determined configuration in step 2 and 3 above.
6. Report any problems immediately to on-line medical control.
7. Document the time and any adjustments or problems in the patient care report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Standards Procedure (Skill) Parenteral Access Section

Parenteral Access: Existing Catheters

Clinical Indications:

P **PARAMEDIC** **P**

- Inability to obtain adequate peripheral access.
- Access of an existing venous catheter for medication or fluid administration.
- Central venous access in a patient in cardiac arrest.

Procedure:

1. Clean the port of the catheter with alcohol wipe.
2. Using sterile technique, withdraw 5-10 ml of blood and discard syringe in sharps container.
3. Using 5cc of normal saline, access the port with sterile technique and gently attempt to flush the saline.
4. If there is no resistance, no evidence of infiltration (e.g., no subcutaneous collection of fluid), and no pain experienced by the patient, then proceed to step 5. If there is resistance, evidence of infiltration, pain experienced by the patient, or any concern that the catheter may be clotted or dislodged, do not use the catheter.
5. Begin administration of medications or IV fluids slowly and observe for any signs of infiltration. If difficulties are encountered, stop the infusion and reassess.
6. Record procedure, any complications, and fluids/medications administered in the Patient Care Report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Standards Procedure (Skill) Parenteral Access Section

Parenteral Access: External Jugular Access

A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- External jugular vein cannulation is indicated in a critically ill patient ≥ 8 years of age who requires intravenous access for fluid or medication administration and in whom an extremity vein is not obtainable.
- External jugular cannulation can be attempted initially in life threatening events where no obvious peripheral site is noted.

Procedure:

1. Place the patient in a supine head down position. This helps distend the vein and prevents air embolism.
2. Turn the patient's head toward the opposite side if no risk of cervical injury exists.
3. Prep the site as per peripheral IV site.
4. Align the catheter with the vein and aim toward the same side shoulder.
5. "Tourniqueting" the vein lightly with one finger above the clavicle, puncture the vein midway between the angle of the jaw and the clavicle and cannulate the vein in the usual method.
6. Attach the IV and secure the catheter avoiding circumferential dressing or taping.
7. Document the procedure, time, and result (success) on/with the patient care report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Parenteral Access: Venous-Extremity

Clinical Indications:

- Any patient where intravenous access is indicated (significant trauma, emergent or potentially emergent medical condition).

A	AEMT	A
P	PARAMEDIC	P

Procedure:

- Saline locks may be used as an alternative to an IV tubing and IV fluid in every protocol at the discretion of the ALS professional.
- Paramedic/AEMT can use intraosseous access where threat to life exists as provided for in the Venous Access-Intraosseous procedure.
- Use the largest catheter bore necessary based upon the patient's condition and size of veins.
- Fluid and setup choice is preferably:
 - Lactated Ringers with a macro drip (10 gtt/cc) for burns
 - Normal Saline with a macro drip (10 gtt/cc) for medical conditions, trauma or hypotension
 - Normal Saline with a micro drip (60 gtt/cc) for medication infusions
- Inspect the IV solution for expiration date, cloudiness, discoloration, leaks, or the presence of particles.
- Connect IV tubing to the solution in a sterile manner. Fill the drip chamber half full and then flush the tubing bleeding all air bubbles from the line.
- Place a tourniquet around the patient's extremity to restrict venous flow only.
- Select a vein and an appropriate gauge catheter for the vein and the patient's condition.
- Prep the skin with an antiseptic solution.
- Insert the needle with the bevel up into the skin in a steady, deliberate motion until the bloody flashback is visualized in the catheter.
- Advance the catheter into the vein. **Never** reinsert the needle through the catheter. Dispose of the needle into the proper container without recapping.
- Draw blood samples when appropriate.
- Remove the tourniquet and connect the IV tubing or saline lock.
- Open the IV to assure free flow of the fluid and then adjust the flow rate as per protocol or as clinically indicated.

Rates are preferably:

- Adult: KVO: 60 cc/hr (1 gtt/ 6 sec for a macro drip set)
- Pediatric: KVO: 30 cc/hr (1 gtt/ 12 sec for a macro drip set)

If shock is present:

- Adult: 500 cc fluid boluses repeated as long as lungs are dry and BP < 90. Consider a second IV line.
- Pediatric: 20 cc/kg boluses repeated PRN for poor perfusion.

- Cover the site with a sterile dressing and secure the IV and tubing.
- Label the IV with date and time, catheter gauge, and name/ID of the person starting the IV.
- Document the procedure, time and result (success) on/with the patient care report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.

Standards Procedure (Skill) Parenteral Access Section

Parenteral Access: Intraosseous

A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Rapid, regular IV access is unavailable with any of the following:
- Cardiac arrest.
- Multisystem trauma with severe hypovolemia.
- Severe dehydration with vascular collapse and/or loss of consciousness.
- Respiratory failure / Respiratory arrest.
- Burns.

Contraindications:

- Fracture proximal to proposed intraosseous site.
- History of Osteogenesis Imperfecta
- Current or prior infection at proposed intraosseous site.
- Previous intraosseous insertion or joint replacement at the selected site.

Procedure:

1. Don personal protective equipment (gloves, eye protection, etc.).
 - **Proximal tibia:** Identify anterior-medial aspect of the proximal tibia (bony prominence below the knee cap). The insertion location will be 1-2 cm (2 finger widths) below this.
 - **Distal tibia:** Not approved for BCEMS
 - **Distal femur:** Not approved for BCEMS
 - **Proximal humerus:** Acceptable insertion site for adult patients. Locate the insertion site 1 – 2 cm above the surgical neck on the most prominent aspect of the greater tubercle. This is located on the lateral aspect of the ball of the humerus. Direct the needle at a 45 degree angle or toward the opposite hip.
2. Prep the site recommended by the device manufacturer with an antiseptic solution.
3. For manual pediatric devices, hold the intraosseous needle at a 90 degree angle to bone surface, aimed away from the nearby joint and epiphyseal plate, twist the needle handle with a rotating grinding motion applying controlled downward force until a “pop” or “give” is felt indicating loss of resistance. Do not advance the needle any further.
5. For the EZ-IO intraosseous device, hold the intraosseous needle at a 90 degree angle to bone surface, aimed away from the nearby joint and epiphyseal plate, power the driver until a “pop” or “give” is felt indicating loss of resistance. Do not advance the needle any further. Blue (25mm) IO needle is typically recommended for tibial IO placement (adults and children), Yellow (45 mm) IO needle is typically utilized for proximal humerus, and Pink (15 mm) should only be utilized in neonates. IO needle choice may vary based on a patients body habitus, or abnormal weight for age.
6. For the Bone Injection Gun (BIG), find and mark the manufacturers recommended site. Position the device and pull out the safety latch. Trigger the BIG at 90° to the bone surface and remove the injection device.

Standards Procedure (Skill) Parenteral Access Section

Parenteral Access: Intraosseous

A	AEMT	A
P	PARAMEDIC	P

7. Remove the stylette and place in an approved sharps container.
8. Attach a syringe filled with at least 5 ml NS; For IO manual devices only, verify placement by aspirating bone marrow. Inject at least 5 ml of NS to clear the lumen of the needle. Look for infiltrations around site.
9. Attach the IV line and adjust flow rate. A pressure bag may assist with achieving desired flows.
10. Stabilize and secure the needle with dressings and tape.
11. Paramedics and AEMT may administer 20 to 50 mg (1 to 2.5 ml) of 2% Lidocaine in adult patients who experience infusion-related pain. This may be repeated prn to a maximum of 60 mg (3 ml).
For infant/child dose is 0.5mg/kg, Not to exceed 40mg.
12. Following the administration of any IO medications, flush the IO line with 10 ml of IV fluid.
13. Document the procedure, time, and result (success) on/with the patient care report (PCR).

Certification Requirements:

Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System. Assessment should include direct observation at least once per certification cycle.



Standards Procedure (Skill) Parenteral Access Section Parenteral Access: Swan-Ganz Catheter Maintenance

P PARAMEDIC P

Clinical Indications:

- Transport of a patient with a Swan-Ganz catheter that is in place prior to transport.

Procedure:

1. Make certain catheter is secure prior to transport.
2. Under the supervision of the nurse or physician caring for the patient, make certain the transport personnel are aware of the depth at which the catheter is secured.
3. **UNDER NO CIRCUMSTANCES SHOULD TRANSPORT PERSONNEL ADVANCE THE SWAN-GANZ CATHETER.**
4. The sterile plastic sheath that surrounds the catheter should not be manipulated.
5. The ports of the catheter may be used to continue administration of medications or IV fluids that were initiated prior to transport. These should be used as any other IV port with attention to sterile technique.
6. If applicable, measurements from the catheter may be obtained during transport and used to guide care as per local protocols and medical control orders.
7. If at anytime during the transport difficulties with the function of the Swan-Ganz catheter is noted, contact medical control.
8. Document the time and any adjustments or problems associated with the catheter in the patient care report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Standards Procedure (Skill) Respiratory Section

Airway: Suctioning-Advanced

A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Obstruction of the airway (secondary to secretions, blood, or any other substance) in a patient currently being assisted by an airway adjunct such as a naso-tracheal tube, endotracheal tube, Combitube, tracheostomy tube, or a cricothyrotomy tube.

Procedure:

1. Ensure suction device is in proper working order.
2. Preoxygenate the patient as is possible.
3. Attach suction catheter to suction device, keeping sterile plastic covering over catheter.
4. Using the suprasternal notch and the end of the airway into the catheter will be placed as guides, measure the depth desired for the catheter (judgment must be used regarding the depth of suctioning with cricothyrotomy and tracheostomy tubes).
5. If applicable, remove ventilation devices from the airway.
6. With the thumb port of the catheter uncovered, insert the catheter through the airway device.
7. Once the desired depth (measured in #4 above) has been reached, occlude the thumb port and remove the suction catheter slowly.
8. A small amount of Normal Saline (10 ml) may be used if needed to loosen secretions for suctioning.
9. Reattach ventilation device (e.g., bag-valve mask) and ventilate the patient
10. Document time and result in the patient care report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Standards Procedure (Skill) Respiratory Section

Respiratory: Suctioning-Basic

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Obstruction of the airway (secondary to secretions, blood, or any other substance) in a patient who cannot maintain or keep the airway clear.

Procedure:

1. Ensure suction device is in proper working order with suction tip in place.
2. Preoxygenate the patient as is possible.
3. Explain the procedure to the patient if they are coherent.
4. Examine the oropharynx and remove any potential foreign bodies or material which may occlude the airway if dislodged by the suction device.
5. If applicable, remove ventilation devices from the airway.
6. Use the suction device to remove any secretions, blood, or other substance.
7. The alert patient may assist with this procedure.
8. Reattach ventilation device (e.g., bag-valve mask) and ventilate or assist the patient
9. Record the time and result of the suctioning in the patient care report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Standards Procedure (Skill) Respiratory Section

Respiratory: Nebulizer Inhalation Therapy

Clinical Indications:

- Patients experiencing bronchospasm.

B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Procedure:

1. Gather the necessary equipment.
2. Assemble the nebulizer kit.
3. Instill the premixed drug (such as Albuterol or other approved drug) into the reservoir well of the nebulizer.
4. Connect the nebulizer device to oxygen at 4 - 6 liters per minute or adequate flow to produce a steady, visible mist.
5. Instruct the patient to inhale normally through the mouthpiece of the nebulizer. The patient needs to have a good lip seal around the mouthpiece.
6. The treatment should last until the solution is depleted. Tapping the reservoir well near the end of the treatment will assist in utilizing all of the solution.
7. Monitor the patient for medication effects. This should include the patient's assessment of his/her response to the treatment and reassessment of vital signs, ECG, and breath sounds.
8. Assess and document peak flows before and after nebulizer treatments.
9. Document the treatment, dose, and route on/with the patient care report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Standards Procedure (Skill) Respiratory Section

Respiratory: NIPPV (Non-Invasive Positive Pressure)

Clinical Indications:

B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

- Non-Invasive Positive Airway Pressure (NIPPV) is indicated in all patients whom inadequate ventilation is suspected.

This could be as a result of Pulmonary Edema, CHF, COPD, Pneumonia, or Asthma.

- Agencies may utilize Continuous and/or Bi-Level Positive Airway Pressure Devices

Clinical Contraindications:

- Decreased Mental Status.
- Facial features or deformities that prevent an adequate mask seal.
- Excessive respiratory secretions.

Procedure:

1. Ensure adequate oxygen supply to ventilation device.
2. Explain the procedure to the patient.
3. Consider placement of a nasopharyngeal airway.
4. Place the delivery mask over the mouth and nose. Oxygen should be flowing through the device at this point.
5. Secure the mask with provided straps starting with the lower straps until minimal air leak occurs.
6. If the Positive Pressure is adjustable on the NIPPV device adjust and slowly titrate to achieve a positive pressure as follows:

Continuous pressure device:

5 – 25 cmH₂O for Pulmonary Edema, CHF, COPD, Asthma, Drowning, possible aspiration, or pneumonia.

Bi-Level pressure device:

IPAP 10 – 15 over EPAP 5 – 7 cmH₂O for Pulmonary Edema, CHF, COPD, Asthma, Drowning, possible aspiration, or pneumonia.

During titration keep IPAP – EPAP at least a difference of 5 cmH₂O

25 cmH₂O is maximum pressure that should be utilized with NIPPV.

Increasing positive pressure can cause hypotension.

Use caution or remove and re-evaluate with Systolic Blood Pressures consistently < 100 mmHg.

7. Evaluate the response of the patient assessing breath sounds, oxygen saturation, and general appearance.
8. Titrate oxygen levels to the patient's response. Many patients respond to low FIO₂ (30-50%).
9. Encourage the patient to allow forced ventilation to occur. Observe closely for signs of complications. The patient must be breathing for use of the NIPPV device.
10. Document time and response on patient care report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Childbirth

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Imminent delivery with crowning

Procedure:

1. Delivery should be controlled so as to allow a slow controlled delivery of the infant. This will prevent injury to the mother and infant.
2. Support the infant's head as needed.
3. Check the umbilical cord surrounding the neck. If it is present, slip it over the head. If unable to free the cord from the neck, double clamp the cord and cut between the clamps.
4. Suction the airway with a bulb syringe.
5. Grasping the head with hands over the ears, gently pull down to allow delivery of the anterior shoulder.
6. Gently pull up on the head to allow delivery of the posterior shoulder.
7. Slowly deliver the remainder of the infant.
8. Clamp the cord 2 inches from the abdomen with 2 clamps and cut the cord between the clamps.
9. Record APGAR scores at 1 and 5 minutes.
10. Follow the **Newly Born Protocol** for further treatment.
11. The placenta will deliver spontaneously, usually within 5 minutes of the infant. Do not force the placenta to deliver.
12. Massaging the uterus may facilitate delivery of the placenta and decrease bleeding by facilitating uterine contractions.
13. Continue transport to the hospital.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Standards Procedure (Skill) Universal Section

Decontamination

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Any patient who may have been exposed to significant hazardous materials, including chemical, biological, or radiological weapons.

Procedure:

- In coordination with HazMAT and other Emergency Management personnel, establish hot, warm and cold zones of operation.
- Ensure that personnel assigned to operate within each zone have proper personal protective equipment.
- In coordination with other public safety personnel, assure each patient from the hot zone undergoes appropriate initial decontamination. This is specific to each incident; such decontamination may include:
 - Removal of patients from Hot Zone
 - Simple removal of clothing
 - Irrigation of eyes
 - Passage through high-volume water bath (e.g., between two fire apparatus) for patients contaminated with liquids or certain solids. Patients exposed to gases, vapors, and powders often will not require this step as it may unnecessarily delay treatment and/or increase dermal absorption of the agent(s).
- Initial triage of patients should occur after step #3. Immediate life threats should be addressed prior to technical decontamination.
- Assist patients with technical decontamination (unless contraindicated based on #3 above). This may include removal of all clothing and gentle cleansing with soap and water. All body areas should be thoroughly cleansed, although overly harsh scrubbing which could break the skin should be avoided.
- Place triage identification on each patient. Match triage information with each patient's personal belongings which were removed during technical decontamination. Preserve these personnel effects for law enforcement.
- Monitor all patients for environmental illness.
- Transport patients per local protocol.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Gastric Tube Insertion

Clinical Indications:

P **PARAMEDIC** **P**

- Gastric decompression in intubated patients or for administration of activated charcoal in patients with altered mental status.

Procedure:

1. Estimate insertion length by superimposing the tube over the body from the nose to the stomach.
2. Flex the neck **if not contraindicated** to facilitate esophageal passage.
3. Liberally lubricate the distal end of the tube and pass through the patient's nostril along the floor of the nasal passage. Do not orient the tip upward into the turbinates. This increases the difficulty of the insertion and may cause bleeding.
4. In the setting of an intubated patient or a patient with facial trauma, oral insertion of the tube may be considered or preferred after securing airway.
5. Continue to advance the tube gently until the appropriate distance is reached.
6. Confirm placement by injecting 20cc of air and auscultate for the swish or bubbling of the air over the stomach. Additionally, aspirate gastric contents to confirm proper placement.
7. Secure the tube.
8. Decompress the stomach of air and food either by connecting the tube to suction or manually aspirating with the large catheter tip syringe.
9. Document the procedure, time, and result (success) on/with the patient care report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Standards Procedure (Skill) Universal Section

Gastric Tube Insertion of i-gel

Clinical Indications:

- Gastric tube insertion with use of the i-gel through the gastric channel. The gastric channel runs through the device from its proximal opening at the side of the flat connector wing to the distal tip of the non-inflatable cuff. Since the distal tip of the device fits snugly and anatomically correctly into the upper oesophageal opening, the distal opening of the gastric channel allows for the passing of a nasogastric tube to empty the stomach contents and can facilitate the venting of gas from the stomach. The gastric channel can also provide an early indication of regurgitation. Please note the size one i-gel does not have a gastric channel.

B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Procedure:

- Select the appropriate size gastric tube.
- Estimate insertion length by superimposing the tube over the body from the nose, around the ear and to the lowest part of the sternum (xiphoid process). This point can be marked with a piece of tape.
- Liberally lubricate the distal end of the tube and pass through the proximal end of the gastric tube of the i-gel.
- Continue to advance the tube gently until the appropriate distance is reached.
- Confirm placement via injecting air (20-30 ml) into proximal end of tube while performing epigastric auscultation.
- Secure the tube.
- Place tube to appropriate suction.
- Document the procedure, time, and result (success) on/with the patient care report (PCR).

i-gel size	Maximum size of Nasogastric Tube
1	N/A
1.5	10
2	12
2.5	12
3	12
4	12
5	14

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Standards Procedure (Skill) Universal Section

Injections: Subcutaneous and Intramuscular

B	<i>EMT*</i>	B
A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- When medication administration is necessary and the medication must be given via the SQ or IM route (not auto-injector), or as an alternative route in selected medications.

Procedure:

1. Receive and confirm medication order or perform according to standing orders.
2. Prepare equipment and medication expelling air from the syringe.
3. Explain the procedure to the patient and reconfirm patient allergies.
4. The most common site for subcutaneous injection is the arm.
 - Injection volume should not exceed 1 cc.
5. The possible injection sites for intramuscular injections include the arm, buttock and thigh.
 - Injection volume should not exceed 1 cc for the arm
 - Injection volume should not exceed 2 cc in the thigh or buttock.
6. The thigh should be used for injections in pediatric patients and injection volume should not exceed 1 cc.
7. Expose the selected area and cleanse the injection site with alcohol.
8. Insert the needle into the skin with a smooth, steady motion

SQ: 45-degree angle
skin pinched

IM: 90-degree angle
skin flattened

9. Aspirate for blood
10. Inject the medication.
11. Withdraw the needle quickly and dispose of properly without recapping.
12. Apply pressure to the site.
13. Monitor the patient for the desired therapeutic effects as well as any possible side effects.
14. Document the medication, dose, route, and time on/with the patient care report (PCR).

** EMT may administer Epinephrine for anaphylaxis, by IM route, if approved by the system medical director.*

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Standards Procedure (Skill) Universal Section

Restraints: Physical

Agency Name: _____
 Provider Name: _____ Paramedic
 Instructor Name: _____ EMT AEMT Paramedic Physician

SATISFACTORY
 UNSATISFACTORY



Instructor:

- Evaluate providers skill performance using the check off list below.
- Circle performance indicator.
 YES = Provider completed skill with no assistance from instructor.
 NO = Provider unable to complete skill satisfactorily following instructor intervention.
 IL = Provider able to complete skill satisfactorily following Instructor Led (teaching) intervention.

	EMR		
B	EMT	B	
A	AEMT	A	
P	Paramedic	P	

Satisfactory performance indicated with ≥ 12 YES / IL completions. (Combination of both YES and IL)

YES NO IL	<u>Verbalizes indications for physical restraints:</u> 1. Used to ensure the physical safety of the patient, provider, or others 2. Clear and immediate danger to the patient (self), provider, or others 3. When less restrictive alternatives are unsuccessful (e.g.. verbal de-escalation) 4. Delay in restraint will subject patient (self) , providers, or others to risk of serious harm
YES NO IL	<u>Verbalizes contraindications for physical restraints:</u> 1. Patient has medical decision-making capacity and refuses care 2. Patient is not a danger to self, provider, or others 3. Less restrictive alternatives have not been considered or used
YES NO IL	<u>Verbalizes assessment of resource needs:</u> Request Law Enforcement if indicated Contact Medical Control if indicated Call for additional providers if indicated Withdraw from scene if unsafe
YES NO IL	<u>Assemble appropriate equipment and personnel:</u> 1. 3 – 6 providers preferably 2. Don appropriate PPE 3. Soft nylon or leather restraints specifically manufactured for use as restraints
YES NO IL	<u>Remove potential items from all providers that can be used as weapons:</u> 1. Stethoscope, shears or scissors, hemostats, writing pens, badges, pins 2. Window punch, pocket knives, communication devices
YES NO IL	<u>Team leader assign roles to providers and discusses plans and strategies:</u> Team leader explains procedure to patient: If patient standing and will not follow directions use Procedure USP – 6. <u>If patient already on cot or flat surface:</u> <ul style="list-style-type: none"> 1 Provider to control the head and airway 1 Provider for each extremity <u>Team leader attempts verbal instructions to move patient to cot if possible:</u> <ul style="list-style-type: none"> 2 Providers take control of both wrists and elbows 2 Providers take control of both ankles and knees 1 Provider controls head/airway and 1 Provider is available for medications May place in lateral decubitus position – DO NOT place prone:
YES NO IL	Soft nylon or leather manufacture restraints are applied to wrist and ankles Secure restraints to cot with quick-release tie Examine patient for potential injuries following restrain application
YES NO IL	Both lower extremities restrained extended, cross restraints beneath lower extremities One upper extremity restrained extended by patient's side One upper extremity restrained flexed over patient's head Do not tie restraint to cot undercarriage
YES NO IL	Assess pulse, motor, and sensory immediately following application Perform pulse, motor, and sensory assessments every 15 minutes afterwards
YES NO IL	Patient must remain under constant observation by EMS at all times Appropriate monitoring equipment required based on clinical circumstances
YES NO IL	<u>Patient care report documentation requirements (restraint checklist recommended):</u> Indication for restraint use Type of restrain applied and time of application Pulse, motor, and sensory exams and time of exam



1

2

2

Instructor notes:



Restraints: Physical

Clinical Information for physical restraints

Objective of Procedure:

To protect a patient from self-harm and/or protection of providers or others on scene
Used when less restrictive alternatives have failed
Used as last resort

Scope of Practice: EMR, EMT, AEMT, and Paramedic

Indications:

Physically combative patient not responding to less restrictive means of de-escalation
Immediate danger of self-harm or harm to providers, or others on scene

Contraindications:

Less restrictive techniques have not been used or considered prior to physical restraint
Intact medical decision-making capacity refusing treatment and not a danger to self or others

Clinical Presentation:

Behavioral health crisis
Altered Mental Status with combativeness
Agitation and violence

Potential Complications:

Positional asphyxiation
Injury to patient, providers, or others
Increased mental stress to patient
Injury following escape from restraints
Bodily fluid exposure

Positioning Considerations:

Do not place patient in a supine position or place objects on top of patient
One arm should be restrained above the head
Both legs should be restrained fully extended
May place in a lateral decubitus position, supine is preferred
Head of bed should be elevated to about 30°

Procedure references:

1. Kowalski JM. (2019). Physical and Chemical Restraint. Roberts and Hedges' Clinical Procedures in Emergency Medicine and Acute Care. 7th ed.(pp 1481 - 1498). Philadelphia, PA. Elsevier.
2. Heiner JD, Moore GP. (2018). The combative and difficult. Rosen's Emergency Medicine: Concepts and Clinical Practice. 9th ed. (pp 2375 - 2386). Philadelphia, PA. Elsevier.
3. Booth JS. (2018, Dec 19). Four-Point Restraint. Retrieved from <https://emedicine.medscape.com/article/1941454-overview>.
4. Bradley S. (2017). Psychiatric Emergencies. AAOS Emergency Care and Transportation of the Sick and Injured. 11th ed. (pp.802 – 827). Burlington, MA. Jones and Bartlett Learning.



Standards Procedure (Skill) Universal Section

Restraints: Therapeutic Take Down

Agency Name: _____

Provider Name: _____ Paramedic

Instructor Name: _____ EMT AEMT Paramedic Physician

SATISFACTORY

UNSATISFACTORY



Instructor:

- Evaluate providers skill performance using the check off list below.
- Circle performance indicator.
 - YES = Provider completed skill with no assistance from instructor.
 - NO = Provider unable to complete skill satisfactorily following instructor intervention.
 - IL = Provider able to complete skill satisfactorily following Instructor Led (teaching) intervention.

Satisfactory performance indicated with ≥ 12 YES / IL completions. (Combination of both YES and IL)

	EMR	
B	EMT	B
A	AEMT	A
P	Paramedic	P

YES NO IL	<u>Verbalizes indications for physical restraints:</u> 1. Used to ensure the physical safety of the patient, provider, or others 2. Clear and immediate danger to the patient (self), provider, or others 3. When less restrictive alternatives are unsuccessful (e.g., verbal de-escalation) 4. Delay in restraint will subject patient (self), providers, or others to risk of serious harm	1
YES NO IL	<u>Verbalizes contraindications for physical restraints:</u> 1. Patient has medical decision-making capacity and refuses care 2. Patient is not a danger to self, provider, or others 3. Less restrictive alternatives have not been considered or used	1
YES NO IL	<u>Verbalizes assessment of resource needs:</u> Request Law Enforcement if indicated Contact Medical Control if indicated Call for additional providers if indicated Withdraw from scene if unsafe	2
YES NO IL	<u>Assemble appropriate equipment and personnel:</u> 1. 4 – 6 providers preferably 2. Don appropriate PPE 3. Soft nylon or leather restraints specifically manufactured for use as restraints	2
YES NO IL	<u>Remove potential items from all providers that can be used as weapons:</u> 1. Stethoscope, shears or scissors, hemostats, writing pens, badges, pins 2. Window punch, pocket knives, communication devices	
YES NO IL	<u>Team leader assign roles to providers and discusses plans and strategies:</u> Team leader explains procedure to patient If patient already on cot: 1 Provider to control the head and airway 1 Provider for each extremity 1 Provider to administer medications, if indicated If patient standing or walking: Team leader attempts verbal instructions to move patient to cot if possible 2 Providers approach from front and take control of both wrists and elbows 2 Providers approach from rear and take control of both ankles and knees 1 Provider controls head/airway and 1 Provider is available for medications	3
YES NO IL	<u>With patient supine on cot (may place in lateral decubitus – DO NOT place prone):</u> 2 Providers approach from front and take control of both wrists and elbows 2 Providers approach from rear and take control of both ankles and knees 1 Provider controls head/airway and 1 Provider is available for medications	
YES NO IL	Soft nylon or leather manufacture restraints are applied to wrist and ankles Secure restraints to cot with quick-release tie Examine patient for potential injuries following restrain application	
YES NO IL	Assess pulse, motor, and sensory immediately following application Perform pulse, motor, and sensory assessments every 15 minutes afterwards	
YES NO IL	Patient must remain under constant observation by EMS at all times Appropriate monitoring equipment required based on clinical circumstances	
YES NO IL	<u>Patient care report documentation requirements (restraint checklist recommended):</u> Indication for restraint use Type of restrain applied and time of application Pulse, motor, and sensory exams and time of exam	



Instructor notes:



Standards Procedure (Skill) Universal Section

Restraints: Therapeutic Take Down

Clinical Information for physical restraints

Objective of Procedure:

To protect a patient from self-harm and/or protection of providers or others on scene
Used when less restrictive alternatives have failed
Used as last resort

Scope of Practice: EMR, EMT, AEMT, and Paramedic

Indications:

Physically combative patient not responding to less restrictive means of de-escalation
Immediate danger of self-harm or harm to providers, or others on scene

Contraindications:

Less restrictive techniques have not been used or considered prior to physical restraint
Intact medical decision-making capacity refusing treatment and not a danger to self or others

Clinical Presentation:

Behavioral health crisis
Altered Mental Status with combativeness
Agitation and violence

Potential Complications:

Positional asphyxiation
Injury to patient, providers, or others
Increased mental stress to patient
Injury following escape from restraints
Bodily fluid exposure

Positioning Considerations:

Do not place patient in a supine position or place objects on top of patient
One arm should be restrained above the head
May place in a lateral decubitus position, supine is preferred
Head of bed should be elevated to about 30°

Procedure references:

1. Kowalski JM. (2019). Physical and Chemical Restraint. Roberts and Hedges' Clinical Procedures in Emergency Medicine and Acute Care. 7th ed.(pp 1481 - 1498). Philadelphia, PA. Elsevier.
2. Heiner JD, Moore GP. (2018). The combative and difficult. Rosen's Emergency Medicine: Concepts and Clinical Practice. 9th ed. (pp 2375 - 2386). Philadelphia, PA. Elsevier.
3. Booth JS. (2018, Dec 19). Four-Point Restraint. Retrieved from <https://emedicine.medscape.com/article/1941454-overview>.
4. Bradley S. (2017). Psychiatric Emergencies. AAOS Emergency Care and Transportation of the Sick and Injured. 11th ed. (pp.802 – 827). Burlington, MA. Jones and Bartlett Learning.



Chest Decompression

P **PARAMEDIC** **P****Clinical Indications:**

- Patients with hypotension (SBP <90), clinical signs of shock, and at least one of the following signs:
 - Jugular vein distention.
 - Tracheal deviation away from the side of the injury (often a late sign).
 - Absent or decreased breath sounds on the affected side.
 - Hyper-resonance to percussion on the affected side.
 - Increased resistance when ventilating a patient.
- Patients in traumatic arrest with chest or abdominal trauma for whom resuscitation is indicated. These patients may require bilateral chest decompression even in the absence of the signs above.

Procedure:

1. Don personal protective equipment (gloves, eye protection, etc.).
2. Administer high flow oxygen.
3. Identify and prep the site:
 - Locate the second intercostals space in the mid-clavicular line on the same side as the pneumothorax.
 - If unable to place anteriorly, lateral placement may be used at the fourth ICS mid-axillary line.
 - Prepare the site with providone-iodine ointment or solution.
4. Insert the catheter (14 gauge for adults) into the skin over the third rib and direct it just over the top of the rib (superior border) into the interspace.
5. Advance the catheter through the parietal pleura until a “pop” is felt and air or blood exits under pressure through the catheter, then advance catheter only to chest wall.
6. Remove the needle, leaving the plastic catheter in place.
7. Secure the catheter hub to the chest wall with dressings and tape.
8. Consider placing a finger cut from an exam glove over the catheter hub. Cut a small hole in the end of the finger to make a flutter valve. Secure the glove finger with tape or a rubber band. (Note – don’t waste much time preparing the flutter valve; if necessary control the air flow through the catheter hub with your gloved thumb.)

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System. Assessment should include direct observation once per certification cycle.



Chest Decompression (SPEAR)

Clinical Indications:

P PARAMEDIC P

- Adult patients with hypotension (SBP <90), clinical signs of shock, and at least one of the following signs:
 - Jugular vein distention.
 - Tracheal deviation away from the side of the injury (often a late sign).
 - Absent or decreased breath sounds on the affected side.
 - Hyper-resonance to percussion on the affected side.
 - Increased resistance when ventilating a patient.
- Patients in traumatic arrest with chest or abdominal trauma for whom resuscitation is indicated. These patients may require bilateral chest decompression even in the absence of the signs above.

Procedure:

1. Don personal protective equipment (gloves, eye protection, etc.).
2. Administer high flow oxygen.
3. Identify and prep the site:
 - Locate the second intercostal space in the mid-clavicular line on the same side as the tension pneumothorax.
 - If unable to place anteriorly, lateral placement may be used at the fourth ICS mid-axillary line.
 - Prepare the site with providone-iodine ointment or solution.
4. With the needle set stabilized between thumb and middle finger penetrate the skin and adipose tissue, it should be firmly placed against the rib. Once the SPEAR needle tip has been placed against the targeted rib, the provider should mark and hold ≤ 3 centimeters above the skin.
5. Proceed with insertion while ensuring that penetration is limited by fingertip control. Move the needle tip superiorly over the targeted rib and guide assembly into the thoracic cavity. Insert no more than 3 centimeters.
6. Remove the needle by releasing the spin-lock 1/4 turn and advance the catheter completely into the pleural cavity. The needle should remain stationary.
7. Place a one-way valve on proximal end of catheter. Secure the needle in a bio-hazard sharps container. The catheter and one-way valve assemble may be attached to low pressure suction as needed.
8. Secure the catheter hub to the chest wall with dressings and tape.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System. Assessment should include direct observation once per certification cycle.



Spinal Motion Restriction

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Need for Spinal Motion Restriction as determined by protocol.
- **Guidelines for appropriate use of long spine board (LSB) OR any equivalent device below:**

1. **Spine boards or similar rigid devices, should NOT be used during transport or during inter-facility transfers.** They should be utilized for extrication and / or patient transfers, as well as support for chest compressions. They DO NOT improve outcomes and can induce pain, agitation / anxiety, respiratory compromise, and decreased tissue perfusion at pressure points.
2. Devices such as the long or short spine board, scoop stretcher, soft-body splints, etc., should be considered extrication devices rather than transport-devices. Instead, use of Spinal Motion Restriction which includes a rigid cervical collar, manual in-line spine stabilization, maintaining spinal alignment with movement and transfers, and securing to the ambulance stretcher.
3. Penetrating trauma to head, torso, or back with no evidence of spinal injury does not require Spinal Motion Restriction.

Procedure:

1. Gather LSB, scoop, ambulance cot, or other Spinal Motion Restriction device, securing devices, and appropriate C-collar.
2. Explain the procedure to the patient and assess / record neurological exam and pulse status.
3. Place the patient in an appropriately sized C-collar while maintaining in-line stabilization of the C-spine by second provider. In-line stabilization should not involve traction / tension, but rather maintain the head in a neutral, midline position while the first rescuer applies the collar.
4. Once the collar is secure, the second rescuer should still maintain their position to ensure stabilization (the collar is helpful but will not do the job by itself.)
5. If indicated, place patient on a Spinal Motion Restriction device with log-roll or similar technique dependent on circumstances, if patient is supine or prone. During extrication or where otherwise unable to be placed prone or supine, place on Spinal Motion Restriction device by the safest method available that allows maintenance of in-line spinal stability.
6. Stabilize the patient with straps / head rolls / tape / other devices as needed. Once the head is secured to the Spinal Motion Restriction device / stretcher, the second rescuer may release manual in-line stabilization. **Once the patient arrives at the stretcher, REMOVE the rigid Spinal Motion Restriction device while maintaining spinal alignment using log-roll or multi-rescuer lift techniques and transfer and secure to the stretcher for transport.**
7. NOTE: Spinal precautions may be achieved by many methods. Never force a patient into a certain position to immobilize them. Such situations may require a second rescuer to maintain manual stabilization throughout the transport to the hospital. Special equipment such as football players in full pads and helmet may remain immobilized with helmet and pads in place.
8. Document the time of the procedure in the patient care report (PCR).

Certification Requirements:

Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Splinting

Clinical Indications:

- Immobilization of an extremity for transport, either due to suspected fracture, sprain, or injury.
- Immobilization of an extremity for transport to secure medically necessary devices such as intravenous catheters

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Procedure:

1. Assess and document pulses, sensation, and motor function prior to placement of the splint. If no pulses are present and a fracture is suspected, consider reduction of the fracture prior to placement of the splint.
2. Remove all clothing from the extremity.
3. Select a site to secure the splint both proximal and distal to the area of suspected injury, or the area where the medical device will be placed.
4. Do not secure the splint directly over the injury or device.
5. Place the splint and secure with Velcro, straps, or bandage material (e.g., kling, kerlex, cloth bandage, etc.) depending on the splint manufacturer and design.
6. Document pulses, sensation, and motor function after placement of the splint. If there has been a deterioration in any of these 3 parameters, remove the splint and reassess
7. If a femur fracture is suspected and there is no evidence of pelvic fracture or instability, the following procedure may be followed for placement of a femoral traction splint:
 - Assess neurovascular function as in #1 above.
 - Place the ankle device over the ankle.
 - Place the proximal end of the traction splint on the posterior side of the affected extremity, being careful to avoid placing too much pressure on genitalia or open wounds. Make certain the splint extends proximal to the suspected fracture. If the splint will not extend in such a manner, reassess possible involvement of the pelvis
 - Extend the distal end of the splint at least 6 inches beyond the foot.
 - Attach the ankle device to the traction crank.
 - Twist until moderate resistance is met.
 - Reassess alignment, pulses, sensation, and motor function. If there has been deterioration in any of these 3 parameters, release traction and reassess.
8. Document the time, type of splint, and the pre and post assessment of pulse, sensation, and motor function in the patient care report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Wound Care-General

Clinical Indications:

- Protection and care for open wounds prior to and during transport.

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Procedure:

1. Use personal protective equipment, including gloves, gown, and mask as indicated.
2. If active bleeding, elevate the affected area if possible and hold direct pressure. Do not rely on “compression” bandage to control bleeding. Direct pressure is much more effective.
3. Once bleeding is controlled, irrigate contaminated wounds with saline as appropriate (this may have to be avoided if bleeding was difficult to control). Consider analgesia per protocol prior to irrigation.
4. Cover wounds with sterile gauze/dressings. Check distal pulses, sensation, and motor function to ensure the bandage is not too tight.
5. Monitor wounds and/or dressings throughout transport for bleeding.
6. Document the wound and assessment and care in the patient care report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Wound Care-Hemostatic Agent

Clinical Indications:

- Serious hemorrhage that can not be controlled by other means.

Contraindications:

- Wounds involving open thoracic or abdominal cavities.

Procedure:

1. Apply approved non-heat-generating hemostatic agent per manufacturer's instructions.
2. Supplement with direct pressure and standard hemorrhage control techniques.
3. Apply dressing.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P



Wound Care-Conducted Electrical Weapon Removal

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Patient with uncomplicated conducted electrical weapon probes embedded subcutaneously in non-sensitive areas of skin.
- Conducted electrical weapon probes are barbed metal projectiles that may embed themselves up to 13 mm into the skin.

Contraindications:

- Patients with conducted electrical weapon probe penetration in vulnerable areas of body as mentioned below should be transported for further evaluation and probe removal
- Probes embedded in skin above level of clavicles, female breasts, or genitalia
- Suspicion that probe might be embedded in bone, blood vessel, or other sensitive structure.

Procedure:

- Ensure wires are disconnected from weapon.
- Stabilize skin around probe using non-dominant hand.
- Grasp probe by metal body with pliers or hemostats to prevent puncture wounds to EMS personnel.
- Remove probe in single quick motion.
- Wipe wound with antiseptic wipe and apply dressing.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Wound Care-Tourniquet

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Life threatening extremity hemorrhage that can not be controlled by other means.
- Serious or life threatening extremity hemorrhage and tactical considerations prevent the use of standard hemorrhage control techniques.

Contraindications:

- Non-extremity hemorrhage
- Proximal extremity location where tourniquet application is not practical

Procedure:

1. Place tourniquet proximal to wound.
2. Route the band around the limb, pass the red tip through the slit of the buckle, and position the tourniquet above the bleeding site directly top the skin OR Insert the limb through the loop in the band and position the tourniquet above the bleeding site directly to the skin.
3. Pull band tightly and fasten it back on itself all the way around the limb, but not over the rod clips. Band should be tight enough that tips of three (3) fingers cannot be slid between the band and the limb. If the tips of three (3) fingers slide under band, re tighten and re-secure.
4. Twist the rod until bleeding has stopped.
5. Secure the rod inside a clip to lock it in place. Check for bleeding and distal pulse. If bleeding is not controlled, or distal pulse is present, consider additional tightening or applying a second tourniquet above and side-by-side to the first. Reassess.
6. Route the band between the clips and over the rod. Secure rod and band with TIME strap.
7. Note time of tourniquet application and communicate this to receiving care providers.
8. Dress wounds per standard wound care protocol. Minimal dressing should be placed over the wound so that inspection of the wound for continued bleeding can be performed.
9. If delayed or prolonged transport and tourniquet application time > 45 minutes: consider reattempting standard hemorrhage control techniques and removing tourniquet. Contact Medical Control.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Trauma Hand- Off Tool

Clinical Indications:

Used upon transfer of care in the ED to a Trauma Team to facilitate a rapid and standardized communication during the time -out period.

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Procedure:

- Mechanism** - simple description of mechanism. GSW with location, Stab with location, Fall with height, MVC, etc. While the mechanism of injury is important and increases index of suspicion of injury it will not impact the immediate actions of the trauma surgeon. Keep it simple! The intimate details of the mechanism are important but not in the first 30 seconds of acute resuscitation. These details are used for data collection and secondary treatment decisions. Place them in your PCR.
- Known / Suspected Injuries** - List these in a head-to-toe fashion. Be sure to clearly notate any suspected open fractures or uncontrolled bleeding. Start with what you know (i.e. open fracture, location of bullet hole) and progress to what you think (i.e. TBI, Pelvis fracture)
- Highest Heart Rate** - Provide the highest heart rate obtained during the patient encounter.
- Lowest BP** - Provide the lowest systolic pressure recorded during the patient encounter. This is used universally as a key indicator for shock in the trauma patient. Drops in blood pressure below 90 systolic at any time substantially increase the chances of death due to shock in the trauma patient.
*NOTE- Trauma surgeons use the shock index (divide the highest heart rate by the lowest systolic BP) to predict the severity of illness in the acute trauma patient.
- Best GCS** - provide the best GCS score recorded during the patient encounter. This will provide the surgeon with perfusion status and noted improvement or decline.
- Lines, Fluids, Interventions** - provide number, size and location of IV's, amount and type of fluid administered, any medications administered, and any other interventions performed.

The intent of this hand off tool is to streamline the communication process. The information the pre-hospital provider has pertaining to the patient is a very important component of the patient's recovery and all the information needed in the first moments of acute trauma resuscitation during that critical time of passing information verbally and passing off the patient physically. It is the hope that this tool will provide for that moment of focused silence in the trauma bay allowing the pre-hospital providers the time needed to articulate this vital information and the trauma surgeon the time to process it and make decisions. This process's goal is to increase chances of survival for the patient.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Pelvic Binder

Clinical Indications:

- Any suspected pelvic fracture or injury
- Pelvic pain with trauma mechanism
- Unconscious with trauma mechanism
- Signs of shock with trauma mechanism


	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Contraindications:


When there is an impaled object that would be covered by the binder.

Procedure:


1. Remove objects from patient's pocket or pelvic area. Place SAM Pelvic Sling II gray side up beneath patient at level of trochanters (hips).



2. Place BLACK STRAP through buckle and pull completely through.



3. Hold ORANGE STRAP and pull BLACK STRAP in opposite direction until you hear and feel the buckle click. Maintain tension and immediately press BLACK STRAP onto surface of SAM Pelvic Sling II to secure.



Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Topical Application of TXA

A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Patients with uncontrolled hemorrhage due to avulsed tooth or epistaxis.

Procedure:

1. Don personal protective equipment (gloves, eye protection, etc.).
2. Soak one 4x4 in 1gram of TXA.
3. Dental: fashion gauze into small square and place into socket with patient closing teeth to exert pressure.
Epistaxis: After failed attempt at use of Oxymetazoline and Direct Pressure. Have the patient blow their nose to remove any remaining clots. Spray Oxymetazoline twice in the affected nostril. Insert the TXA soaked gauze into the affected nostril straight back with forceps.
4. Check for bleeding by visualizing the posterior pharynx for blood.
5. Gently remove the gauze after 20 minutes.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Low Titer O+ Whole Blood Administration

P PARAMEDIC P

Clinical Indications:

- For patients in hemorrhagic shock, and at least one of the following signs:
 - Systolic BP <70 mmHg
 - Systolic BP <90 mmHg AND HR >110 BPM
 - Witnessed traumatic arrest <10 min PTA and continuous CPR throughout downtime
 - Age >65: Systolic BP <100 AND HR >100 BPM
 - Shock Index Criteria >1 (SI = HR/SBP)
 - For pediatric patients <10 years see PM4 for criteria

Contraindications:

- Religious objection to receiving whole blood - Consult On-line Medical Control

Procedure:

1. Don personal protective equipment (gloves, eye protection, etc.).
2. Assess or initiate venous access with IV/IO greater than 22g diameter.
3. Visually inspect the blood product to ensure that there is no evidence of clots, discoloration or leakage of the unit.
4. Connect Normal Saline and Blood bag to the infusion set.
5. Open Normal Saline bag to flow to prepare to prime the tubing.
6. Connect Life Flow tubing to the Compact Disposable Unit (CDU) where it is labeled N.
7. Prime the air check chamber by squeezing it until you have removed all air from the lower and upper chambers and it is filled all the way to the top of the chamber with NS.
8. Load the syringe into the Life Flow Plus with the numbers facing up.
9. Prime the tubing. Point the handle up and squeeze the handle about six times until the CDU is primed as well.
10. Insert the CDU in the top of the Qin Flow Fluid Warmer with arrows facing the front and press down. Lights will turn green when it is ready.
11. Turn off the NS and turn on the blood.
12. Begin administering the blood by squeezing the handle of the Life Flow Plus.
13. Repeat complete vital signs every 10 minutes to include temperature.
14. **Remove a single label sticker off the blood product and place on the PCR make a copy and leave at the hospital. Upload the original into the ePCR.**
15. If entire unit of blood is adm., flush the tubing with NS to ensure all product has been given.
16. During administration, monitor the patient for Transmission Associated Circulatory Overload (TACO) and/or transfusion reaction. TACO s/s are pulmonary edema. If allergic reaction occurs exit to protocol AM1 or PM1. Discontinue administration of blood product and notify receiving facility.
 - a. Do not discard tubing or unit of blood. Disconnect from the patient and place in bio-hazard bag. Turn over to hospital for testing.
17. Blood should be administered until a return of a strong radial pulse OR a patient has a SBP of 90 or SBP of 110 if age >65.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System. Assessment should include direct observation once per certification cycle.



Refusal Form Instructions

Instructions

The EMS Patient Refusal Form has been designed to be used by EMS personnel to legally document a variety of situations. The front of the page is used to describe the situation and the back lists a variety of specific patient instructions by complaint.

The form should be used to document any refusal of care by a patient (complete refusal or refusal of specific aspects of care) and to document the patient / guardian's understanding of medical instructions.

To understand the intent of this form, it is probably simplest to walk through several common patient encounter situations.

1. Complete refusal of EMS care or transport: In the first section, the appropriate blocks for "paramedic recommendation" should be marked. This section should be explained to the patient or guardian, who should understand that their refusal may result in complications up to and including death. The patient or guardian should be asked to sign the form, indicating that he/she understands the seriousness of the situation and the information provided. If the situation warrants, the paramedic should explain the risks of the refusal using the patient instructions section. If the instructions section is used, the appropriate blocks should also be checked.
2. Refusal of a specific procedure (IV therapy, for example): In the first section, the appropriate blocks for "paramedic recommendation" should be marked. The first section should be explained to the patient or guardian, who should understand the potential consequences of their refusal. The patient or guardian should be asked to sign the form, indicating that he/she understands the seriousness of the situation.
3. The appropriate blocks in the instructions given section should be marked. This section and the specific instructions should all be carefully explained to the patient and/or guardian, who must understand them. The patient or guardian should be asked to sign the form, indicating that he/she understands the instructions and the seriousness of the situation.

In all situations, the top part of the form should be completed, and as much of the signature portion as necessary. It is preferable to have witnesses, particularly if the patient or guardian refuses to sign.

Discharge Instructions

UNIVERSAL INSTRUCTIONS:

- YOU HAVE NOT RECEIVED A COMPLETE MEDICAL EVALUATION. SEE A PHYSICIAN AS SOON AS POSSIBLE.
- IF AT ANY TIME AFTER YOU HAVE TAKEN ANY MEDICATION, YOU HAVE TROUBLE BREATHING, START WHEEZING, GET HIVES OR A RASH, OR HAVE ANY UNEXPECTED REACTION, CALL 911 IMMEDIATELY.
- IF YOUR SYMPTOMS WORSEN AT ANY TIME, YOU SHOULD SEE YOUR DOCTOR, GO TO THE EMERGENCY DEPARTMENT OR CALL 911.

ABDOMINAL PAIN:

- Abdominal pain is also called belly pain. Many illnesses can cause abdominal pain and it is very difficult for EMS to identify the cause.
- Take your temperature every 4 hours.

Call or see a physician, go to the emergency department, or call 911 immediately if:

- Your pain gets worse or is now only in 1 area
- You vomit (throw up) blood or find blood in your bowel movement
- You become dizzy or faint
- Your abdomen becomes distended or swollen
- You have a temperature over 100° F
- You have trouble passing urine
- You have trouble breathing

BACK PAIN:

- Apply heat to the painful area to help relieve pain. You may use a warm heating pad, whirlpool bath, or warm, moist towels for 10 to 20 minutes every hour.
- Stay in bed as much as possible the first 24 hours.
- Begin normal activities when you can do them without causing pain.
- When picking things up, bend at the hips and knees. Never bend from the waist only.

Call or see a physician, go to the emergency department, or call 911 immediately if:

- You have shooting pains into your buttocks, groin, legs, or arms or the pain increases.
- You have trouble urinating or lose control of your stools or urine.
- You have numbness or weakness in your legs, feet, arms, or hands.

FEVER:

- Always take medications as directed. Tylenol and Ibuprofen can be taken at the same time.
- If you are taking antibiotics, take them until they are gone, not until you are feeling better.
- Drink extra liquids (1 glass of water, soft drink or gatorade per hour of fever for an adult)
- If the temperature is above 103° F, it can be brought down by a sponge bath with room temperature water. Do not use cold water, a fan, or an alcohol bath.
- Temperature should be taken every 4 hours.

Call or see a physician, go to the emergency department, or call 911 immediately if:

- Temperature is greater than 101° F for 24 hours
- A child becomes less active or alert.
- The Temperature does not come down with Acetaminophen (Tylenol) or Ibuprofen with the appropriate dose.

HEAD INJURY:

- Immediately after a blow to the head, nausea, and vomiting may occur.
- Individuals who have sustained a head injury must be checked, and if necessary awakened, every 2 hours for the first 24 hours.
- Ice may be placed on the injured area to decrease pain and swelling.
- Only drink clear liquids such as juices, soft drinks, or water the first 12 hours after injury..
- Acetaminophen (Tylenol) or Ibuprofen only may be used for pain.

Call or see a physician, go to the emergency department, or call 911 immediately if:

- The injured person has persistent vomiting, is not able to be awakened, has trouble walking or using an arm or leg, has a seizure, develops unequal pupils, has a clear or bloody fluid coming from the ears or nose, or has strange behavior.

INSECT BITE/STING:

- A bite or sting typically is a red lump which may have a hole in the center. You may have pain, swelling and a rash. Severe stings may cause a headache and an upset stomach (vomiting).
- Some individuals will have an allergic reaction to a bite or sting. Difficulty breathing or chest pain is an emergency requiring medical care.
- Elevation of the injured area and ice (applied to the area 10 to 20 minutes each hour) will decrease pain and swelling.
- Diphenhydramine (Benadryl) may be used as directed to control itching and hives.

Call or see a physician, go to the emergency department, or call 911 immediately if:

- You develop any chest pain or difficulty breathing.
- The area becomes red, warm, tender, and swollen beyond the area of the bite or sting.
- You develop a temperature above 101° F.

RESPIRATORY DISTRESS:

- Respiratory Distress is also known as shortness of breath or difficulty breathing.
- Causes of Respiratory Distress include reactions to pollen, dust, animals, molds, foods, drugs, infections, smoke, and respiratory conditions such as Asthma and COPD. If possible avoid any causes which produce respiratory distress.
- If you have seen a physician for this problem, take all medication's as directed.

Call or see a physician, go to the emergency department, or call 911 immediately if:

- Temperature is greater than 101° F.
- The cough, wheezing, or breathing difficulty becomes worse or does not improve even when taking medications.
- You have Chest Pain.
- Sputum (spit) changes from clear to yellow, green, grey, or becomes bloody.
- You are not able to perform normal activities.

EXTREMITY INJURY:

- Extremity Injuries may consist of cuts, scrapes, bruises, sprains, or broken bones (fractures).
- Apply ice on the injury for 15 to 20 minutes each hour for the first 1 to 2 days.
- Elevate the extremity above the heart as possible for the first 48 hours to decrease pain and swelling.
- Use the extremity as pain allows.

Call or see a physician, go to the emergency department, or call 911 immediately if:

- Temperature is greater than 101° F.
- The bruising, swelling, or pain gets worse despite the treatment listed above.
- Any problems listed on the **Wound Care instructions** are noted.
- You are unable to move the extremity or if numbness or tingling is noted.
- You are not improved in 24 to 48 hours or you are not normal in 7 to 10 days.

VOMITING/DIARRHEA:

- Vomiting (throwing up) can be caused by many things. It is common in children, but should be watched closely.
- Diarrhea is most often caused by either a food reaction or infection.
- Dehydration is the most serious problem associated with vomiting or diarrhea.
- Drink clear liquids such as water, apple juice, soft drinks, or gatorade for the first 12 hours or until things improve. Adults should drink 8 to 12 glasses of fluids per day with diarrhea. Children should drink 1 cup of fluid for each loose bowel movement.

Call or see a physician, go to the emergency department, or call 911 immediately if:

- Temperature is greater than 101° F.
- Vomiting or Diarrhea lasts longer than 24 hours, gets worse, or blood is noted.
- You cannot keep fluids down or no urination is noted in 8 hours.

WOUND CARE:

- Wounds include cuts, scrapes, bites, abrasions, or puncture wounds.
- If the wound begins to bleed, apply pressure over the wound with a clean bandage and elevate the wound above the heart for 5 to 10 minutes.
- Unless instructed otherwise, clean the wound twice daily with soapy water, and keep the wound dry. It is safe to take a shower but do not place the wound in bath or dish water.
- See a physician for a tetanus shot if it has been 10 years or more since your last one.

Call or see a physician, go to the emergency department, or call 911 immediately if:

- See the **Extremity Injury instructions**.
- Temperature is greater than 101° F.
- Bruising, swelling, or pain gets worse or bleeding is not controlled as directed above.
- Any signs of infection, such as redness, drainage of yellow fluid or pus, red streaks extending from the wound, or a bad smell is noted.



On-Scene Physician Form

This EMS service would like to thank you for your effort and assistance. Please be advised that the EMS Professionals are operating under strict protocols and guidelines established by their medical director and the State of North Carolina. As a licensed physician, you may assume medical care of the patient. In order to do so, you will need to:

1. Receive approval to assume the patient's medical care from the EMS Agencies Online Medical Control physician.
2. Show proper identification including current North Carolina Medical Board Registration/ Licensure.
3. Accompany the patient to the hospital.
4. Carry out any interventions that do not conform to the EMS Agencies Protocols. EMS personnel cannot perform any interventions or administer medications that are not included in their protocols.
5. Sign all orders on the EMS Patient Care Report.
6. Assume all medico-legal responsibility for all patient care activities until the patient's care is transferred to another physician at the destination hospital.
7. Complete the "Assumption of Medical Care" section of this form below.

Assumption of Medical Care

I, _____, MD; License #: _____
(Please Print your Name Here)

have assumed authority and responsibility for the medical care and patient management for

(Insert Patient's Name Here)

I understand that I must accompany the patient to the Emergency Department. I further understand that all EMS personnel must follow North Carolina EMS Rules and Regulations as well as local EMS System protocols.

_____, MD Date: ____/____/____ Time: ____ AM/PM
(Physician Signature Here)

_____, EMS _____ Witness
(EMS Lead Crew Member Signature Here) (Witness Signature Here)



Apgar Score

The Apgar score should be obtained and recorded initially and at 5 minutes with the birth of delivery of any infant.

- Each of the 5 parameters should be scored and then totaled.
- The Minimum score is 0
- The Maximum score is 10

Sign	0	1	2
Heart Rate	Absent	<100 min.	>100 min.
Respiratory Effort	Absent	Weak Cry	Strong Cry
Muscle Tone	Limp	Some Flexion	Good Flexion
Reflex Irritability (when feet stimulated)	No Response	Some Motion	Cry
Color	Blue; Pale	Body Pink Extremities Blue	Pink



Los Angeles Prehospital Stroke Screen (LAPSS)

1. Patient Name: _____
(last name) (first name)

2. Information/History from: Patient Family Member Other

(name - if other than patient) (phone)

3. Last known time patient was at baseline or deficit free and awake:

(military time) (date)

SCREENING CRITERIA

	Yes	Unknown	No
4. Age > 45	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
5. History of seizures or epilepsy absent	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
6. Symptom duration less than 24 hours	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
7. At baseline, patient is not wheelchair bound or bedridden	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
8. Blood glucose between 60 and 400	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
9. Exam: LOOK FOR OBVIOUS ASYMMETRY			
	Normal	Right	Left
Facial smile/grimace	<input type="checkbox"/>	<input type="checkbox"/> Droop	<input type="checkbox"/> Droop
Hand grip	<input type="checkbox"/>	<input type="checkbox"/> Weak	<input type="checkbox"/> Weak
		<input type="checkbox"/> No grip	<input type="checkbox"/> No grip
Arm strength	<input type="checkbox"/>	<input type="checkbox"/> Drifts dn	<input type="checkbox"/> Drifts dn
		<input type="checkbox"/> Falls fast	<input type="checkbox"/> Falls fast

Based on exam, patient has only unilateral (not bilateral) weakness: YES NO

10. Items 4, 5, 6, 7, 8, 9 all YES's (or unknown) --- LAPSS screening criteria met:

YES NO


11. If LAPSS criteria for stroke are met, alert the receiving hospital of a possible stroke patient. If not, then return to the appropriate treatment protocol.

(Note: the patient may be experiencing a stroke even if the LAPSS criteria are not met.)

12. Time LAPSS Exam Performed: Military Time: _____

13. Form Completed by: _____

Pain Scale Forms




No Pain Moderate Pain Worst Possible Pain

0 1 2 3 4 5 6 7 8 9 10

**If you are having pain, tell your doctor or nurse.
Use these pain scales to describe your pain.**

No Hurt	Hurts Little Bit	Hurts Little More	Hurts Even More	Hurts Whole Lot	Hurts Worst
					

UNC Health Care Pain Management Committee
March 2001



Sin dolor Duele todavía más El peor dolor

0 1 2 3 4 5 6 7 8 9 10

**Si tiene dolor, dígaselo a su doctor o enfermera.
Use esta escala para describir su dolor.**

Sin Dolor	Duele un poquito	Duele un poquito más	Duele todavía más	Duele mucho	El peor dolor
					

UNC Health Care
Pain Management Committee
March 2001

From Hockenberry MJ, Wilson D, Winkelstein ML; Wong's Essentials of Pediatric Nursing, ed. 7, St. Louis, 2005, p. 1259. Used with permission. Copyright, Mosby.



Restraint Checklist

Patient's Name: _____

PCR Number: _____ Date: _____

It is recommended that a Restraint Checklist be completed with any restraint use.

1. Reason for restraint (check all that apply):

- Patient attempting to hurt self
- Patient attempting to hurt others
- Patient attempting to remove medically necessary devices

2. Attempted verbal reassurance / redirection?

- Yes
- No

3. Attempted environmental modification? (i.e. remove patient from stressful environment)

- Yes
- No

4. Received medical control order for restraints?

- Yes _____, MD
- No (Medical Control Physician Name Here)

5. Time and Type of restraint applied (check all that apply):

Date: ____/____/____ Time: ____AM/PM

Limb restraints:

- LUE
- RUE
- LLE
- RLE

Chemical Restraint:

- Yes
- No

If Yes: Drug Used: _____

Total Dose: _____

6. Vital signs and extremity neurovascular exam should be taken every 15 minutes.

7. Transport Position (Patient should NOT be in prone position)

- Supine position for transport
- Lateral recumbent position for transport

Signature: _____

(EMS Lead Crew Member)



Approved Medical Abbreviations

The following is a list of approved medical abbreviations. Providers should use only these abbreviations in medical documentation or other agency specific approved abbreviations.

Most commonly used abbreviations:

A/O x3	Alert and oriented to person, place and time
A/O x4	Alert and oriented to person, place, time and situation
AOSTF	Arrived on scene to find
BBS	Bilateral Breath Sounds
BILAT or B/L	Bilateral
BG or BGL	Blood Glucose
BL	Bilevel Positive Airway Device
BS	Breath Sounds
CC or C/C	Chief Complaint
C/O	Complains Of
CP	Chest Pain
CTA	Clear to Auscultation
DNEB	Duoneb
ET	Endotracheal
ETCO2	End Tidal Carbon Dioxide
ETT	Endotracheal Tube
HI	Homicidal Ideation
LKW	Last Known Well
MAE	Moves All Extremities
PT	Patient
SI	Suicidal Ideation
SZ or Sz	Seizure
T	Temperature
TK or TQ	Tourniquet
TOR	Termination of Resuscitation
VS or V/S	Vital Signs
VSS	Vital Signs Stable

Agency specific hospital/ facilities abbreviations:

NHBMC	Novant Health Brunswick Medical Center
NHNHRMC	Novant Health New Hanover Regional Medical Center
GSRMC	Grand Strand regional Medical Center
MHSC	McLeod Health Seacoast



Approved Medical Abbreviations

The following is a list of approved medical abbreviations. Providers should use only these abbreviations in medical documentation or other agency specific approved abbreviations.

A:

AAA	Abdominal Aortic Aneurysm
AAL	Anterior Axillary Line
ABC	Airway, Breathing, Circulation
ABD	Abdomen or Abdominal
ABCH	Airway, Breathing, Circulation, Hemorrhage
AC	Antecubital fossa
ACLS	Advanced Cardiac Life Support
ACV	Assist-Control Volume
ACP	Assist-Control Pressure
A/O x 3	Alert and oriented to person, place, and time
A/O x 4	Alert and oriented to person, place, time, and situation
AED	Automated External Defibrillator
AEMT	Advanced Emergency Medical Technician
AFIB	Atrial Fibrillation
AFLT	Atrial Flutter
AFIBRVR	Atrial Fibrillation with Rapid Ventricular Response
AFLTRVR	Atrial Flutter with Rapid Ventricular Response
AI	Adrenal Insufficiency
AICD	Automated Internal Cardioverter – Defibrillator
AIDS	Acquired Immunodeficient Syndrome
AIVR	Accelerated Idioventricular Rhythm
AKA	Above Knee Amputation
ALS	Advanced Life Support
AOSTF	Arrived on scene to find
AM	Morning
AMA	Against Medical Advice
AMB	Ambulance
AMI	Anterior Myocardial Infarction
AMS	Altered Mental Status
AMT	Amount
APGAR	Appearance, Pulse, Grimace, Activity, Respiratory
APPROX or ~	Approximately



Approved Medical Abbreviations

The following is a list of approved medical abbreviations. Providers should use only these abbreviations in medical documentation or other agency specific approved abbreviations.

A:

ASA	Aspirin
ASSOC	Associated or Association
AVB	AV Block / Atrioventricular Block
AVPU	Alert, Responsive to Verbal, Responsive to Pain, Unresponsive

B:

BB	Beta-blockers
BBS	Bilateral Breath Sounds
BILAT or B/L	Bilateral
BIAD	Blind Insertion Airway Device
BI-VAD	Bi-Ventricular Assist Device
BG or BGL	Blood Glucose
BKA	Below Knee Amputation
BL	Bilevel Positive Airway Pressure
BLS	Basic Life Support
BM	Bowel Movement
BP	Blood Pressure
BPM	Breaths per Minute or Beats per Minute
BS	Breath Sounds
BSI	Body Substance Isolation
BVM	Bag-Valve-Mask

C:

CA	CANCER
CABG	Coronary Artery Bypass Graft
CAD	Coronary Artery Disease
CATH	Catheter
Cap refill	Capillary refill
CBG	Capillary Blood Glucose
CC or C/C	Chief Complaint



Approved Medical Abbreviations

The following is a list of approved medical abbreviations. Providers should use only these abbreviations in medical documentation or other agency specific approved abbreviations.

C:

CCP	Casualty Collection Point
CCU	Coronary Care Unit
CHF	Congestive Heart Failure
CHI	Closed Head Injury
CKD	Chronic Kidney Disease
CNA	Certified Nursing Assistant
CNS	Central Nervous System
COMM	Command
COMMP	Command Post
CO2	Carbon Dioxide
C/O	Complains Of
COPD	Chronic Obstructive Pulmonary Disease
CO	Carbon Monoxide
CP	Chest Pain
CPAP	Continuous Positive Airway Pressure
CPR	Cardiopulmonary Resuscitation
CVA	Cerebrovascular Accident (Stroke)
CS	Cervical Spine
CSIC	Cervical Spine Immobilization Collar
CSF	Cerebrospinal Fluid
C-SECT	Cesarean Section
CT	Cat Scan
CTA	Clear to Auscultation

D:

D5W	5% Dextrose in Water
D5NS	5% Dextrose in Normal Saline
D5LR	5% Dextrose in Lactate Ringers
D10	10% Dextrose in Water or 10% Dextrose
D50	50% Dextrose



Approved Medical Abbreviations

The following is a list of approved medical abbreviations. Providers should use only these abbreviations in medical documentation or other agency specific approved abbreviations.

D:	
D25	25% Dextrose
DBP	Diastolic Blood Pressure
DC or D/C	Discontinue
DCAPBTLS	Deformities, contusions, abrasions, penetrations, burns, tenderness, lacerations and swelling
Defib	Defibrillation
DKA	Diabetic Ketoacidosis
DL	Direct Laryngoscopy
DNEB	DuoNeb
DNI	Do Not Intubate
DNR	Do Not Resuscitate
DM	Diabetes Mellitus
DOA	Dead on Arrival
DOB	Date of Birth
DOE	Dyspnea on Exertion
DAA	Drug Assisted Airway
DT	Delirium Tremens
DVT	Deep Venous Thrombosis
DX or Dx	Diagnosis

E:	
EBL	Estimated Blood Loss
ECG or EKG	Electrocardiogram
ED	Emergency Department
eFAST	Enhanced Focused Assessment with Sonography in Trauma
EEG	Electroencephalogram
EGA	Extra-glottic Airway
EJ	External Jugular
eKit	Emergency (Hospice) Kit
EQ or =	Equal



Approved Medical Abbreviations

The following is a list of approved medical abbreviations. Providers should use only these abbreviations in medical documentation or other agency specific approved abbreviations.

E:

EMD	Emergency Medical Dispatcher
EMR	Emergency Medical Responder
EMT	Emergency Medical Technician
EOC	Emergency Operations Center
ePCR	Electronic Patient Care Report
ESRD	End Stage Renal Disease
ET	Endotracheal
ETA	Estimated Time of Arrival
ETCO ₂	End-Tidal Carbon Dioxide
ETT	Endotracheal Tube
ETOH	Ethanol or Alcohol
EXT	External or Extension

F:

F or ♀	Female
FAST	Focused Assessment with Sonography in Trauma
FB	Foreign body
FF	Firefighter
FiO ₂	Fraction of Inspired Oxygen Concentration
FR	First Responder
FSBS	Finger Stick Blood Sugar
FLEX	Flexion
FX	Fracture

G:

g	Gram(s)
G	Gravida
gtts	Drops
>	Greater
≥	Greater than or equal to



Approved Medical Abbreviations

The following is a list of approved medical abbreviations. Providers should use only these abbreviations in medical documentation or other agency specific approved abbreviations.

G:

GCS	Glasgow Coma Score
GI	Gastrointestinal
GIB	Gastrointestinal Bleed (ing)
GSW	Gunshot wound
GU	Genitourinary
GYN	Gynecology or Gynecological

H:

HA	Headache
HazMat	Hazardous Material(s)
HCPOA	Health Care Power of Attorney
HEENT	Head, Eyes, Ears, Nose, Throat
HD	Hemodialysis
HI	Homicidal Ideation
HIV	Human Immunodeficiency Virus
HOB	Head of Bed
HOSP	Hospital
HR	Heart rate
HTN	Hypertension
HTX	Hemothorax
HX	History

I:

IABP	Intra-Aortic Balloon Pump
ICP	Intracranial pressure
ICS	Incident Command System
ICU	Intensive care unit
I:E	Inspiratory to Expiratory Ratio
IM	Intramuscular
IN	Intranasal



Approved Medical Abbreviations

The following is a list of approved medical abbreviations. Providers should use only these abbreviations in medical documentation or other agency specific approved abbreviations.

I:

IO	Intraosseous
IU	International Unit
IUD	Intrauterine Device
IV	Intravenous
IVP	Intravenous Push
IVR	Idioventricular Rhythm
IVPB	IV Piggy Back
IOP	Intraosseous Push

J:

J	Joules
JVD	Jugular Vein Distension

K:

KED	Kendrick Extrication Device
Kg or kg	Kilogram
KTD	Kendrick Traction Device
KVO	Keep Vein Open

L:

L	Liter
LAC	Laceration
LBBB	Left Bundle Branch Block
LEO	Law Enforcement Officer
<	Less
≤	Less than or equal to
LKW	Last Known Well
L-SPINE	Lumbar spine
LS	Lumbosacral spine
L&D	Labor and delivery



Approved Medical Abbreviations

The following is a list of approved medical abbreviations. Providers should use only these abbreviations in medical documentation or other agency specific approved abbreviations.

L:

LAT	Lateral
Lbs.	Pound or Pounds
LLE	Left Lower Extremity
LLQ	Left Lower Quadrant
LUE	Left Upper Extremity
LMA	Laryngeal Mask Airway
LMP	Last Menstrual Period
LOC	Level of Consciousness
LPN	Licensed Practical Nurse
LR	Lactated ringers
LS	Lumbar Spine
LSB	Long Spine Board
LSN	Last Seen Normal
LUQ	Left Upper Quadrant
LVAD	Left Ventricular Assist Device

M:

M or ♂	Male
MAE	Moves All Extremities
MAP	Mean Arterial Pressure
MCI	Mass Casualty Incident
MCL	Mid Clavicular Line
mcg	Microgram(s)
MDI	Metered Dose Inhaler
ME	Medical Examiner
MED	Medicine
MERT	Medical Emergency Response Team
mEq	Milliequivalent
Mg or mg	Milligram(s)
mg/dL	milligrams per deciliter



Approved Medical Abbreviations

The following is a list of approved medical abbreviations. Providers should use only these abbreviations in medical documentation or other agency specific approved abbreviations.

M:

MI	Myocardial Infarction (heart attack)
min	Minute
mL	Milliliter
MOI	Mechanism of Injury
MOST	Medical Order for Scope of Treatment
mm	Millimeter
MS	Mental status
MSC	Mental status change
msec	Miliseconds
MV	Mechanical Ventilation
MVC	Motor Vehicle Crash

N:

N/A	Not applicable
N/V	Nausea/Vomiting
N/V/D	Nausea/Vomiting/Diarrhea
NAD	No Apparent (or Acute) Distress
NAEMSP	National Association of EMS Physicians
NC	Nasal Cannula
NCCEP	North Carolina Chapter of Emergency Physicians
NCOEMS	North Carolina Office of EMS
NEB	Nebulizer
NEG or -	Negative
NGT	Nasogastric Tube
NH	Nursing Home
NIPPV	Non-Invasive Positive Pressure Ventilation
NKDA	No Known Drug Allergies
NO	Nitrous Oxide
NPO	Nothing by Mouth
NPA	Nasopharyngeal Airway



Approved Medical Abbreviations

The following is a list of approved medical abbreviations. Providers should use only these abbreviations in medical documentation or other agency specific approved abbreviations.

N:

NRB	Non-rebreather Face Mask
NREMT	National Registry of EMT
NS	Normal Saline
NSAID	Non-steroidal Anti-inflammatory Drug
NSR	Normal Sinus Rhythm
NSTEMI	Non ST-Segment Myocardial Infarction
NTG	Nitroglycerin
NTI	Nasotracheal Intubation

O:

O2	Oxygen
OBGYN	Obstetrics and Gynecology
OHCA	Out-of-Hospital Cardiac Arrest
OD	Overdose
OGT	Orogastric Tube
OME	Oral Morphine Equivalents
OPA	Oropharyngeal Airway
OPO	Organ Procurement Organization
OTI	Orotracheal Intubation
OTC	Over-the-counter (medications)
OZ	Ounces

P:

P	Para
PA	Physician Assistant
PALP	Palpation
PAC	Premature Atrial Contraction
PBW	Predicted Body Weight
PCI	Percutaneous Coronary Intervention
PCP	Primary Care Provider
PCR	Patient Care Report



Approved Medical Abbreviations

The following is a list of approved medical abbreviations. Providers should use only these abbreviations in medical documentation or other agency specific approved abbreviations.

P:	
PD	Peritoneal Dialysis
PE	Pulmonary embolus
PEA	Pulseless Electrical Activity
PEEP	Positive End Expiratory Pressure
PEARL	Pupils equal and reactive to light
PICC	Peripheral Inserted Central Catheter
PIP	Peak Inspiratory Pressure
PJC	Premature Junctional Contraction
PMH	Past Medical History
PM	Evening
PMS	Pulse, Motor, Sensory
PO	Oral or By Mouth
POCUS	Point of Care Ultrasound
POS or +	Positive
POV	Privately Owned Vehicle
PP	Plateau Pressure
PPE	Personal Protective Equipment
PPH	Post partum Hemorrhage
PRN	As needed
PSVT	Paroxysmal Supraventricular Tachycardia
PSY or Ψ	Psychiatric
PT	Patient
PTA	Prior to Arrival
PTX	Pneumothorax
PVC	Premature Ventricular Contraction
Q:	
q	Every
QRV	Quick Response Vehicle
QUES or ?	Question or questionable



Approved Medical Abbreviations

The following is a list of approved medical abbreviations. Providers should use only these abbreviations in medical documentation or other agency specific approved abbreviations.

R:	
RBBB	Right Bundle Branch Block
ROM	Range of Motion
RLE	Right Lower Extremity
RLQ	Right Lower Quadrant
RUE	Right Upper Ex
ROSC	Return of Spontaneous Circulation
RN	Registered Nurse
RR	Respiratory Rate
RSA	Rapid Sequence Airway
RSI	Rapid Sequence Intubation
RT or RCP	Respiratory Therapist or Respiratory Care Provider
RTF	Rescue Task Force
RUQ	Right Upper Quadrant
RVAD	Right Ventricular Assist Device
RXN	Reaction

S:	
SA	Sinus Arrhythmia
SBP	Systolic Blood Pressure
SCBA	Self-Contained Breathing Appartus
S/P	Status Post
sec	Seconds
SGA	Supraglottic Airway
SI	Suicidal Ideation
Six	Shock Index
SIMV	Synchronized Intermittent Mandatory Ventilation
SL	Sublingual
SOB	Shortness of Breath
SQ	Subcutaneous
SNF	Skill Nursing Facility



Approved Medical Abbreviations

The following is a list of approved medical abbreviations. Providers should use only these abbreviations in medical documentation or other agency specific approved abbreviations.

S:

SOG	Standard Operating Guideline(s)
SOP	Standard Operating Procedure(s)
SPO2	Pulse Oximetry
SMR	Spinal Motion Restriction
SSB	Short Spine Board
SSN	Social Security Number
SSS	Sick Sinus Syndrome
ST	Sinus Tachycardia
STEMI	ST Segment Elevation Myocardial Infarction
SVD	Spontaneous Vaginal Delivery
SVT	Supraventricular Tachycardia
SWAT	Special Weapons and Tactics Team
SX or Sx	Symptom(s)
Sync	Synchronized
SZ or Sz	Seizure

T:

T	Temperature
TAH	Total Artificial Heart
TBSA	Total Body Surface Area
TCP	Transcutaneous Pacing
TFCPR	Team Focused Cardiopulmonary Resuscitation
TIA	Transient Ischemic Attack
TK or TQ	Tourniquet
TKO	To Keep Open
TOB	Time of Birth
TOD	Time of Death
TOI	Time of Ingestion
TOO	Time of Onset
TOR	Termination of Resuscitation



Approved Medical Abbreviations

The following is a list of approved medical abbreviations. Providers should use only these abbreviations in medical documentation or other agency specific approved abbreviations.

T:

tPA	Tissue Plasminogen Activator
T-POD	Trauma Pelvic Orthotic Device
TS	Thoracic Spine
TTP	Targeted Temperature Management
TV	Tidal Volume
TX or Tx	Treatment
TXA	Tranexamic Acid

U:

UOA	Upon Our Arrival
URI	Upper Respiratory Infection
US	Ultrasound
USIV	Ultrasound IV
UTI	Urinary Tract Infection

V:

VAD	Ventricular Assist Device
VF	Ventricular Fibrillation
VL	Video Laryngoscopy
VS or V/S	Vital Signs
VSS	Vital Signs Stable
VT	Ventricular Tachycardia

W:

WAP	Wandering Atrial Pacemaker
WCD	Wearable Cardio-Defibrillator Vest
WMD	Weapon of Mass Destruction
WNL	Within Normal Limits
WPW	Wolf Parkinson White syndrome
W/S	Watts per Second



Approved Medical Abbreviations

The following is a list of approved medical abbreviations. Providers should use only these abbreviations in medical documentation or other agency specific approved abbreviations.

W:

WT or Wt Weight

Y:

Y/O Year(s) Old

YOF Year Old Female

YOM Year Old Male

YONB Year Old Nonbinary

Z:

ZED Zee Extrication Device

Symbol Chart:

~	Approximate, approximately, approximation
&	And
@	At
Δ	Change
°	Degree (s)
=	Equal (s)
♀	Female
>	Greater than
≥	Greater than or equal to
<	Less than
≤	Less than or equal to
♂	Male
-	Negative
#	Number
%	Percent, percentage
+	Positive
ψ	Psychiatry, psychiatric
?	Question, questionable



Difficult Airway Evaluation

Evaluating for the difficult airway

Between 1 – 3% of patients who require endotracheal intubation have airways that make intubation difficult. Recognizing those patients who may have a difficult airway allows the paramedic to proceed with caution and to keep as many options open as possible. It also allows the paramedic to prepare additional equipment (such as a cricothyrotomy kit) that may not ordinarily be part of a standard airway kit. The mnemonic LEMON is useful in evaluating patients for signs that may be consistent with a difficult airway and should raise the paramedic's index of suspicion.

Look externally


External indicators of either difficult intubation or difficult ventilation include: presence of a beard or moustache, abnormal facial shape, extreme cachexia, edentulous mouth, facial trauma, obesity, large front teeth or "buck teeth", high arching palate, receding mandible, short bull neck.

Evaluate 3-3-2 Rule

- 3 fingers between the patient's teeth (patient's mouth should open adequately to permit three fingers to be placed between the upper and lower teeth)
- 3 fingers between the tip of the jaw and the beginning of the neck (under the chin)
- 2 fingers between the thyroid notch and the floor of the mandible (top of the neck)

Mallampati

This scoring system is based on the work of Mallampati et al published in the Canadian Anaesthesia Society Journal in 1985. The system takes into account the anatomy of the mouth and the view of various anatomical structures when the patient opens his mouth as wide as possible. This test is performed with the patient in the sitting position, the head held in a neutral position, the mouth wide open, and the tongue protruding to the maximum. Inappropriate scoring may occur if the patient is in the supine position (instead of sitting), if the patient phonates or if the patient arches his or her tongue.



Class I (easy) = visualization of the soft palate, fauces, uvula, anterior and posterior pillars.

Class II = visualization of the soft palate, fauces and uvula.

Class III = visualization of the soft palate and the base of the uvula.

Class IV (difficult) = soft palate is not visible at all.

Obstruction?

Besides the obvious difficulty if the airway is obstructed with a foreign body, the paramedic should also consider other obstructers such as tumor, abscess, epiglottitis, or expanding hematoma.

Neck Mobility

Ask the patient to place their chin on their chest and to tilt their head backward as far as possible. Obviously, this will not be possible in the immobilized trauma patient.



Burns Resources

Fluid Formula

Formula for Fluid Resuscitation of the Burn Patient (Also known as the Parkland Formula)

Pts Wt kg x %TBSA x 4.0cc LR infused over 24 hours with half given in the first 8 hours.

(For the equation, the abbreviations are: PW x TBSA x 4.0 cc)

EMS focuses on the care given during the 1st hour or several hours following the event. Thus the formula as adapted for EMS and the first 8 hours is:

$$PW \times TBSA \times 4.0 \text{ cc, divide by 2}$$

to take this to the hourly rate, divide that solution by 8 and the equation becomes:

$$PW \times TBSA \times 4.0 \text{ cc} / 2 / 8 = \text{total to be infused for each of the first 8 hours.}$$

Another way to state the equation is to use:

$$PW \times TBSA \times 0.25 \text{ cc} = \text{total to be infused for each hour of the first 8 hours.}$$

Example. 80 kg patient with 50 %TBSA x 0.25 cc = 1000 cc/hr.


Remember:

Patient's Weight in kg (2.2 lbs = 1.0 kg) example: 220 lbs adult = 100 kg

% TBSA = Rule of Nine Total Body Surface Area

Factor for the 1st hr. and each hr. for the 1st 8 hrs. = 0.25

(Reminder, if two IV's are running, divide total amount to be infused each hr. by 2)



Estimate spotty areas by using the size of the patient's palm as 1%.

Wt (kg)	% TBSA	Factor	/Hr for 1st 8 Hrs of Care	60 gtt set, gtt/min	20 gtt set, gtt/min	15 gtt set, gtt/min	10 gtt set, gtt/min
10	10	0.25	25	25	8.3	6.3	4.2
10	20	0.25	50	50	16.7	12.5	8.3
10	30	0.25	75	75	25.0	18.8	12.5
10	40	0.25	100	100	33.3	25.0	16.7
10	50	0.25	125	125	41.7	31.3	20.8
20	10	0.25	50	50	16.7	12.5	8.3
20	20	0.25	100	100	33.3	25.0	16.7
20	30	0.25	150	150	50.0	37.5	25.0
20	40	0.25	200	200	66.7	50.0	33.3
20	50	0.25	250	250	83.3	62.5	41.7
30	10	0.25	75	75	25.0	18.8	12.5
30	20	0.25	150	150	50.0	37.5	25.0
30	30	0.25	225	225	75.0	56.3	37.5
30	40	0.25	300	300	100.0	75.0	50.0
30	50	0.25	375	375	125.0	93.8	62.5
40	10	0.25	100	100	33.3	25.0	16.7
40	20	0.25	200	200	66.7	50.0	33.3
40	30	0.25	300	300	100.0	75.0	50.0
40	40	0.25	400	400	133.3	100.0	66.7
40	50	0.25	500	500	166.7	125.0	83.3
50	10	0.25	125	125	41.7	31.3	20.8
50	20	0.25	250	250	83.3	62.5	41.7
50	30	0.25	375	375	125.0	93.8	62.5
50	40	0.25	500	500	166.7	125.0	83.3
50	50	0.25	625	625	208.3	156.3	104.2
60	10	0.25	150	150	50.0	37.5	25.0
60	20	0.25	300	300	100.0	75.0	50.0
60	30	0.25	450	450	150.0	112.5	75.0
60	40	0.25	600	600	200.0	150.0	100.0
60	50	0.25	750	750	250.0	187.5	125.0
70	10	0.25	175	175	58.3	43.8	29.2
70	20	0.25	350	350	116.7	87.5	58.3
70	30	0.25	525	525	175.0	131.3	87.5
70	40	0.25	700	700	233.3	175.0	116.7
70	50	0.25	875	875	291.7	218.8	145.8
80	10	0.25	200	200	66.7	50.0	33.3
80	20	0.25	400	400	133.3	100.0	66.7
80	30	0.25	600	600	200.0	150.0	100.0
80	40	0.25	800	800	266.7	200.0	133.3
80	50	0.25	1000	1000	333.3	250.0	166.7
90	10	0.25	225	225	75.0	56.3	37.5
90	20	0.25	450	450	150.0	112.5	75.0
90	30	0.25	675	675	225.0	168.8	112.5
90	40	0.25	900	900	300.0	225.0	150.0
90	50	0.25	1125	1125	375.0	281.3	187.5
100	10	0.25	250	250	83.3	62.5	41.7
100	20	0.25	500	500	166.7	125.0	83.3
100	30	0.25	750	750	250.0	187.5	125.0
100	40	0.25	1000	1000	333.3	250.0	166.7
100	50	0.25	1250	1250	416.7	312.5	208.3

Critical (Red)

>15% TBSA 2nd/3rd Degree Burn
Burns with Multiple Trauma
Burns with definitive airway compromise
(When reasonable accessible, transport to a Burn Center)

Serious (Yellow)

5-15% TBSA 2nd/3rd Degree Burn
Suspected Inhalation injury or requiring intubation for airway stabilization
Hypotension
GCS < 14
(When reasonable accessible, transport to either a Level I Burn Center or a Trauma Center)

Minor (Green)

< 5% TBSA 2nd/3rd Degree Burn
No inhalation injury, Not Intubated, Normotensive
GCS > 14
(Transport to the Local Hospital)

Drug Drip Charts

Amiodarone

- **AC6 Adult Tachycardia – Narrow**
- **AC7 Adult Monomorphic Tachycardia – Wide**
- **AC8 Adult Polymorphic Tachycardia – Wide**

Initial Dose: 150mg IV/IO infused over 10 minutes

Mix Amiodarone 150mg in 100cc bag D5W; use 10gtt set and infuse over 10 minutes.

Recurrent Tachycardia:

Mix 450mg in 250cc bag D5W; use 10gtt set and administer 6gtts/min

Diltiazem (Cardizem)

- **AC6 Adult Tachycardia – Narrow**

Initial Dose: 0.25 mg/kg IV/IO infused over 2-3 minutes; Max 25 mg

1. Pull tab on Cardizem vial
2. Pull tab on 100cc bag NS
3. Screw Cardizem vial into NS bag
4. Pull inner plug/stopper on NS bag
5. Mix well
6. Pull port cover and insert clave
7. Draw up dose of Cardizem by calculation. This may take 2-3 10cc syringes.
8. Give total dose over 2-3 minutes not to exceed 25 mg

Recurrent Tachycardia: 0.35 mg/kg w/ Max 25mg

Drug Drip Charts

Epinephrine- Adult

- **AC2 Bradycardia**
- **AM5 Hypotension / Shock**

Dose: 1-10 mcg/min IV/IO, titrate to SBP \geq 90

Mix 1mg of Epi 1:1 concentration into 500cc bag of NS; use 10gtt set	
Concentration = 1 mcg per ½ cc	
5gtts = ½ cc = 1 mcg/min	30gtts = 3cc = 6 mcg/min
10gtts = 1cc = 2 mcg/min	40gtts = 4cc = 8 mcg/min
20gtts = 2cc = 4 mcg/min	50gtts = 5cc = 10 mcg/min
Push Dose Epi, draw up 5cc from mixed bag = 10mcg/min	

- ❖ Start at 10mcg/min and titrate down once desired effect is obtained, may need maintenance therapies or repeat doses.

Epinephrine- Pediatric

- **PM1 Pediatric Allergic reaction**
- **PM3 Pediatric Hypotension/Shock**

Dose: 0.1 mcg/kg/min IV/IO, titrate to SBP \geq 70 + 2(age)

Mix 1mg Epi 1:1 concentration into 500cc bag NS; use a 10gtt set	
Concentration = 1cc = 2 mcg	
Example: 10kg child	
$0.1 \text{ mcg/kg} \times 10\text{kg} = 1 \text{ mcg} = 0.5\text{cc} = 5\text{gtts/ min}$	
Example: 20kg child	
$0.1 \text{ mcg/kg} \times 20 \text{ kg} = 2 \text{ mcg} = 1\text{cc} = 10 \text{ gtts/ min}$	

Drug Drip Charts

Levetiracetam (Keppra)

- **UP13 Seizure**
- **AR8 Post Intubation**

Dose: 1g IV/IO

Mix 1 gram in 100cc bag of NS, use 60gtt set

Infuse over 15 minutes

Magnesium Sulfate

- **AR4 Adult COPD / Asthma Respiratory Distress**

Dose: 2g IV/IO

Mix 2 grams in 100cc bag NS

Use 10gtt set

Infuse over 10-20 minutes

- **AO3 OB Emergency Seizure**

Mix 2 grams in 100cc bag NS Infuse
over 2-3 minutes

Metoprolol

- **AC6 Adult Tachycardia – Narrow**

Mix 5 mg in 100cc bag NS

Use 10gtt set

Infuse over 5 minutes

Drug Drip Charts

Norepinephrine (Levophed)

- **AM5 Hypotension/Shock- Adult ONLY**

Dose: 0.01 – 1.0 mcg/kg/min IV/IO via pump ONLY

Mix 4mg in 250cc bag of D5W

Enter the weight in kg

Enter mcg you wish to start at (start at 1mcg/kg/min and titrate down once you have reached SBP goal of ≥ 90)

Enter the volume of 250cc

Procainamide

- **AC7 Adult Monomorphic Tachycardia – Wide**

Dose 20-50 mg/min IV/IO

Maximum = 17 mg/kg

Mix 1000 mg in 100cc bag NS; use 10gtt set
20gtts = 2cc = 20mg
30gtts = 3cc = 30mg
40gtts = 4cc = 40mg
50gtts = 5cc = 50mg

Drug Drip Charts

TXA

- **A03 Obstetrical Emergencies**

Criteria: postpartum hemorrhage where birth occurred <3 hours

Dose: 1-2 grams

Mix 1-2 grams in 100mL NS

Infuse over 10 minutes

- **TB6 Multiple Trauma**

Criteria: Adult- SBP \leq 70 **OR** SBP \leq 90 and HR > 110

Age > 65- SBP < 100 and HR > 100

Dose:

Adult
1-2 grams in 100mL NS IV/IO
Mix 1-2 grams in 100mL NS
Infuse over 10 minutes

Drug List

ONLY medications that are included by name and dose in the 2025 NCCEP Protocols are included in this document; the only purpose of this document is to serve as a reference. For a full list of medications approved for use by EMS professionals, please refer to the NC Medical Board document titled: Approved Medications for Credentialed EMS Personnel.

Medication	Adult Dosing	Pediatric Dosing
<p><u>Acetaminophen</u> Analgesic, Antipyretic NCCEP Protocol: *UP10- Fever / Infection Control *UP11- Pain Control <u>Indications / Contraindications:</u></p> <ul style="list-style-type: none"> • Analgesic and antipyretic • Patients with liver failure should not receive 	<ul style="list-style-type: none"> • 325 – 1000 mg PO • Up to 1000 mg IV/IO infused over 15 minutes 	<ul style="list-style-type: none"> • 15 mg/kg PO / IV • Maximum single dose 750 mg • IV/IO dose should be infused over 15 minutes
<p><u>Activated Charcoal</u> NCCEP Protocol: *TE7 Overdose / Toxic Ingestion <u>Indications</u> Used to treat oral poisonings following excessive oral ingestion ≤ 1 Hour</p>	<ul style="list-style-type: none"> • 1 gm/kg PO • Max dose 50 gm 	<ul style="list-style-type: none"> • 1 gm/kg PO • Max dose 50 gm
<p><u>Adenosine</u> (Adenocard) Antidysrhythmic NCCEP Protocol: *AC6- Adult Tachycardia Narrow *AC7- Adult Monomorphic Tachycardia Wide *PC5- Pediatric Tachycardia Narrow <u>Indications / Contraindications:</u></p> <ul style="list-style-type: none"> • Specifically for treatment or diagnosis of Supraventricular Tachycardia • Use caution in patients with WPW 	<ul style="list-style-type: none"> • 6 mg IV/IO rapid push with flush • May repeat 12 mg IV/IO • May repeat 12 mg IV/IO 	<ul style="list-style-type: none"> • 0.1 mg/kg IV/IO Maximum 6 mg • May repeat 0.2 mg/kg IV/IO Maximum 12 mg • Use stopcock or both ports and a NS flush with each dose

Drug List

Medication	Adult Dosing	Pediatric Dosing
<p><u>Albuterol</u> Beta- Agonist</p> <p>NCCEP Protocol: *AM1- Allergic Reaction / Anaphylaxis *TB3- Crush Syndrome Trauma *PM1- Pediatric Allergic Reaction *AR4- Adult COPD / Asthma</p> <p><u>Indications / Contraindications:</u></p> <ul style="list-style-type: none"> For use in respiratory distress with bronchospasm 	<ul style="list-style-type: none"> 2.5 – 5 mg in nebulizer May repeat 3 doses 	<ul style="list-style-type: none"> 2.5 – 5 mg in nebulizer May repeat 3 doses
<p><u>Amiodarone</u> Antiarrhythmic NCCEP Protocol: *AC6- Adult Tachycardia Narrow *AC7- Adult Monomorphic Tachycardia Wide Complex *AC8- Adult Polymorphic Tachycardia Wide Torsade's *AC9- V-Fib / Pulseless V-Tach *PC6- Pediatric Tachycardia Wide Complex</p> <p><u>Indications / Contraindications</u></p> <ul style="list-style-type: none"> Antiarrhythmic agent used to treat ventricular arrhythmias and atrial tachycardias Avoid with hypersensitivity to iodine, cardiogenic shock and high degree AV Blocks 	<ul style="list-style-type: none"> 150 mg in 100mL of D5W IV/IO infused over 10 minutes May repeat if tachycardia recurs or persists May also be drawn up in syringe (not preferred), 150 mg in 10 cc flush SLOW IV push over 10 minutes <p style="text-align: center;"><u>V-Fib / Pulseless V-Tach</u></p> <ul style="list-style-type: none"> 300 mg IV/IO may repeat if refractory @ 150 mg IV/IO 	<p><u>Wide Complex Tachycardia (>= 0.09)</u></p> <ul style="list-style-type: none"> 5 mg/kg over 20-60 minutes Consult med control recommended
<p><u>Aspirin</u> NCCEP Protocol: *UP11- Pain Control *AC4- Chest Pain: Cardiac and STEMI</p> <p><u>Indications / Contraindications:</u></p> <ul style="list-style-type: none"> An antiplatelet drug for use in cardiac chest pain <p>Contraindicated in pediatrics less than 15 years of age</p>	<ul style="list-style-type: none"> 324 to 650 mg PO for pain control 81 mg x4 PO (chewed) OR 325 mg PO 	

Drug List

Medication	Adult Dosing	Pediatric Dosing
<p><u>Atropine</u> Anticholinergic NCCEP Protocol: *AC2- Bradycardia; Pulse Present *PC2- Pediatric Bradycardia w/ Poor Perfusion *TE8- WMD-Nerve Agent <u>Indications / Contraindications:</u></p> <ul style="list-style-type: none"> • Anticholinergic drug used in bradycardia • Ineffective and potentially harmful in cardiac transplantation 	<ul style="list-style-type: none"> • 0.5 mg IV/IO • May repeat every 3-5 minutes • Max dose 3mg <p><u>Organophosphate Poisoning</u></p> <ul style="list-style-type: none"> • 2 mg IV/IO/IM • Repeat every 3-5 minutes until symptoms resolve 	<ul style="list-style-type: none"> • 0.02 mg/kg IV/IO • May repeat once • Minimum single dose 0.1 mg • Maximum single dose 0.5 mg <p><u>Organophosphate Poisoning</u></p> <ul style="list-style-type: none"> • Weight based • <= 40 lbs.- 0.5 mg • 40- 90 lbs.- 1 mg • >90 lbs.- 2 mg
<p><u>Calcium Gluconate</u> Electrolyte NCCEP Protocol: *AM6 – Blood Adm. *AM3- Dialysis / Renal failure *PM 4 – Blood Adm. *TB3- Crush Syndrome Trauma *TE7- Overdose / Toxic Ingestion <u>Indications / Contraindications:</u></p> <ul style="list-style-type: none"> • Indicated for severe hyperkalemia • Sodium Bicarb and Calcium should not be mixed, ideally given in separate lines. • Administer with whole blood in separate IV line. 	<ul style="list-style-type: none"> • 2gm IV/IO over 2-3 minutes 	<ul style="list-style-type: none"> • 30 mg/kg IV/IO over 10 minutes
<p><u>Dexamethasone (Decadron)</u> Corticosteroid NCCEP Protocol: *AM1-Allergic Reaction/ Anaphylaxis *PM1- Pediatric Allergic reaction *AR4- Adult COPD/ Asthma *AR7- Pediatric Asthma <u>Indications / Contraindications:</u></p> <ul style="list-style-type: none"> • Indicated Anaphylaxis, COPD, Asthma, Croup • Caution: diabetes/hyperglycemia 	<ul style="list-style-type: none"> • 4 mg IV / IO/ IM 	<ul style="list-style-type: none"> • 0.15 mg/kg IV / IO / IM

Drug List

Medication	Adult Dosing	Pediatric Dosing
<p>Dextrose Carbohydrate NCCEP Protocol: *AM2- Diabetic; Adult *PM2- Pediatric Diabetic Indications / Contraindications: Use in AMS or hypoglycemic states</p>	<ul style="list-style-type: none"> • D10 125mL-250mL IV/IO • D50 12.5-25gm IV/IO May repeat until blood glucose is greater than or equal to 80mg/dL • D50 may be given rectally; contact medical control for advice 	<ul style="list-style-type: none"> • < 1 y/o- D10: 5mL/kg IV/IO Repeat as needed • 1-2 years- D25: 2mL/kg IV/IO Repeat as needed • >2 years- D50: 1mL/kg IV/IO Maximum 25g per dose Repeat as needed
<p>Diazepam (Valium) Benzodiazepine NCCEP Protocol: *UP13- Seizure UP18 Behavioral/ Agitation *UP19 Behavioral/ Violent *AC7- Adult Monomorphic Tachycardia *AO3- Obstetrical Emergencies *PC5- Pediatric Tachycardia Wide Complex *TB3- Crush Syndrome Trauma *TE1- Bites and Envenomation's *TE4- Hyperthermia Indications / Contraindications: • seizures and sedation</p>	<ul style="list-style-type: none"> • 5 mg IV / IO • 10 mg PR 	<ul style="list-style-type: none"> • 0.1 mg/kg IV / IO -Max 4 mg • 0.5 mg/kg PR -Max 10 mg
<p>Diltiazem (Cardizem) Calcium Channel Blocker NCCEP Protocol: *AC6- Adult Tachycardia Narrow Indications / Contraindications: • Calcium channel blocker used to treat narrow complex SVT. • Contraindicated in patients with heart block, bradycardia, V-Tach, WPW, and/or acute MI</p>	<ul style="list-style-type: none"> • 0.25mg/kg IV/IO slow IV push over 2-3 minutes • Max dose 25mg • May repeat in 15 minutes if no improvement with 0.35mg/kg slow IV/IO push over 2-3 minutes 	
<p>Diphenhydramine (Benadryl) Antihistamine NCCEP Protocol: *UP17- Behavioral Health Crisis *AM1 Allergic Reaction / Anaphylaxis *PM1- Pediatric allergic Reaction Indications / Contraindications: • allergic reaction • Treatment for extrapyramidal reactions</p>	<ul style="list-style-type: none"> • 25-50mg IV/IM/IO/PO 	<ul style="list-style-type: none"> • 1mg/kg IV/IM/IO/PO • Do not give to infants less than 3 months old

Drug List

Medication	Adult Dosing	Pediatric Dosing
<p>Dopamine NCCEP Protocol: *AC2- Bradycardia; Pulse Present *AM5- Hypotension / Shock <u>Indications / Contraindications:</u> Vasopressor used in shock or hypotensive states</p>	<ul style="list-style-type: none"> • 5-20mcg/kg/min IV/IO • Titrate to SBP > or equal 90 	
<p>Epinephrine 1:1000 Sympathomimetic (Bronchodilator) NCCEP Protocol: *AR4- Adult COPD / Asthma Respiratory Distress *AR7- Pedi Asthma Resp Distress *AC2- Bradycardia; Pulse Present *AM1- Allergic Reaction/Anaphylaxis *AM5- Hypotension *PC1- Pediatric Asystole / PEA *PC2- Pedi Brady w/ Poor Perfusion *PC6- Pedi V Fib Pulseless V tach *PM1- Pediatric Allergic Reaction <u>Indications / Contraindications:</u> Vasopressor used in allergic reactions, hypotension, bradycardia and respiratory distress</p>	<ul style="list-style-type: none"> • Allergic Reaction /COPD / Asthma 0.3-0.5 mg IM • Bradycardia / Hypotension 1-10 mcg / min Mix 1 mg into 500cc bag NS Use 10gtt set 10gtt / min = 2 mcg / min 	<ul style="list-style-type: none"> • Asthma / Respiratory Distress 0.01mg / kg IM Max 0.3 mg • Pediatric Asystole / PEA 0.1 mg / kg ETT w/ Max 2.5mg Repeat every 3-5 minutes • Pediatric V Fib Pulseless V Tach 0.1mg/kg ETT w/ max 2.5mg Repeat every 3-5 minutes • Pediatric Allergic Reaction <30 kg- 0.15 mg IM ≥30 kg- 0.3-0.5 mg IM • Nebulized 1mg in 2 mL NS May repeat x 1
<p>Epinephrine 1: 10,000 Sympathomimetic (Cardiac Stimulant) NCCEP Protocol: *AC1- Adult Asystole / PEA *AC9- V Fib / Pulseless V tach *AO2- Newly Born *PC1- Pediatric Asystole / PEA *PC2- Pedi Brady w/ Poor Perfusion *PC6- Pedi V Fib Pulseless V Tach <u>Indications / Contraindications:</u> Vasopressor used in cardiac arrest</p>	<ul style="list-style-type: none"> • 1mg IV/IO Repeat every 3-5 minutes as needed 	<ul style="list-style-type: none"> • 0.01mg/kg IV/IO • Max single dose 1mg Repeat every 3-5 minutes as needed

Drug List

Medication	Adult Dosing	Pediatric Dosing
<p>Etomidate (Amidate) Induction Agent for RSI NCCEP Protocol: *AR3- Airway, Drug Assisted</p> <p><u>Indications / Contraindications</u> Contraindicated in presence of Hypoxia(<90%) and Hypotension (Systolic <90)</p>	<ul style="list-style-type: none"> • 0.3 mg/kg IV/IO • May repeat x 1 	
<p>Famotidine (Pepcid) Histamine-2 Blocker NCCEP Protocol: *AM1- Allergic Reaction /Anaphylaxis *PM1- Pediatric Allergic reaction</p> <p><u>Indications / Contraindications:</u> Histamine-2 blocker, decreases amount of acid produced by stomach</p>	<ul style="list-style-type: none"> • 20 mg PO • Do not allow the patient to chew, swallow with water 	<ul style="list-style-type: none"> • 20 mg PO ONLY age > 12 y/o
<p>Fentanyl (Sublimaze) Narcotic Analgesic NCCEP Protocol: *UP11- Pain Control *AC4- Chest Pain: Cardiac and STEMI *AR7- Post Intubation / BIAD Mgmt. *PC3- Pedi Pulmonary Edema/CHF</p> <p><u>Indications/Contraindications</u> *Narcotic pain relief. *Contraindications to opioids include severe COPD and respiratory distress, monitor closely.</p>	<ul style="list-style-type: none"> • Pain Control 50-100 mcg IV/IO/IN Repeat 25 mcg every 20 minutes Max dose 200 mcg • Chest Pain 50-75 mcg IV/IO Repeat 25 mcg every 20 minutes Max dose 200 mcg • Post Intubation sedation / Pain 50-75 mcg IV/IO Repeat every 5 minutes Max dose 300 mcg 	<ul style="list-style-type: none"> • 1mcg/kg IV/IO • Max single dose 50 mcg/kg • Requires consultation by Medical Control
<p>Furosemide (Lasix) Loop Diuretic NCCEP Protocol: *AC5- CHF / Pulmonary Edema</p> <p><u>Indications / Contraindications:</u></p> <ul style="list-style-type: none"> • Used in acute decompensated heart failure <p>Contraindicated in hypotensive patients</p>	<ul style="list-style-type: none"> • 40 mg IV ONLY IF: <ol style="list-style-type: none"> 1. Transport Time >30 minutes 2. Known CHF / Daily Lasix 3. Afebrile • BP >100 	

Drug List

Medication	Adult Dosing	Pediatric Dosing
<p>Glucagon Hormone NCCEP Protocol: *UP4- Altered Mental Status *AM2- Diabetic; Adult *PM2- Pediatric Diabetic *TE7- Overdose / Toxic Ingestion</p> <p>Indications / Contraindications:</p> <ul style="list-style-type: none"> • Drug acting to release glucose into blood stream by glycogen breakdown. • Use when no IV access is achievable. • Beta-Blocker or Calcium Channel Blocker overdose 	<ul style="list-style-type: none"> • Hypoglycemia 1-2 mg IM Repeat in 15 minutes if needed • Beta Blocker or calcium Channel Blocker Overdose 2-4 mg IV/IO/IM May repeat in 15 minutes if needed 	<ul style="list-style-type: none"> • Hypoglycemia 0.1mg/kg IM Max dose 1 mg Repeat every 15 minutes if needed to keep BGL >
<p>Glucose (Oral) Carbohydrate *AM2- Diabetic; Adult *PM2- Pediatric Diabetic</p> <p>Indications / Contraindications:</p> <ul style="list-style-type: none"> • Use in conscious hypoglycemic states. • Do not administer to patients that are not able to swallow or protect their own airway 	<ul style="list-style-type: none"> • 1 tube (15 grams) or packet • Repeat based on BGL 	<ul style="list-style-type: none"> • ½ to 1 tube (7.5 - 15 grams) if age appropriate • Repeat based on BGL
<p>Haloperidol (Haldol) Antipsychotic NCCEP Protocol: *UP17- Behavioral Health Crisis *UP18- Behavioral Agitation / Sedation Guide</p> <p>Indications / Contraindications:</p> <ul style="list-style-type: none"> • Antipsychotic used to assist in the sedation of aggressive or agitated patients 	<ul style="list-style-type: none"> • Age 12 – 64 years 2-5 mg IM • Age ≥ 65 2.5 mg IM • May repeat every 5 minutes as needed up to 10 MG 	<ul style="list-style-type: none"> • Age ≥ 12 ONLY 1 – 5 mg IM • Age < 12 <p style="text-align: center; font-size: 2em;">∅</p>
<p>Heparin Anticoagulant NCCEP Protocol: *AC4- STEMI</p> <p>Indications:</p> <ul style="list-style-type: none"> • Used in STEMI PT's to block antithrombin III <p>Contraindication: Stroke patients</p>	<ul style="list-style-type: none"> • 60 units/kg • Max dose 10,000 units 	<p style="text-align: center; font-size: 2em;">∅</p>

Drug List

Medication	Adult Dosing	Pediatric Dosing
<p>Ibuprofen (Motrin) NSAID NCCEP Protocol: *UP10- Fever / Infection Control *UP11- Pain Control</p> <p>Indications / Contraindications:</p> <ul style="list-style-type: none"> Avoid NSAIDS in patients who are pregnant or could be pregnant Not to be used in patients with GI bleeding or renal insufficiency Not to be used in patients with allergies to ASA or other NSAID drugs 	<ul style="list-style-type: none"> 400 – 600 mg PO Max 800 mg PO 	<ul style="list-style-type: none"> 10 mg / kg PO if age > 6 months
<p>Ipratropium (Atrovent) Bronchodilator NCCEP Protocol: *AR4- Adult COPD / Asthma *AR7- Pedi Asthma Resp Distress *AM1- Allergic Reaction/Anaphylaxis *PM1- Pediatric Allergic Reaction</p> <p>Indications / Contraindications:</p> <ul style="list-style-type: none"> Given with albuterol to dry secretions and manage bronchospasms 	<ul style="list-style-type: none"> 0.5 mg May repeat as needed x 3 There is no proven benefit to continual use of Ipratropium 	<ul style="list-style-type: none"> 0.5 mg May repeat as needed x 3 There is no proven benefit to continual use of Ipratropium
<p>Ketamine Dissociative anesthetic Induction Agent for RSI</p> <p>NCCEP Protocol: *UP11- Pain Control *UP19-Behavioral *AR3- Airway, Drug Assisted *AR8- Post Intubation / BIAD Mgmt. <i>(only when DAI protocol has been used)</i></p> <p>Indications / Contraindications:</p> <ul style="list-style-type: none"> Contraindications include patients who have cardiac disease and uncontrolled hypertension Avoid use in combination with benzodiazepines 	<ul style="list-style-type: none"> Pain Control 0.3 mg/kg IV/IO over 10 minutes Repeat every 20 minutes Max single dose 30 mg OR 1 mg/kg IN one time Airway Drug Assisted and Management 2mg/kg IV/IO or 4mg/kg IM May repeat x 1, Max 400mg Airway Management + Dangerously Combative 4mg/kg (Max 400mg)IM OR 2 mg/kg IV/IO Post Intubation/ BIAD Mgmt. 1.5 – 2.0 mg/kg IV/IO 	<ul style="list-style-type: none"> Direct Order ONLY from Dr. Sherrod Protocol AR3 when hypoxic, hypotensive or dangerously combative 2 mg/kg IM Max dose 400 mg Can not be used for RSI procedure

Drug List

Medication	Adult Dosing	Pediatric Dosing
<p><u>Ketorolac</u> (Toradol) NSAID</p> <p>NCCEP Protocol: *UP 11- Pain Control</p> <p><u>Indications / Contraindications:</u></p> <ul style="list-style-type: none"> • Avoid in women who are pregnant or could be pregnant • Not to be used in patients with GI bleeding, renal insufficiency, or patients who may need immediate surgery. • Avoid in patients currently taking anticoagulants such as coumadin. 	<ul style="list-style-type: none"> • 15 mg IV / IO • 30 mg IM • Max 30 mg 	<ul style="list-style-type: none"> • 0.5 mg/kg IV / IO / IM
<p><u>Lactated Ringer's Solution</u> Isotonic crystalloid solution NCCEP Protocol: *AC4- Chest Pain: Cardiac and STEMI *TB1- Blast Injury / Incident *TB9- Thermal Burn</p> <p><u>Indications / Contraindications:</u> Preferred fluid for pre-cath hydration and fluid resuscitation in burn care and Sepsis</p>	<ul style="list-style-type: none"> • Chest Pain 250-500 mL pre-Cath hydration • Fluid Resuscitation in Burns 0.25 mL / kg (x % TBSA) / hr. 	<ul style="list-style-type: none"> • Fluid Resuscitation in Burns 0.25 mL / kg (x % TBSA) / hr.
<p><u>Levetiracetam</u> (Keppra) Anticonvulsant NCCEP Protocol: *UP13 – Seizure *AR8 -Post Intubation</p>	<ul style="list-style-type: none"> • 1 gram in 500mL NS over 15 minutes IV/IO 	<ul style="list-style-type: none"> • 15 mg/kg over 15 minutes IV/IO Max dose 1 gram
<p><u>Lidocaine</u> Antidysrhythmic NCCEP Protocol: *AC8- Adult Polymorphic Tachycardia (Wide \geq 0.12 sec) Torsade's de Pointes *AC9- V Fib Pulseless V Tach *PC6- Pediatric V Fib Pulseless V Tach</p> <p><u>Indications / Contraindications:</u></p> <ul style="list-style-type: none"> • Antiarrhythmic used for control of ventricular dysrhythmias 	<ul style="list-style-type: none"> • 1-1.5 mg/kg IV/IO • May repeat if refractory 0.75 mg/kg IV/IO • Max 3 mg/kg 	<ul style="list-style-type: none"> • 1 mg / kg IV/ IO (first bolus) • Infusion 20-50 mcg /kg/min (second dose) <p>(If infusion is initiated > 15 minutes after first bolus, repeat 1 mg/kg bolus)</p>

Drug List

Medication	Adult Dosing	Pediatric Dosing
<p><u>Lorazepam (Ativan)</u> Benzodiazepine NCCEP Protocol: *UP13- Seizure *UP18 Behavioral/ Agitation *UP19 Behavioral/ Violent *AC7- Adult Monomorphic Tachycardia *AO3- Obstetrical Emergencies *PC5- Pediatric Tachycardia Wide Complex *TB3- Crush Syndrome Trauma *TE1- Bites and Envenomation's *TE4- Hyperthermia</p> <p><u>Indications / Contraindications:</u></p> <ul style="list-style-type: none"> • seizures and sedation • IM injection is not as effective in termination of seizures. 	<ul style="list-style-type: none"> • 1mg IV / IO May repeat every 10 minutes as needed up to 4 mg. • 2 mg IM May repeat every 10 minutes as needed up to 4 mg. 	<ul style="list-style-type: none"> • 0.1 mg / kg IV / IO /IM • May repeat every 10 minutes as needed up to 4 mg. • Maximum single dose 1 mg
<p><u>Magnesium Sulfate</u> Electrolyte NCCEP Protocol: *AR4- Adult COPD / Asthma Respiratory Distress *AR7- Pedi. Asthma Resp. Distress *AC8- Adult Polymorphic Tachycardia Wide (≥ 0.12 sec) Torsade's *AC9- V Fib Pulseless V Tach *AO3- Obstetrical Emergency</p> <p><u>Indications / Contraindications:</u></p> <ul style="list-style-type: none"> • Smooth muscle relaxer used in refractory respiratory distress resistant to beta-agonists • Elemental electrolyte used to treat eclampsia during the third trimester of pregnancy. • May assist with treatment of Torsades, prolonged QT, low magnesium states (malnourished/ alcoholic) and suspected digitalis toxicity. 	<ul style="list-style-type: none"> • Respiratory Distress 2g IV / IO infused over 10 – 20 min • Torsade's de Pointes 2g IV / IO May repeat Max 4g • Refractory V Fib Pulseless V tach 2g IV / IO • Obstetrical Emergency Seizure 2-4 g IV/ IO over 2-3 minutes OR 5-10 g IM May repeat x 1 	<ul style="list-style-type: none"> • Respiratory Distress 40 mg / kg IV / IO infuse over 10 – 20 minutes
<p><u>Methylprednisolone</u>(corticosteroid) *AR4- Adult COPD/Asthma *AR7- Pediatric Asthma Respiratory *AM 1/PM1- Allergic Reaction</p>	<ul style="list-style-type: none"> • 125 mg IV/ IO / IM 	<ul style="list-style-type: none"> • 2 mg/kg IV / IO / IM • Max dose 125 mg

Drug List

Medication	Adult Dosing	Pediatric Dosing
<p><u>Metoprolol</u> (Lopressor) Beta-Blocker NCCEP Protocol: *AC6- Adult Tachycardia Narrow</p> <p><u>Indications / Contraindications:</u></p> <ul style="list-style-type: none"> Used to slow ventricular response in A-fib w/ RVR Contraindicated in AV Blocks or w/ BP < 100 systolic 	<ul style="list-style-type: none"> 5 mg administered over 5 minutes IV/IO May repeat 2x at 5-minute intervals Max 15 mg 	
<p><u>Midazolam</u> (Versed) Benzodiazepine NCCEP Protocol: *UP13- Seizure *UP18 Behavioral/ Agitation *UP19 Behavioral/ Violent *AC7- Adult Monomorphic Tachycardia *AO3- Obstetrical Emergencies *PC5- Pediatric Tachycardia Wide Complex *TB3- Crush Syndrome Trauma *TE1- Bites and Envenomation's *TE4- Hyperthermia</p> <p><u>Indications / Contraindications:</u></p> <ul style="list-style-type: none"> Quick acting Benzo used for seizures and sedation IM injection is effective in termination of seizures. IM preferred over IO. 	<ul style="list-style-type: none"> Seizure 2-2.5 mg IV/ IO May repeat every 3 to 5 minutes as needed Maximum 10 mg <u>NO IV</u> ≥ 49 kg- 10 mg IM, may repeat in 5 minutes, Max 20mg <49 kg- 0.2 mg/kg IM, max single dose 5 mg, may repeat in 5 minutes if needed, Max 10 mg Sedation 2.5 mg IV/IO/ IN May repeat as needed 5 - 10 mg IM Crush Syndrome 0.5mg -2mg IV/IO 1-2 mg IN Max 5 mg Muscle Spasm 2-2.5mg IV / IO over 2-3 minutes Maximum 10mg 	<ul style="list-style-type: none"> Seizure 0.2mg/kg IV / IO Max single dose 2.5 mg May repeat every 3 to 5 minutes as needed Maximum 10 mg <u>NO IV</u> 0.2mg /kg IM with a max of 5mg Sedation 0.1-0.2 mg/kg IV / IO / IN Max single dose 2 mg Max total dose 5 mg Crush Syndrome 0.1-0.2 mg/kg IV / IO / IN Muscle Spasm 0.1-0.2mg/kg IV / IO / IM / IN over 2-3 minutes Maximum 5 mg IM
<p><u>Morphine Sulfate</u> Opioid Narcotic Analgesic NCCEP Protocol: *UP11- Pain Control *AC4- Chest Pain *PC3- Pedi Pulmonary Edema / CHF</p> <p><u>Indications / Contraindications:</u></p> <ul style="list-style-type: none"> Narcotic Pain Relief Avoid if BP < 110 	<ul style="list-style-type: none"> Pain 4mg IV / IO / IM Repeat 2 mg every 5 minutes as needed Maximum 10 mg Chest Pain 2-4mg IV / IO Repeat every 5 minutes as needed Maximum 10 mg 	<ul style="list-style-type: none"> Pain 0.1mg/kg IV / IO / IM May repeat every 5 minutes Maximum 10 mg Pediatric Pulmonary Edema Requires Med Control Order

Drug List

Medication	Adult Dosing	Pediatric Dosing
<p><u>Naloxone</u> (Narcan) Opioid Narcotic Antagonist</p> <p>NCCEP Protocol: *AC3- Cardiac Arrest; Adult *PC1- Pediatric Asystole / PEA *PC4- Pediatric Cardiac Arrest *TE7- Overdose / Toxic Ingestion</p> <p><u>Indications / Contraindications:</u></p> <ul style="list-style-type: none"> • Titrate to adequate respirations and oxygenation. • NOT GIVEN TO RESTORE CONSCIOUSNESS 	<ul style="list-style-type: none"> • 0.4-2mg IV / IO / IN / IM / ETT Repeat as needed 	<ul style="list-style-type: none"> • 0.1 mg / kg IV / IO / IN / IM Repeat as needed
<p><u>Nitroglycerin</u> Vasodilator, Nitrate NCCEP Protocol: *AC4- Chest Pain: Cardiac and STEMI *AC5- CHF / Pulmonary Edema *PC3- Pedi Pulmonary Edema /CHF</p> <p><u>Indications / Contraindications</u></p> <ul style="list-style-type: none"> • Avoid in any patient who has used Viagra or Levitra in the past 24 hours or Cialis in the past 36 hours 	<ul style="list-style-type: none"> • 0.3-0.4mg SL Repeat every 5 minutes as needed • Nitroglycerin Paste SBP > 100 1 inch SBP > 150 1.5 inch SBP > 200 2 inches 	<ul style="list-style-type: none"> • Pediatric Pulmonary Edema Requires Med Control Order
<p><u>Norepinephrine</u> (Levophed) Vasoconstrictor NCCEP Protocols: *AC2 *AM5</p> <p><u>Indications/Contraindications</u></p> <ul style="list-style-type: none"> • Hypotension 	<ul style="list-style-type: none"> • 0.1 – 1.0 mcg/kg/min IV /IO via Pump Only 	
<p><u>Normal Saline</u> Crystalloid Solution NCCEP Protocol: *Multiple</p> <p>Special Consideration in: *UP7- Dental Problems *AC4- Chest Pain: Cardiac and STEMI *AM3- Dialysis Renal Failure *AO2- Newly Born *TB2- Chemical and Electrical Burn *TB4- Extremity Trauma *TB7- Radiation Incident *TB9- Thermal Burn</p>	<ul style="list-style-type: none"> • Chest Pain; Cardiac and STEMI Consider bolus of 250 -500 mL for pre-Cath hydration • Thermal Burn 0.25mL/kg (x % TBSA) / hr. for up to first 8 hours • See individual protocols for dosing and/or bolus rates 	<ul style="list-style-type: none"> • 20 mL / kg IV / IO Repeat as needed x 3 Max dose 60 ml / kg • Newly Born HR < 60 10 mL/ kg IV / IO • See individual protocols for dosing and/or bolus rates

Drug List

Medication	Adult Dosing	Pediatric Dosing
<p><u>Ondansetron</u> (Zofran) Antiemetic NCCEP Protocol: *UP3- Abdominal Pain Vomiting and Diarrhea <u>Indications / Contraindications:</u></p> <ul style="list-style-type: none"> Zofran is the recommended antiemetic for EMS use since it has significantly less side effects and sedation. 	<ul style="list-style-type: none"> 4mg IV / IO / IM / PO / ODT May repeat in 15 minutes 	<ul style="list-style-type: none"> 0.2 mg/kg PO / ODT Maximum 4 mg May repeat in 15 minutes
<p><u>Oxygen</u> <u>Indications / Contraindications:</u></p> <ul style="list-style-type: none"> Indicated in any condition with increased cardiac workload, respiratory distress, or illness or injury resulting in altered ventilation and/or perfusion. Goal oxygen saturation 94-98% Indicated for pre-oxygenation whenever possible prior to endotracheal intubation. Goal oxygen saturation 100%. 	<ul style="list-style-type: none"> 1-6 liters/minute via nasal cannula 10-15 liters/minute via NRB mask 15 liters or > via BVM/ETT/BIAD 	<ul style="list-style-type: none"> 1-6 liters/minute via nasal cannula 10-15 liters/minute via NRB mask 15 liters or > via BVM/ETT/BIAD
<p><u>Oxymetazoline (Afrin)</u> Nasal Decongestant Spray NCCEP Protocol: *UP9- Epistaxis <u>Indications / Contraindications</u> Vasoconstrictor used in nasal intubation and epistaxis</p>	<ul style="list-style-type: none"> 2 sprays in affected nostril followed by direct pressure 	<ul style="list-style-type: none"> 2 sprays in affected nostril followed by direct pressure
<p><u>Oxytocin (Pitocin)</u> Hormone NCCEP Protocol: *AO1- Childbirth/Labor *AO3- OB-GYN Emergency <u>Indications / Contraindications</u></p> <ul style="list-style-type: none"> promote uterine contractions ensure no other fetus is present 	<ul style="list-style-type: none"> 10 IU IM 	
<p><u>Pralidoxime (2-PAM)</u> NCCEP Protocol: WMD – Nerve Agent Protocol - *TE8 <u>Indications/Contraindications</u></p> <ul style="list-style-type: none"> Antidote for nerve agents or organophosphate overdose Administered with Atropine 	<ul style="list-style-type: none"> 600 mg IV / IO / IM over 30 minutes 	<ul style="list-style-type: none"> 15-25 mg / kg IV / IO / IM over 30 minutes

Drug List

Medication	Adult Dosing	Pediatric Dosing
<p>Procainamide (antiarrhythmic) *AC7- Adult Monomorphic Tachycardia Wide Complex</p>	<ul style="list-style-type: none"> • 20-50 mg / min IV / IO • 1-4 mg / min maintenance drip <p>Maximum 17 mg / kg</p>	
<p>Promethazine (Phenergan) Antiemetic NCCEP Protocol: *UP3- Abdominal Pain Vomiting and Diarrhea Indications / Contraindications:</p> <ul style="list-style-type: none"> • IV Phenergan should be given IV only with great caution. Extravasation of this drug can result in severe damage. • IV-Dilute with 100ml NS and administer slowly over 20 minutes. • Zofran is the preferred prehospital anti-emetic. 	<p style="color: red;">** Can only be administered after no response of Zofran and actively vomiting.</p> <ul style="list-style-type: none"> • 12.5 mg IV / IM • IM is preferred • IV- Dilute w/ 100ml NS, adm. over 20 minutes • May repeat x 1 as needed • Consider lower initial dose of 6.25 mg 	
<p>Rocephin (Ceftriaxone) Antibiotic NCCEP Protocol: *UP15- Suspected Sepsis *TB4- Extremity Trauma Indications / Contraindications:</p> <ul style="list-style-type: none"> • Administered when sepsis is suspected, and patient meets SIRS criteria • Open fracture or amputated part with bone fracture <p>Contraindicated if pregnant</p>	<ul style="list-style-type: none"> • 50 mg/kg IV / IO up to 2 grams over 10 minutes • 50 mg/kg <p>Sepsis: Can be administered if unable to draw blood cultures ONLY if unable due to no IV access or no culture vials.</p>	
<p>Rocuronium Neuromuscular Blocker / Paralytic NCCEP Protocol: *AR3- Airway, Drug Assisted Indications / Contraindications:</p> <p>*Non-depolarizing paralytic agent used as a component of drug assisted intubation when succinylcholine is contraindicated.</p> <p>*Onset of action is longer than succinylcholine, up to 3 minutes. Patient will not defasciculate.</p>	<ul style="list-style-type: none"> • 1.5 mg/ kg IV / IO • May repeat x 1 	

Drug List

Medication	Adult Dosing	Pediatric Dosing
<p><u>Sodium Bicarbonate</u> Alkalinizing Agent NCCEP Protocol: *AC2- Bradycardia; Pulse Present *AM3- Dialysis / Renal Failure *UP12- Police Custody *TB3- Crush Syndrome *TE7- Overdose/ Toxic Ingestion <u>Indications / Contraindications:</u></p> <ul style="list-style-type: none"> A buffer used in acidosis to increase the pH in cardiac arrest, hyperkalemia, or tricyclic overdose. <p>Sodium Bicarb and Calcium Chloride should not be mixed. Ideally give in separate lines.</p>	<ul style="list-style-type: none"> 50mEq IV/IO Repeat in 10 minutes as needed See individual protocol for specific dosing algorithm 	<ul style="list-style-type: none"> 1mEq/Kg IV/IO Repeat in 10 minutes as needed Max 50mEq See individual protocol for specific dosing algorithm
<p><u>Suxamethonium</u> (Succinylcholine) Neuromuscular Blocker / Paralytic NCCEP Protocol: *AR3- Airway, Drug Assisted <u>Indications / Contraindications:</u></p> <ul style="list-style-type: none"> Paralytic agent used as a component to Drug Assisted Intubation Avoid in patients with burns >24 hours old, chronic neuromuscular disease (found in Native American Lumbee Tribe), underlying hyperkalemia (dialysis patients), patients with rhabdomyolysis. 	<ul style="list-style-type: none"> 2 mg / kg IV / IO May repeat x 1 	

Drug List

Medication	Adult Dosing	Pediatric Dosing
<p><u>Terbutaline</u> B2 Adrenergic Receptor Agonist NCCEP Protocol: *AM1- Allergic Reaction / Anaphylaxis *TB3- Crush Syndrome Trauma *PM1- Pediatric Allergic Reaction *AR4- Adult COPD / Asthma <u>Indications / Contraindications:</u></p> <ul style="list-style-type: none"> For use in respiratory distress with bronchospasm Used in place of Albuterol and Atrovent during COVID Pandemic 	<ul style="list-style-type: none"> 0.25 mg subcutaneous in lateral deltoid May repeat as needed every 15 minutes 	<ul style="list-style-type: none"> 0.01 mg / kg subcutaneously Max dose 0.25 mg May repeat as needed every 15 minutes
<p><u>Tranexamic Acid</u> <u>(TXA)</u>Antifibrinolytic NCCEP Protocol: *UP7 Dental Problems *UP9 Epistaxis *AO3 Obstetrical Emergencies *TB6 Multiple Trauma Must meet criteria</p> <p><u>Indications / Contraindications:</u></p> <ul style="list-style-type: none"> Contraindicated in trauma that occurred >3 hours from EMS arrival. Contraindicated pregnancy and postpartum hemorrhage where birth occurred >3 hours prior to EMS arrival. 	<p><u>OB Emergencies / Multiple Trauma</u></p> <ul style="list-style-type: none"> 1-2 gram mixed in 100 mL NaCl infused over 10 minutes <p><u>Dental Problems / Epistaxis</u></p> <ul style="list-style-type: none"> 1 gram soaked in 1-4x4 gauze for 20 minutes 	

Drug List

Medication	Adult Dosing	Pediatric Dosing
<p><u>Whole Blood Low Titer O+</u></p> <p>NCCEP Protocol: *AM6-Blood Administration *PM4-Pediatric Blood Admin *AO3 – OB /GYN *TB6-Multiple Trauma *TB10- Traumatic Arrest</p> <p><u>Indications:</u> Hypovolemic/Hemorrhagic Shock Trauma Arrest</p> <p><u>Contraindications:</u> Religious Beliefs</p>	<ul style="list-style-type: none"> • 1 Unit IV/IO 	<ul style="list-style-type: none"> • 10 cc/kg IV/IO

These formularies are provided as a reference only. It does not contain all the contraindications and potential side effects or adverse reactions for each listed drug. It is the responsibility of each EMS professional to become and remain knowledgeable about each drug in this formulary.

Updated 9/30/2025